Full wwPDB X-ray Structure Validation Report (i)

Feb 19, 2016 – 10:02 PM GMT

PDB ID	:	5CAM
Title	:	Crystal Structure of the Cytoplasmic Domain of the Pseudomonas putida Anti-sigma Factor PupR (SeMet)
Authors	:	Jensen, J.L.; Colbert, C.L.
Deposited on	:	2015-06-29
Resolution	:	2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5130 (2.20-2.16)
Clashscore	102246	5965 (2.20-2.16)
Ramachandran outliers	100387	5863 (2.20-2.16)
Sidechain outliers	100360	5864 (2.20-2.16)
RSRZ outliers	91569	5142 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	86	9% 56% 7% • 36%
1	B	86	7% 56% 6% • 37%
1	C	86	2% 62% 6% • 31%
1	D	86	% 64% • • 31%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3796 atoms, of which 1794 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PupR protein.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S	Se			
1	A	55	904	290	439	91	79	1	4	85	3	0
1	B	54	878	284	424	87	78	1	4	62	2	0
1	C	59	944	304	458	95	83	1	3	16	1	0
1	D	59	974	314	473	100	83	1	3	6	3	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q52209
A	-2	ALA	-	expression tag	UNP Q52209
A	-1	MSE	-	expression tag	UNP Q52209
A	0	GLY	-	expression tag	UNP Q52209
B	-3	GLY	-	expression tag	UNP Q52209
B	-2	ALA	-	expression tag	UNP Q52209
B	-1	MSE	-	expression tag	UNP Q52209
B	0	GLY	-	expression tag	UNP Q52209
C	-3	GLY	-	expression tag	UNP Q52209
C	-2	ALA	-	expression tag	UNP Q52209
C	-1	MSE	-	expression tag	UNP Q52209
C	0	GLY	-	expression tag	UNP Q52209
D	-3	GLY	-	expression tag	UNP Q52209
D	-2	ALA	-	expression tag	UNP Q52209
D	-1	MSE	-	expression tag	UNP Q52209
D	0	GLY	-	expression tag	UNP Q52209

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	7	7	7	0	0
			Total	O		
2	B	7	7	7	0	0
			Total	O		
2	C	46	46	46	0	0
			Total	O		
2	D	36	36	36	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PupR protein

Chain A:



- Molecule 1: PupR protein

Chain B:



- Molecule 1: PupR protein

Chain C:



- Molecule 1: PupR protein

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	33.49Å 134.69Å 34.90Å 90.00° 113.78° 90.00°	Depositor
Resolution (Å)	67.35 – 2.17 67.35 – 2.17	Depositor EDS
% Data completeness (in resolution range)	94.8 (67.35-2.17) 94.7 (67.35-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.78 (at 2.16Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.178 , 0.216 0.180 , 0.218	Depositor DCC
R_{free} test set	1434 reflections (10.12%)	DCC
Wilson B-factor (Å^2)	34.0	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.41 , 58.5	EDS
Estimated twinning fraction	0.048 for l,-k,h	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.47$, $< L^2 > = 0.30$	Xtriage
Outliers	1 of 14503 reflections (0.007%)	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	3796	wwPDB-VP
Average B, all atoms (Å^2)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.01% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.71	0/484	0.64	0/650
1	B	0.63	0/467	0.69	0/628
1	C	0.95	0/501	0.76	0/676
1	D	0.90	0/523	0.81	0/705
All	All	0.81	0/1975	0.73	0/2659

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	465	439	439	3	0
1	B	454	424	430	8	0
1	C	486	458	464	6	0
1	D	501	473	484	3	0
2	A	7	0	0	0	0
2	B	7	0	0	0	0
2	C	46	0	0	1	0
2	D	36	0	0	2	0
All	All	2002	1794	1817	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:53[A]:ARG:NH1	2:D:101:HOH:O	2.07	0.86
1:C:7:THR:O	2:C:101:HOH:O	2.05	0.74
1:C:61:LEU:HA	1:C:64:MSE:HE2	1.86	0.57
1:B:38[B]:MSE:HE2	1:B:41:ARG:NH1	2.28	0.48
1:B:38[B]:MSE:HE3	1:C:60:ARG:CD	2.44	0.47
1:B:38[B]:MSE:HE2	1:B:41:ARG:CZ	2.46	0.46
1:B:38[B]:MSE:CE	1:B:41:ARG:CZ	2.94	0.45
1:D:21[B]:HIS:CE1	1:D:55:GLN:HG2	2.51	0.45
1:B:34:LEU:HD13	1:C:61:LEU:HD21	1.99	0.45
1:A:44:HIS:ND1	1:A:45:PRO:HD2	2.32	0.44
1:B:38[A]:MSE:HG3	1:C:64:MSE:HE1	1.99	0.44
1:D:65:ARG:NH2	2:D:104:HOH:O	2.51	0.44
1:B:38[B]:MSE:HE3	1:C:60:ARG:HD2	1.99	0.44
1:B:39:SER:O	1:B:42[A]:GLN:NE2	2.42	0.44
1:A:51:TRP:O	1:A:55:GLN:HG2	2.17	0.43
1:A:38[B]:MSE:HE2	1:A:41:ARG:NH1	2.34	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	56/86 (65%)	56 (100%)	0	0	100	100
1	B	54/86 (63%)	52 (96%)	2 (4%)	0	100	100
1	C	58/86 (67%)	58 (100%)	0	0	100	100
1	D	60/86 (70%)	59 (98%)	1 (2%)	0	100	100
All	All	228/344 (66%)	225 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	47/60 (78%)	43 (92%)	4 (8%)	13	11
1	B	45/60 (75%)	42 (93%)	3 (7%)	20	19
1	C	49/60 (82%)	46 (94%)	3 (6%)	23	23
1	D	51/60 (85%)	49 (96%)	2 (4%)	39	45
All	All	192/240 (80%)	180 (94%)	12 (6%)	35	22

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38[A]	MSE
1	A	38[B]	MSE
1	A	42[A]	GLN
1	A	42[B]	GLN
1	B	38[A]	MSE
1	B	38[B]	MSE
1	B	61	LEU
1	C	7	THR
1	C	8	SER
1	C	46	LEU
1	D	53[A]	ARG
1	D	53[B]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	55	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	52/86 (60%)	0.87	8 (15%)	3	3	39, 69, 101, 110	10 (19%)
1	B	51/86 (59%)	0.93	6 (11%)	6	7	45, 66, 86, 102	9 (17%)
1	C	56/86 (65%)	0.61	2 (3%)	46	48	24, 36, 76, 114	4 (7%)
1	D	56/86 (65%)	0.66	1 (1%)	71	72	23, 36, 69, 91	1 (1%)
All	All	215/344 (62%)	0.76	17 (7%)	15	16	23, 52, 88, 114	24 (11%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	44	HIS	5.2
1	B	42[A]	GLN	4.6
1	A	11	GLY	4.1
1	A	45	PRO	3.9
1	A	44	HIS	3.7
1	B	53	ARG	3.0
1	B	27	PRO	2.9
1	B	46	LEU	2.7
1	A	42[A]	GLN	2.6
1	B	45	PRO	2.5
1	D	61	LEU	2.5
1	A	49	HIS	2.4
1	C	7	THR	2.4
1	C	61	LEU	2.4
1	A	22	LEU	2.2
1	A	53	ARG	2.1
1	A	61	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.