wwPDB X-ray Structure Validation Summary Report (i)

Jan 31, 2016 – 06:46 PM GMT

PDB ID	:	1CDE
Title	:	STRUCTURES OF APO AND COMPLEXED ESCHERICHIA COLI GLYCINAMIDE RIBONUCLEOTIDE TRANSFORMYLASE
Authors	:	Almassy, R.J.; Janson, C.A.; Kan, C.-C.; Hostomska, Z.
Deposited on	:	1992-05-15
Resolution	:	2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	212	3% 67% 25% 6%
1	B	212	4% 66% 26% 6%
1	C	212	4% 65% 26% 7%
1	D	212	2% 67% 25% 6%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6668 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHORIBOSYL-GLYCINAMIDE FORMYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	209	Total	C	N	O	S	0	0	0
			1617	1024	287	301	5			
1	B	209	Total	C	N	O	S	0	0	0
			1617	1024	287	301	5			
1	C	209	Total	C	N	O	S	0	0	0
			1617	1024	287	301	5			
1	D	209	Total	C	N	O	S	0	0	0
			1617	1024	287	301	5			

- Molecule 2 is GLYCINAMIDE RIBONUCLEOTIDE (three-letter code: GAR) (formula: $C_7H_{13}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			18	7	2	8	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C N O P 18 7 2 8 1	0	0
2	C	1	Total C N O P 18 7 2 8 1	0	0
2	D	1	Total C N O P 18 7 2 8 1	0	0

- Molecule 3 is 5-DEAZAFOLIC ACID (three-letter code: DZF) (formula: $C_{20}H_{20}N_6O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 32 20 6 6	0	0
3	B	1	Total C N O 32 20 6 6	0	0
3	C	1	Total C N O 32 20 6 6	0	0
3	D	1	Total C N O 32 20 6 6	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PHOSPHORIBOSYL-GLYCINAMIDE FORMYLTRANSFERASE

Chain A:



- Molecule 1: PHOSPHORIBOSYL-GLYCINAMIDE FORMYLTRANSFERASE

Chain B:



- Molecule 1: PHOSPHORIBOSYL-GLYCINAMIDE FORMYLTRANSFERASE

Chain C:



• Molecule 1: PHOSPHORIBOSYL-GLYCINAMIDE FORMYLTRANSFERASE

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	76.70Å 72.60Å 57.00Å 111.50° 82.80° 62.60°	Depositor
Resolution (Å)	(Not available) – 2.50 8.00 – 2.51	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.50) 66.4 (8.00-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.60 (at 2.52Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.250 , (Not available) 0.228 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å^2)	27.6	Xtriage
Anisotropy	0.404	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 68.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 25408 reflections	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	6668	wwPDB-VP
Average B, all atoms (Å^2)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.41% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAR, DZF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	1.03	1/1654 (0.1%)	1.82	35/2248 (1.6%)
1	B	1.03	1/1654 (0.1%)	1.82	35/2248 (1.6%)
1	C	1.03	1/1654 (0.1%)	1.82	35/2248 (1.6%)
1	D	1.03	1/1654 (0.1%)	1.82	35/2248 (1.6%)
All	All	1.03	4/6616 (0.1%)	1.82	140/8992 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	84	VAL	CA-CB	5.67	1.66	1.54
1	A	84	VAL	CA-CB	5.66	1.66	1.54
1	D	84	VAL	CA-CB	5.65	1.66	1.54
1	C	84	VAL	CA-CB	5.63	1.66	1.54

The worst 5 of 140 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	64	ARG	NE-CZ-NH1	12.63	126.61	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	D	64	ARG	NE-CZ-NH1	12.59	126.59	120.30
1	C	64	ARG	NE-CZ-NH1	12.56	126.58	120.30
1	A	64	ARG	NE-CZ-NH1	12.56	126.58	120.30
1	A	168	ARG	NE-CZ-NH2	-11.53	114.53	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	177	TYR	Sidechain
1	B	177	TYR	Sidechain
1	C	177	TYR	Sidechain
1	D	177	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1617	0	1592	19	0
1	B	1617	0	1592	20	4
1	C	1617	0	1592	22	4
1	D	1617	0	1592	20	0
2	A	18	0	13	1	0
2	B	18	0	13	1	0
2	C	18	0	13	1	0
2	D	18	0	13	1	0
3	A	32	0	19	1	0
3	B	32	0	19	1	0
3	C	32	0	19	1	0
3	D	32	0	19	1	0
All	All	6668	0	6496	81	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 81 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:LEU:HD13	1:B:165:ILE:HD11	1.61	0.83
1:C:125:LEU:HD13	1:C:165:ILE:HD11	1.61	0.83
1:A:125:LEU:HD13	1:A:165:ILE:HD11	1.61	0.82
1:D:125:LEU:HD13	1:D:165:ILE:HD11	1.61	0.81
1:A:4:VAL:HG13	1:A:80:PRO:HB3	1.73	0.70

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:HIS:NE2	1:C:70:GLU:OE1[1_465]	1.80	0.40
1:B:70:GLU:OE1	1:C:73:HIS:NE2[1_465]	1.86	0.34
1:B:73:HIS:CE1	1:C:70:GLU:OE1[1_465]	1.96	0.24
1:B:70:GLU:OE1	1:C:73:HIS:CE1[1_465]	1.97	0.23

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/212 (98%)	194 (94%)	11 (5%)	2 (1%)	19	34
1	B	207/212 (98%)	194 (94%)	11 (5%)	2 (1%)	19	34
1	C	207/212 (98%)	194 (94%)	11 (5%)	2 (1%)	19	34
1	D	207/212 (98%)	194 (94%)	11 (5%)	2 (1%)	19	34
All	All	828/848 (98%)	776 (94%)	44 (5%)	8 (1%)	19	34

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	193	GLU
1	B	193	GLU
1	C	193	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	193	GLU
1	A	62	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/171 (99%)	144 (85%)	25 (15%)	4	7
1	B	169/171 (99%)	144 (85%)	25 (15%)	4	7
1	C	169/171 (99%)	144 (85%)	25 (15%)	4	7
1	D	169/171 (99%)	144 (85%)	25 (15%)	4	7
All	All	676/684 (99%)	576 (85%)	100 (15%)	4	7

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	160	ASP
1	C	43	LEU
1	D	143	LEU
1	B	177	TYR
1	C	1	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	174	HIS
1	C	13	ASN
1	D	36	ASN
1	B	108	HIS
1	B	127	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	GAR	A	222	-	17,18,18	2.72	5 (29%)	21,26,26	1.43	5 (23%)
3	DZF	A	225	-	28,34,34	3.15	8 (28%)	31,47,47	3.37	15 (48%)
2	GAR	B	222	-	17,18,18	2.73	5 (29%)	21,26,26	1.42	5 (23%)
3	DZF	B	225	-	28,34,34	3.15	8 (28%)	31,47,47	3.37	15 (48%)
2	GAR	C	222	-	17,18,18	2.72	5 (29%)	21,26,26	1.42	5 (23%)
3	DZF	C	225	-	28,34,34	3.15	8 (28%)	31,47,47	3.37	15 (48%)
2	GAR	D	222	-	17,18,18	2.72	5 (29%)	21,26,26	1.43	5 (23%)
3	DZF	D	225	-	28,34,34	3.14	8 (28%)	31,47,47	3.37	15 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAR	A	222	-	-	0/12/28/28	0/1/1/1
3	DZF	A	225	-	-	0/16/22/22	0/3/3/3
2	GAR	B	222	-	-	0/12/28/28	0/1/1/1
3	DZF	B	225	-	-	0/16/22/22	0/3/3/3
2	GAR	C	222	-	-	0/12/28/28	0/1/1/1
3	DZF	C	225	-	-	0/16/22/22	0/3/3/3
2	GAR	D	222	-	-	0/12/28/28	0/1/1/1
3	DZF	D	225	-	-	0/16/22/22	0/3/3/3

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	222	GAR	P15-O18	-6.74	1.40	1.51
2	C	222	GAR	P15-O18	-6.74	1.40	1.51
2	A	222	GAR	P15-O18	-6.73	1.40	1.51
2	D	222	GAR	P15-O18	-6.73	1.40	1.51
3	B	225	DZF	C8A-N8	-3.48	1.31	1.37

The worst 5 of 80 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	225	DZF	C9-C6-C5	-10.57	105.18	121.98
3	B	225	DZF	C9-C6-C5	-10.56	105.19	121.98
3	A	225	DZF	C9-C6-C5	-10.55	105.20	121.98
3	D	225	DZF	C9-C6-C5	-10.54	105.22	121.98
3	C	225	DZF	C6-C7-N8	-7.57	110.96	123.82

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	222	GAR	1	0
3	A	225	DZF	1	0
2	B	222	GAR	1	0
3	B	225	DZF	1	0
2	C	222	GAR	1	0
3	C	225	DZF	1	0
2	D	222	GAR	1	0
3	D	225	DZF	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	209/212 (98%)	-0.19	6 (2%) 55 60	2, 19, 59, 86	0
1	B	209/212 (98%)	-0.16	9 (4%) 39 44	2, 19, 59, 86	0
1	C	209/212 (98%)	-0.16	9 (4%) 39 44	2, 19, 59, 86	0
1	D	209/212 (98%)	-0.29	5 (2%) 62 66	2, 19, 59, 86	0
All	All	836/848 (98%)	-0.20	29 (3%) 48 53	2, 20, 60, 86	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	60	ALA	8.6
1	C	59	SER	5.8
1	B	62	ASP	5.5
1	C	62	ASP	5.4
1	C	60	ALA	5.3

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	GAR	A	222	18/18	0.93	0.16	1.41	16,23,36,44	0
2	GAR	B	222	18/18	0.90	0.16	1.10	16,23,36,44	0
3	DZF	C	225	32/32	0.91	0.17	0.71	13,56,102,106	0
3	DZF	D	225	32/32	0.90	0.19	0.53	13,56,102,106	0
2	GAR	C	222	18/18	0.95	0.13	0.47	16,23,36,44	0
3	DZF	B	225	32/32	0.91	0.17	0.47	13,56,102,106	0
3	DZF	A	225	32/32	0.92	0.16	0.26	13,56,102,106	0
2	GAR	D	222	18/18	0.96	0.11	-0.02	16,23,36,44	0

6.5 Other polymers (i)

There are no such residues in this entry.