



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2CDR
Title : Crystal structures of caspase-3 in complex with aza-peptide epoxide inhibitors.
Authors : Ganesan, R.; Jelakovic, S.; Campbell, A.J.; Li, Z.Z.; Asgian, J.L.; Powers, J.C.; Gruetter, M.G.
Deposited on : 2006-01-27
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	49% 87% 13%
2	B	103	69% 80% 20%
3	I	5	40% 80% 20%

2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 2429 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CASPASE-3 SUBUNIT P17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C 1161	N 716	O 208	S 228	9	0	0

- Molecule 2 is a protein called CASPASE-3 SUBUNIT P12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	103	Total	C 843	N 547	O 135	S 154	7	0	0

- Molecule 3 is a protein called AZA-PEPTIDE EXPOXIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	I	5	Total	C 62	N 42	O 6	S 14	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	126	Total O 126 126	0	0
4	A	231	Total O 231 231	0	0
4	I	6	Total O 6 6	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CASPASE-3 SUBUNIT P17



- Molecule 2: CASPASE-3 SUBUNIT P12



- Molecule 3: AZA-PEPTIDE EXPOXIDE



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	67.47 Å 83.78 Å 96.25 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.49 – 1.70 19.80 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.9 (19.49-1.70) 95.6 (19.80-1.70)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.37 (at 1.70 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R , R_{free}	0.177 , 0.204 0.368 , 0.366	Depositor DCC
R_{free} test set	2876 reflections (11.07%)	DCC
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.655	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 28859 reflections	Xtriage
F_o, F_c correlation	0.82	EDS
Total number of atoms	2429	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MY1, PHQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	0/1176	0.58	0/1574
2	B	0.34	0/868	0.63	0/1171
3	I	0.25	0/23	0.62	0/30
All	All	0.31	0/2067	0.60	0/2775

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1161	0	1152	19	12
2	B	843	0	807	13	15
3	I	62	0	44	3	0
4	A	231	0	0	8	6
4	B	126	0	0	6	4
4	I	6	0	0	0	0
All	All	2429	0	2003	31	21

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:185:HIS:O	4:B:2044:HOH:O	1.77	1.01
1:A:61:MET:HG2	3:I:905:MY1:H20	1.48	0.92
1:A:174:THR:OG1	4:A:2221:HOH:O	1.60	0.74
2:B:185:HIS:HB2	4:B:2044:HOH:O	1.88	0.74
1:A:87:ASN:HB2	4:A:2121:HOH:O	1.89	0.71
2:B:241:ARG:NH1	4:B:2096:HOH:O	2.19	0.70
2:B:252:PHE:HB2	4:B:2110:HOH:O	1.90	0.70
2:B:175:ALA:N	4:B:2028:HOH:O	2.27	0.67
2:B:228:ASP:OD1	2:B:229:LYS:HG3	1.97	0.64
1:A:100:MET:HG3	1:A:139:ILE:HG23	1.84	0.59
2:B:177:GLY:HA3	4:B:2035:HOH:O	2.04	0.58
1:A:74:LEU:HD13	1:A:117:VAL:HG11	1.87	0.57
1:A:153:GLY:HA3	4:A:2196:HOH:O	2.07	0.54
1:A:149:ARG:HH11	1:A:149:ARG:HA	1.74	0.53
1:A:165:GLY:HA2	3:I:905:MY1:H27	1.94	0.50
2:B:240:ASN:OD1	2:B:263:PRO:HB2	2.12	0.50
1:A:93:ARG:NH1	4:A:2131:HOH:O	2.45	0.50
2:B:206:TRP:HH2	2:B:256:PHE:HB3	1.76	0.50
2:B:213:SER:O	2:B:217:GLN:HG3	2.12	0.49
2:B:207:ARG:HA	2:B:213:SER:HA	1.96	0.47
1:A:34:ASP:OD1	2:B:271:LYS:HE2	2.15	0.46
2:B:262:ILE:O	2:B:262:ILE:HG23	2.15	0.46
1:A:89:ASN:O	4:A:2124:HOH:O	2.20	0.46
1:A:75:ARG:HD2	4:A:2121:HOH:O	2.16	0.45
1:A:149:ARG:NH1	1:A:149:ARG:HB3	2.32	0.45
1:A:149:ARG:HH11	1:A:149:ARG:CA	2.31	0.43
1:A:89:ASN:HB3	4:A:2038:HOH:O	2.18	0.43
1:A:153:GLY:CA	4:A:2196:HOH:O	2.65	0.43
1:A:74:LEU:CD1	1:A:117:VAL:HG11	2.48	0.42
1:A:41:TYR:HB2	1:A:112:SER:OG	2.21	0.41
1:A:61:MET:SD	3:I:905:MY1:H19	2.60	0.41

All (21) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:SER:OG	2:B:253:ASP:OD2[2_665]	1.16	1.04
1:A:173:GLU:OE2	2:B:186:LYS:CG[3_656]	1.49	0.71
1:A:173:GLU:OE2	2:B:186:LYS:CD[3_656]	1.53	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:ILE:N	2:B:187:ILE:O[3_656]	1.59	0.61
2:B:234:HIS:CE1	2:B:234:HIS:CE1[3_656]	1.74	0.46
1:A:172:ILE:O	2:B:187:ILE:N[3_656]	1.75	0.45
1:A:173:GLU:CD	2:B:186:LYS:CG[3_656]	1.76	0.44
2:B:234:HIS:NE2	2:B:234:HIS:NE2[3_656]	1.78	0.42
4:A:2126:HOH:O	4:B:2043:HOH:O[6_554]	1.79	0.41
4:A:2063:HOH:O	4:B:2060:HOH:O[2_665]	1.85	0.35
1:A:58:SER:OG	2:B:253:ASP:CG[2_665]	1.86	0.34
1:A:174:THR:CG2	2:B:187:ILE:CG2[3_656]	1.87	0.33
1:A:89:ASN:N	2:B:181:ASP:OD2[6_554]	1.90	0.30
4:A:2085:HOH:O	4:A:2085:HOH:O[3_656]	1.91	0.29
1:A:173:GLU:OE1	2:B:185:HIS:O[3_656]	2.00	0.20
4:A:2156:HOH:O	4:A:2156:HOH:O[3_656]	2.06	0.14
1:A:87:ASN:ND2	2:B:181:ASP:O[6_554]	2.09	0.11
4:A:2079:HOH:O	4:B:2045:HOH:O[6_554]	2.09	0.11
2:B:234:HIS:CE1	2:B:234:HIS:NE2[3_656]	2.13	0.07
1:A:173:GLU:OE1	2:B:186:LYS:CG[3_656]	2.14	0.06
4:A:2059:HOH:O	4:B:2107:HOH:O[2_665]	2.17	0.03

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	145/147 (99%)	140 (97%)	5 (3%)	0	100 100
2	B	101/103 (98%)	100 (99%)	1 (1%)	0	100 100
3	I	1/5 (20%)	1 (100%)	0	0	100 100
All	All	247/255 (97%)	241 (98%)	6 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/132 (100%)	132 (100%)	0	100	100
2	B	90/90 (100%)	90 (100%)	0	100	100
3	I	3/3 (100%)	3 (100%)	0	100	100
All	All	225/225 (100%)	225 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/147 (100%)	2.17	72 (48%) 0 0	12, 18, 29, 56	0
2	B	103/103 (100%)	2.75	71 (68%) 0 0	11, 15, 28, 34	0
3	I	3/5 (60%)	2.81	2 (66%) 0 0	23, 23, 26, 31	0
All	All	253/255 (99%)	2.42	145 (57%) 0 0	11, 17, 29, 56	0

All (145) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	168	LEU	7.4
2	B	204	TYR	7.1
2	B	180	ASP	6.5
1	A	61	MET	6.1
2	B	256	PHE	5.8
1	A	175	ASP	5.6
2	B	183	ALA	5.5
1	A	172	ILE	5.3
2	B	252	PHE	5.2
1	A	174	THR	5.2
2	B	184	CYS	5.0
2	B	262	ILE	4.9
1	A	170	CYS	4.8
2	B	250	PHE	4.8
2	B	206	TRP	4.8
3	I	904	VAL	4.8
1	A	33	LEU	4.8
2	B	200	ALA	4.7
2	B	255	THR	4.6
1	A	128	PHE	4.6
2	B	215	PHE	4.6
1	A	29	SER	4.5
2	B	203	TYR	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	210	LYS	4.3
2	B	253	ASP	4.1
1	A	173	GLU	4.1
1	A	169	ASP	4.0
1	A	34	ASP	3.9
1	A	58	SER	3.8
1	A	96	ILE	3.8
2	B	199	THR	3.8
2	B	176	SER	3.8
2	B	246	GLU	3.7
2	B	266	VAL	3.7
2	B	201	PRO	3.7
2	B	273	LEU	3.6
1	A	60	GLY	3.6
2	B	247	PHE	3.6
1	A	139	ILE	3.5
2	B	270	THR	3.4
2	B	182	MET	3.4
2	B	254	ALA	3.4
2	B	177	GLY	3.4
2	B	243	VAL	3.4
2	B	264	CYS	3.4
2	B	214	TRP	3.3
2	B	195	TYR	3.3
2	B	276	TYR	3.3
2	B	235	ILE	3.3
1	A	62	THR	3.3
2	B	241	ARG	3.3
1	A	119	LEU	3.3
2	B	230	LEU	3.3
1	A	171	GLY	3.3
1	A	151	LEU	3.2
2	B	265	ILE	3.2
2	B	261	GLN	3.2
1	A	118	LEU	3.2
1	A	147	ARG	3.2
1	A	162	ALA	3.1
2	B	189	VAL	3.1
1	A	55	PHE	3.1
2	B	249	SER	3.1
1	A	93	ARG	3.1
2	B	187	ILE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	134	VAL	3.0
2	B	219	LEU	3.0
2	B	236	LEU	3.0
1	A	36	SER	3.0
2	B	185	HIS	3.0
2	B	216	ILE	2.9
2	B	258	ALA	2.9
2	B	213	SER	2.9
2	B	239	VAL	2.9
1	A	114	PHE	2.9
1	A	127	ILE	2.9
1	A	165	GLY	2.8
2	B	188	PRO	2.8
1	A	59	THR	2.8
2	B	220	CYS	2.8
1	A	67	THR	2.8
1	A	163	CYS	2.8
2	B	227	ALA	2.8
2	B	178	VAL	2.8
1	A	65	SER	2.7
1	A	155	PRO	2.7
2	B	257	HIS	2.7
1	A	143	PHE	2.7
1	A	46	LEU	2.7
2	B	244	ALA	2.7
1	A	152	THR	2.7
2	B	193	PHE	2.7
2	B	208	ASN	2.6
1	A	100	MET	2.6
1	A	31	ILE	2.6
1	A	123	GLU	2.6
1	A	91	LEU	2.6
2	B	223	LEU	2.6
1	A	83	TYR	2.6
1	A	72	ALA	2.6
1	A	109	SER	2.5
1	A	99	LEU	2.5
1	A	160	ILE	2.5
1	A	136	LEU	2.5
2	B	275	PHE	2.5
2	B	226	TYR	2.4
2	B	181	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	50	ILE	2.4
1	A	71	ALA	2.4
2	B	197	TYR	2.3
1	A	142	PHE	2.3
2	B	175	ALA	2.3
2	B	196	ALA	2.3
1	A	110	LYS	2.3
2	B	269	LEU	2.3
2	B	263	PRO	2.3
1	A	66	GLY	2.3
1	A	101	ARG	2.2
2	B	259	LYS	2.2
1	A	97	VAL	2.2
1	A	73	ASN	2.2
1	A	75	ARG	2.2
1	A	49	ILE	2.2
1	A	44	MET	2.2
2	B	194	LEU	2.2
2	B	240	ASN	2.2
1	A	159	ILE	2.2
2	B	207	ARG	2.2
1	A	158	PHE	2.2
1	A	167	GLU	2.2
1	A	92	THR	2.2
1	A	135	ASP	2.2
3	I	902	ASP	2.1
2	B	268	MET	2.1
1	A	74	LEU	2.1
1	A	103	VAL	2.1
1	A	41	TYR	2.1
1	A	117	VAL	2.0
2	B	274	TYR	2.0
1	A	86	ARG	2.0
2	B	234	HIS	2.0
2	B	245	THR	2.0
1	A	30	GLY	2.0
1	A	51	ASN	2.0
1	A	150	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.