



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:31 PM GMT

PDB ID : 4CEX
Title : 1.59 Å resolution Fluoride inhibited *Sporosarcina pasteurii* urease
Authors : Benini, S.; Cianci, M.; Ciurli, S.
Deposited on : 2013-11-12
Resolution : 1.59 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3815 (1.60-1.56)
Clashscore	102246	4131 (1.60-1.56)
Ramachandran outliers	100387	4021 (1.60-1.56)
Sidechain outliers	100360	4018 (1.60-1.56)
RSRZ outliers	91569	3823 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	100	<div> <div style="width: 97%;"></div> <div>97%</div> </div>
2	B	126	<div> <div style="width: 91%;"></div> <div>91%</div> <div style="width: 6%;"></div> <div>6%</div> </div>
3	C	570	<div> <div style="width: 2%;"></div> <div>2%</div> <div style="width: 93%;"></div> <div>93%</div> <div style="width: 7%;"></div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	1102	-	-	-	X
4	EDO	A	1103	-	-	-	X
4	EDO	B	1127	-	-	-	X
4	EDO	B	1128	-	-	-	X
4	EDO	C	1574	-	-	-	X
4	EDO	C	1575	-	-	-	X
4	EDO	C	1576	-	-	-	X
4	EDO	C	1578	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UREASE SUBUNIT GAMMA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	2	0
			784	496	130	151	7			

- Molecule 2 is a protein called UREASE SUBUNIT BETA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	122	Total	C	N	O	S	0	1	0
			948	589	172	186	1			


- Molecule 3 is a protein called UREASE SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	570	Total	C	N	O	S	0	13	0
			4372	2752	745	848	27			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	19	GLU	ARG	CONFLICT	UNP P41020
C	28	TRP	GLY	CONFLICT	UNP P41020
C	29	ILE	-	INSERTION	UNP P41020
C	36	THR	TYR	CONFLICT	UNP P41020
C	37	THR	TYR	CONFLICT	UNP P41020
C	38	TYR	LEU	CONFLICT	UNP P41020
C	263	LEU	VAL	CONFLICT	UNP P41020
C	420	ILE	MET	CONFLICT	UNP P41020

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	2	Total	Ni	0	0
			2	2		

- Molecule 7 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	2	Total	F	0	0
			2	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	126	Total	O	0	0
			126	126		
8	B	182	Total	O	0	0
			182	182		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	518	Total	O	0	0
			518	518		

• Molecule 1: UREASE SUBUNIT GAMMA

• Molecule 2: UREASE SUBUNIT BETA

- Molecule 3: UREASE SUBUNIT ALPHA

Chain C:

Amino Acid	Frequency Category
M1	Green
R5	Green
D26	Green
Q47	Green
K48	Red
V49	Green
E64	Green
L70	Green
D86	Green
D104	Green
G132	Green
H137	Green
T155	Green
F158	Green
P177	Green
H201	Green
H222	Green
V225	Green
H249	Orange
F274	Green
G281	Green
S300	Green
I316	Green
D317	Green
K320	Red
K321	Red
C322	Green
H323	Red
H324	Red
L325	Red
K326	Red
Q327	Red
H328	Red
I329	Red
H352	Green
H360	Green
M367	Green
K383	Green
K386	Green
Q387	Green
K388	Green
G389	Green
P390	Red
N396	Red
K409	Green
T470	Green
P471	Green
R513	Green
C520	Green
R566	Green
F570	Green

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	131.33Å 131.33Å 188.88Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	113.74 – 1.59 113.74 – 1.59	Depositor EDS
% Data completeness (in resolution range)	99.8 (113.74-1.59) 99.8 (113.74-1.59)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.82 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.130 , 0.150 0.145 , 0.163	Depositor DCC
R_{free} test set	6467 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	11.2	Xtriage
Anisotropy	0.534	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 46.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 128733 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6993	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, F, EDO, KCX, SO4, CXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/790	0.75	0/1064
2	B	0.53	0/963	0.71	0/1296
3	C	0.56	1/4480 (0.0%)	0.79	5/6071 (0.1%)
All	All	0.55	1/6233 (0.0%)	0.77	5/8431 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	520	CYS	CB-SG	-8.89	1.67	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	566	ARG	NE-CZ-NH1	6.76	123.68	120.30
3	C	5	ARG	NE-CZ-NH2	-6.45	117.08	120.30
3	C	388	ARG	NE-CZ-NH1	5.74	123.17	120.30
3	C	520	CYS	CB-CA-C	-5.71	98.98	110.40
3	C	360	MET	CG-SD-CE	-5.27	91.77	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	784	0	795	2	0
2	B	948	0	933	4	0
3	C	4372	0	4351	31	0
4	A	12	0	18	0	0
4	B	8	0	12	2	0
4	C	24	0	36	1	1
5	B	5	0	0	0	0
5	C	10	0	0	0	0
6	C	2	0	0	0	0
7	C	2	0	0	0	0
8	A	126	0	0	1	0
8	B	182	0	0	1	0
8	C	518	0	0	4	3
All	All	6993	0	6145	35	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:320[A]:MET:HA	3:C:325:LEU:HD23	1.48	0.95
3:C:320[B]:MET:HA	3:C:325:LEU:HD23	1.56	0.86
3:C:201:HIS:HE1	3:C:222:HIS:H	1.26	0.84
3:C:64[A]:GLU:OE2	8:C:2104:HOH:O	1.99	0.80
3:C:201:HIS:CE1	3:C:222:HIS:H	2.03	0.76

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:C:2116:HOH:O	8:C:2214:HOH:O[2_665]	1.51	0.69
8:C:2034:HOH:O	8:C:2034:HOH:O[11_555]	1.80	0.40
4:C:1576:EDO:O1	8:C:2432:HOH:O[12_565]	1.91	0.29

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/100 (100%)	100 (100%)	0	0	100	100
2	B	121/126 (96%)	117 (97%)	3 (2%)	1 (1%)	24	6
3	C	579/570 (102%)	554 (96%)	24 (4%)	1 (0%)	52	27
All	All	800/796 (100%)	771 (96%)	27 (3%)	2 (0%)	46	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	367	MET
2	B	99	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/84 (99%)	83 (100%)	0	100	100
2	B	99/105 (94%)	98 (99%)	1 (1%)	82	65
3	C	465/460 (101%)	461 (99%)	4 (1%)	84	69
All	All	647/649 (100%)	642 (99%)	5 (1%)	86	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	5	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	158	PHE
3	C	249	HIS
3	C	317	ASP
3	C	325	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	267	ASN
3	C	418	GLN
3	C	352	HIS
3	C	201	HIS
3	C	387	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CXM	A	1	1	6,10,11	0.63	0	4,11,13	0.38	0
3	KCX	C	220	3,6	7,11,12	0.62	0	7,12,14	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	A	1	1	-	0/6/10/12	0/0/0/0
3	KCX	C	220	3,6	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	1101	-	3,3,3	0.52	0	2,2,2	0.64	0
4	EDO	A	1102	-	3,3,3	0.59	0	2,2,2	0.07	0
4	EDO	A	1103	-	3,3,3	0.83	0	2,2,2	0.19	0
4	EDO	B	1127	-	3,3,3	0.54	0	2,2,2	0.15	0
4	EDO	B	1128	-	3,3,3	0.31	0	2,2,2	0.73	0
5	SO4	B	1129	-	4,4,4	0.47	0	6,6,6	0.27	0
4	EDO	C	1573	-	3,3,3	0.41	0	2,2,2	0.43	0
4	EDO	C	1574	-	3,3,3	0.35	0	2,2,2	0.06	0
4	EDO	C	1575	-	3,3,3	0.53	0	2,2,2	0.21	0
4	EDO	C	1576	-	3,3,3	0.57	0	2,2,2	0.12	0
4	EDO	C	1577	-	3,3,3	0.47	0	2,2,2	0.46	0
4	EDO	C	1578	-	3,3,3	0.32	0	2,2,2	0.33	0
5	SO4	C	1579	-	4,4,4	0.72	0	6,6,6	0.90	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	C	1580	-	4,4,4	0.39	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1101	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1102	-	-	0/1/1/1	0/0/0/0
4	EDO	A	1103	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1127	-	-	0/1/1/1	0/0/0/0
4	EDO	B	1128	-	-	0/1/1/1	0/0/0/0
5	SO4	B	1129	-	-	0/0/0/0	0/0/0/0
4	EDO	C	1573	-	-	0/1/1/1	0/0/0/0
4	EDO	C	1574	-	-	0/1/1/1	0/0/0/0
4	EDO	C	1575	-	-	0/1/1/1	0/0/0/0
4	EDO	C	1576	-	-	0/1/1/1	0/0/0/0
4	EDO	C	1577	-	-	0/1/1/1	0/0/0/0
4	EDO	C	1578	-	-	0/1/1/1	0/0/0/0
5	SO4	C	1579	-	-	0/0/0/0	0/0/0/0
5	SO4	C	1580	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1128	EDO	2	0
4	C	1576	EDO	0	1
4	C	1578	EDO	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	99/100 (99%)	-0.81	0 100 100	9, 12, 18, 23	0
2	B	122/126 (96%)	-0.71	0 100 100	10, 13, 22, 41	0
3	C	569/570 (99%)	-0.70	10 (1%) 71 72	8, 11, 22, 50	0
All	All	790/796 (99%)	-0.71	10 (1%) 79 80	8, 11, 22, 50	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	396	ASN	3.8
3	C	325	LEU	3.4
3	C	329	ILE	3.1
3	C	328	ASN	2.9
3	C	321	VAL	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CXM	A	1	11/12	0.98	0.06	-	10,13,16,17	0
3	KCX	C	220	12/13	0.98	0.05	-	7,8,9,9	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	C	1576	4/4	0.86	0.23	12.06	28,28,29,30	0
4	EDO	B	1127	4/4	0.92	0.13	10.36	29,29,30,35	0
4	EDO	B	1128	4/4	0.86	0.32	10.03	33,33,34,35	0
4	EDO	A	1102	4/4	0.86	0.19	9.81	33,39,40,42	0
4	EDO	A	1103	4/4	0.86	0.12	9.53	20,23,23,23	0
4	EDO	C	1578	4/4	0.84	0.17	2.71	36,36,38,40	0
4	EDO	C	1574	4/4	0.98	0.10	2.28	21,22,23,24	0
4	EDO	C	1575	4/4	0.80	0.25	2.13	37,41,44,51	0
4	EDO	A	1101	4/4	0.97	0.07	0.81	13,16,18,20	0
4	EDO	C	1573	4/4	0.96	0.05	0.27	23,24,25,26	0
6	NI	C	601	1/1	1.00	0.04	-0.90	8,8,8,8	0
6	NI	C	600	1/1	1.00	0.05	-1.15	9,9,9,9	0
7	F	C	1571	1/1	1.00	0.04	-1.27	8,8,8,8	0
7	F	C	1572	1/1	0.99	0.04	-2.23	13,13,13,13	0
5	SO4	B	1129	5/5	0.90	0.20	-	47,49,52,54	0
5	SO4	C	1580	5/5	0.93	0.15	-	61,62,62,62	5
4	EDO	C	1577	4/4	0.88	0.18	-	33,33,34,34	0
5	SO4	C	1579	5/5	0.87	0.23	-	19,19,27,29	5

6.5 Other polymers

There are no such residues in this entry.