



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:52 AM GMT

PDB ID : 3CF1
Title : Structure of P97/vcp in complex with ADP/ADP.alfx
Authors : Davies, J.M.; Delabarre, B.; Brunger, A.T.; Weis, W.I.
Deposited on : 2008-03-01
Resolution : 4.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1067 (5.20-3.60)
Clashscore	102246	1175 (5.20-3.60)
Ramachandran outliers	100387	1114 (5.20-3.60)
Sidechain outliers	100360	1096 (5.20-3.60)
RSRZ outliers	91569	1071 (5.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	806	<div> <div>3%</div> <div>31%</div> <div>55%</div> <div>11%</div> </div>
1	B	806	<div> <div>3%</div> <div>32%</div> <div>55%</div> <div>10%</div> </div>
1	C	806	<div> <div>2%</div> <div>30%</div> <div>57%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AF3	A	915	-	-	X	-
3	AF3	B	915	-	-	X	-
3	AF3	C	915	-	-	X	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 17126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transitional endoplasmic reticulum ATPase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	719	Total	C	N	O	S	0	0	0
			5634	3547	990	1067	30			
1	B	723	Total	C	N	O	S	0	0	0
			5659	3561	996	1072	30			
1	C	723	Total	C	N	O	S	0	0	0
			5659	3561	996	1072	30			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 3 is ALUMINUM FLUORIDE (three-letter code: AF3) (formula: AlF_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Al	F	0	0
			4	1	3		
3	B	1	Total	Al	F	0	0
			4	1	3		
3	C	1	Total	Al	F	0	0
			4	1	3		



• Molecule 1: Transitional endoplasmic reticulum ATPase





4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	162.66Å 178.02Å 321.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.87 – 4.40 49.60 – 4.40	Depositor EDS
% Data completeness (in resolution range)	82.2 (29.87-4.40) 82.0 (49.60-4.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 4.45Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.229 , 0.286 0.309 , 0.332	Depositor DCC
R_{free} test set	1865 reflections (8.12%)	DCC
Wilson B-factor (Å ²)	176.9	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 208.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 28099 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	17126	wwPDB-VP
Average B, all atoms (Å ²)	272.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, AF3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	6/5724 (0.1%)	0.61	1/7727 (0.0%)
1	B	0.42	2/5751 (0.0%)	0.61	2/7767 (0.0%)
1	C	0.42	1/5751 (0.0%)	0.61	2/7767 (0.0%)
All	All	0.43	9/17226 (0.1%)	0.61	5/23261 (0.0%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	755	TYR	CE1-CZ	7.97	1.49	1.38
1	A	755	TYR	CD2-CE2	-6.38	1.29	1.39
1	A	755	TYR	CG-CD1	6.23	1.47	1.39
1	A	755	TYR	CE2-CZ	6.20	1.46	1.38
1	B	625	ARG	CD-NE	-6.10	1.36	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	431	ASP	N-CA-C	8.06	132.75	111.00
1	B	431	ASP	N-CA-C	6.58	128.76	111.00
1	B	706	GLU	N-CA-C	5.90	126.92	111.00
1	C	433	GLU	N-CA-C	5.25	125.16	111.00
1	C	625	ARG	NE-CZ-NH2	-5.05	117.78	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5634	0	5705	657	0
1	B	5659	0	5731	660	0
1	C	5659	0	5731	634	0
2	A	54	0	24	8	0
2	B	54	0	24	14	0
2	C	54	0	24	11	0
3	A	4	0	0	2	0
3	B	4	0	0	9	0
3	C	4	0	0	2	0
All	All	17126	0	17239	1892	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 55.

The worst 5 of 1892 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:427:MET:HG3	1:A:431:ASP:CB	1.69	1.22
1:A:464:LEU:HD11	1:A:466:GLU:HB2	1.22	1.17
1:A:203:TYR:O	1:A:206:ILE:HG12	1.49	1.13
1:A:427:MET:HG3	1:A:431:ASP:HB2	1.20	1.09
1:A:66:GLU:HB2	1:A:147:ARG:NH1	1.68	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	711/806 (88%)	536 (75%)	155 (22%)	20 (3%)	6	46
1	B	719/806 (89%)	548 (76%)	148 (21%)	23 (3%)	5	43
1	C	719/806 (89%)	533 (74%)	163 (23%)	23 (3%)	5	43
All	All	2149/2418 (89%)	1617 (75%)	466 (22%)	66 (3%)	5	44

5 of 66 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	360	PHE
1	B	360	PHE
1	C	360	PHE
1	A	185	GLU
1	A	353	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	612/678 (90%)	605 (99%)	7 (1%)	80	90
1	B	615/678 (91%)	610 (99%)	5 (1%)	86	93
1	C	615/678 (91%)	609 (99%)	6 (1%)	82	91
All	All	1842/2034 (91%)	1824 (99%)	18 (1%)	82	91

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	183	HIS
1	B	283	GLU
1	C	434	ASP
1	A	556	GLU
1	B	173	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	260	ASN
1	B	458	GLN
1	C	490	GLN
1	B	285	ASN
1	B	348	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	ADP	A	807	-	22,29,29	1.45	4 (18%)	27,45,45	2.81	3 (11%)
2	ADP	A	900	-	22,29,29	1.38	3 (13%)	27,45,45	2.77	4 (14%)
3	AF3	A	915	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ADP	B	807	-	22,29,29	1.44	5 (22%)	27,45,45	2.79	4 (14%)
2	ADP	B	900	-	22,29,29	1.40	4 (18%)	27,45,45	2.88	3 (11%)
3	AF3	B	915	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ADP	C	807	-	22,29,29	1.43	4 (18%)	27,45,45	3.02	3 (11%)
2	ADP	C	900	-	22,29,29	1.42	3 (13%)	27,45,45	2.89	4 (14%)
3	AF3	C	915	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	807	-	-	0/12/32/32	0/3/3/3
2	ADP	A	900	-	-	0/12/32/32	0/3/3/3
3	AF3	A	915	-	-	0/0/0/0	0/0/0/0
2	ADP	B	807	-	-	0/12/32/32	0/3/3/3
2	ADP	B	900	-	-	0/12/32/32	0/3/3/3
3	AF3	B	915	-	-	0/0/0/0	0/0/0/0
2	ADP	C	807	-	-	0/12/32/32	0/3/3/3
2	ADP	C	900	-	-	0/12/32/32	0/3/3/3
3	AF3	C	915	-	-	0/0/0/0	0/0/0/0

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	807	ADP	C5-N7	-3.28	1.28	1.39
2	C	900	ADP	C5-N7	-3.24	1.28	1.39
2	A	900	ADP	C5-N7	-3.17	1.28	1.39
2	B	807	ADP	C5-N7	-3.10	1.28	1.39
2	A	807	ADP	C5-N7	-2.98	1.29	1.39

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	807	ADP	N3-C2-N1	-14.77	117.58	128.89
2	B	900	ADP	N3-C2-N1	-14.06	118.13	128.89
2	C	900	ADP	N3-C2-N1	-13.89	118.26	128.89
2	A	807	ADP	N3-C2-N1	-13.69	118.41	128.89
2	B	807	ADP	N3-C2-N1	-13.40	118.64	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

9 monomers are involved in 39 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	807	ADP	4	0
2	A	900	ADP	4	0
3	A	915	AF3	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	807	ADP	6	0
2	B	900	ADP	8	0
3	B	915	AF3	9	0
2	C	807	ADP	7	0
2	C	900	ADP	4	0
3	C	915	AF3	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	719/806 (89%)	0.18	24 (3%) 50 39	50, 267, 356, 574	0
1	B	723/806 (89%)	0.24	27 (3%) 45 35	49, 264, 348, 571	0
1	C	723/806 (89%)	0.18	17 (2%) 62 52	51, 266, 367, 616	0
All	All	2165/2418 (89%)	0.20	68 (3%) 52 41	49, 266, 360, 616	0

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	731	ILE	4.1
1	B	102	ILE	4.0
1	C	729	PRO	4.0
1	A	36	ASN	3.7
1	A	25	ARG	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ADP	B	807	27/27	0.86	0.40	1.62	306,306,306,306	0
2	ADP	C	807	27/27	0.88	0.34	0.77	306,306,306,306	0
2	ADP	A	807	27/27	0.91	0.31	0.49	306,306,306,306	0
3	AF3	C	915	4/4	0.91	0.27	0.41	306,306,306,306	0
3	AF3	A	915	4/4	0.86	0.24	0.16	306,306,306,306	0
2	ADP	C	900	27/27	0.79	0.32	0.05	306,306,306,306	0
2	ADP	B	900	27/27	0.88	0.23	-0.30	306,306,306,306	0
2	ADP	A	900	27/27	0.86	0.24	-0.34	306,306,306,306	0
3	AF3	B	915	4/4	0.92	0.19	-0.34	306,306,306,306	0

6.5 Other polymers [i](#)

There are no such residues in this entry.