



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:16 AM GMT

PDB ID : 2CG9
Title : CRYSTAL STRUCTURE OF AN HSP90-SBA1 CLOSED CHAPERONE COMPLEX
Authors : Ali, M.M.U.; Roe, S.M.; Prodromou, C.; Pearl, L.H.
Deposited on : 2006-03-01
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	677	
1	B	677	
2	X	134	
2	Y	134	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ATP	B	1678	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-DEPENDENT MOLECULAR CHAPERONE HSP82.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	609	Total	C	N	O	S	0	0	0
			4923	3147	804	963	9			
1	B	618	Total	C	N	O	S	0	0	0
			4997	3195	816	977	9			

- Molecule 2 is a protein called CO-CHAPERONE PROTEIN SBA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	115	Total	C	N	O	S	0	0	0
			962	621	154	185	2			
2	Y	115	Total	C	N	O	S	0	0	0
			962	621	154	185	2			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	127	ALA	GLU	ENGINEERED MUTATION	UNP P28707
Y	127	ALA	GLU	ENGINEERED MUTATION	UNP P28707

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	126.73 Å 126.73 Å 279.78 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	115.47 – 3.10 39.82 – 3.50	Depositor EDS
% Data completeness (in resolution range)	68.2 (115.47-3.10) 83.9 (39.82-3.50)	Depositor EDS
R_{merge}	0.19	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 3.48 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.312 , 0.353 0.351 , 0.371	Depositor DCC
R_{free} test set	1249 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	87.3	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 24801 reflections	Xtriage
F_o, F_c correlation	0.78	EDS
Total number of atoms	11906	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/5004	0.67	2/6743 (0.0%)
1	B	0.46	0/5080	0.67	1/6844 (0.0%)
2	X	0.50	1/986 (0.1%)	0.62	0/1335
2	Y	0.46	0/986	0.62	0/1335
All	All	0.46	1/12056 (0.0%)	0.67	3/16257 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	4
All	All	0	9

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	X	134	GLU	CG-CD	5.53	1.60	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	50	LEU	CA-CB-CG	6.01	129.13	115.30
1	B	50	LEU	CA-CB-CG	5.92	128.92	115.30
1	A	24	TYR	N-CA-CB	5.08	119.75	110.60

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	GLU	Peptide
1	A	271	ASN	Peptide
1	A	384	GLN	Peptide
1	A	531	GLU	Peptide
1	A	536	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4923	0	4960	296	0
1	B	4997	0	5037	316	0
2	X	962	0	936	62	0
2	Y	962	0	936	66	0
3	A	31	0	12	5	0
3	B	31	0	12	13	0
All	All	11906	0	11893	701	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 30.

The worst 5 of 701 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:331:LEU:HB3	1:B:332:PHE:HA	1.18	1.17
2:Y:18:THR:HG23	2:Y:120:ASP:HB2	1.18	1.17
1:B:117:ILE:HB	1:B:346:ARG:HD2	1.21	1.13
1:A:475:THR:HG23	1:A:588:ASN:HD21	1.00	1.11
2:X:18:THR:HG23	2:X:120:ASP:HB2	1.16	1.11

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	601/677 (89%)	447 (74%)	107 (18%)	47 (8%)	1	6
1	B	612/677 (90%)	447 (73%)	117 (19%)	48 (8%)	1	6
2	X	109/134 (81%)	83 (76%)	18 (16%)	8 (7%)	1	7
2	Y	109/134 (81%)	82 (75%)	19 (17%)	8 (7%)	1	7
All	All	1431/1622 (88%)	1059 (74%)	261 (18%)	111 (8%)	1	6

5 of 111 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	23	VAL
1	A	24	TYR
1	A	25	SER
1	A	98	LYS
1	A	164	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	546/615 (89%)	454 (83%)	92 (17%)	2	11
1	B	554/615 (90%)	457 (82%)	97 (18%)	2	10
2	X	106/124 (86%)	84 (79%)	22 (21%)	1	6
2	Y	106/124 (86%)	85 (80%)	21 (20%)	1	7
All	All	1312/1478 (89%)	1080 (82%)	232 (18%)	2	10

5 of 232 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	104	PHE
1	B	274	LYS
2	Y	24	ASN
1	B	113	ASP
1	B	160	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	467	HIS
1	A	588	ASN
1	B	561	GLN
1	A	561	GLN
1	B	72	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ATP	A	1678	1	24,33,33	1.12	2 (8%)	31,52,52	2.21	6 (19%)
3	ATP	B	1678	-	24,33,33	0.98	1 (4%)	31,52,52	2.27	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	A	1678	1	-	0/18/38/38	0/3/3/3
3	ATP	B	1678	-	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1678	ATP	O4'-C1'	2.01	1.43	1.41
3	B	1678	ATP	C5-C4	2.93	1.47	1.40
3	A	1678	ATP	C5-C4	3.63	1.48	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1678	ATP	N3-C2-N1	-6.79	123.69	128.89
3	A	1678	ATP	N3-C2-N1	-6.66	123.80	128.89
3	A	1678	ATP	PA-O3A-PB	-6.45	114.63	132.73
3	B	1678	ATP	PA-O3A-PB	-6.03	115.80	132.73
3	B	1678	ATP	PB-O3B-PG	-5.49	114.27	132.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1678	ATP	5	0
3	B	1678	ATP	13	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	609/677 (89%)	1.04	92 (15%) 3 1	63, 72, 74, 79	0
1	B	618/677 (91%)	0.86	71 (11%) 6 2	66, 72, 75, 88	0
2	X	115/134 (85%)	0.82	17 (14%) 3 1	69, 71, 74, 82	0
2	Y	115/134 (85%)	0.94	17 (14%) 3 1	69, 71, 74, 83	0
All	All	1457/1622 (89%)	0.94	197 (13%) 4 2	63, 72, 74, 88	0

The worst 5 of 197 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	538	ALA	9.0
1	A	25	SER	7.3
1	B	673	SER	7.0
1	A	532	GLU	6.8
1	A	549	PRO	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ATP	A	1678	31/31	0.94	0.19	-2.15	21,30,32,33	0
3	ATP	B	1678	31/31	0.94	0.20	-2.44	18,29,31,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.