



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:31 PM GMT

PDB ID : 5CH1
Title : Crystal structure of an active polycomb repressive complex 2 in the stimulated state
Authors : Jiao, L.; Liu, X.
Deposited on : 2015-07-10
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5193 (2.30-2.26)
Clashscore	102246	5929 (2.30-2.26)
Ramachandran outliers	100387	5851 (2.30-2.26)
Sidechain outliers	100360	5850 (2.30-2.26)
RSRZ outliers	91569	5204 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	605	<div> <div></div> <div> <div></div> <div>73%</div> <div>6%</div> <div>20%</div> </div> </div>
2	B	937	<div> <div>6%</div> <div> <div></div> <div>76%</div> <div>9%</div> <div>15%</div> </div> </div>
3	D	11	<div> <div>9%</div> <div> <div></div> <div>73%</div> <div>27%</div> </div> </div>
4	E	20	<div> <div>10%</div> <div> <div></div> <div>40%</div> <div>5%</div> <div>55%</div> </div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative polycomb protein Eed.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	481	Total	C	N	O	S	0	0	0
			3782	2417	647	699	19			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	MET	-	initiating methionine	UNP G0S8H7
A	-38	ALA	-	expression tag	UNP G0S8H7
A	-37	SER	-	expression tag	UNP G0S8H7
A	-36	ALA	-	expression tag	UNP G0S8H7
A	-35	TRP	-	expression tag	UNP G0S8H7
A	-34	SER	-	expression tag	UNP G0S8H7
A	-33	HIS	-	expression tag	UNP G0S8H7
A	-32	PRO	-	expression tag	UNP G0S8H7
A	-31	GLN	-	expression tag	UNP G0S8H7
A	-30	PHE	-	expression tag	UNP G0S8H7
A	-29	GLU	-	expression tag	UNP G0S8H7
A	-28	LYS	-	expression tag	UNP G0S8H7
A	-27	GLY	-	expression tag	UNP G0S8H7
A	-26	GLY	-	expression tag	UNP G0S8H7
A	-25	GLY	-	expression tag	UNP G0S8H7
A	-24	SER	-	expression tag	UNP G0S8H7
A	-23	GLY	-	expression tag	UNP G0S8H7
A	-22	GLY	-	expression tag	UNP G0S8H7
A	-21	GLY	-	expression tag	UNP G0S8H7
A	-20	SER	-	expression tag	UNP G0S8H7
A	-19	GLY	-	expression tag	UNP G0S8H7
A	-18	GLY	-	expression tag	UNP G0S8H7
A	-17	SER	-	expression tag	UNP G0S8H7
A	-16	ALA	-	expression tag	UNP G0S8H7
A	-15	TRP	-	expression tag	UNP G0S8H7
A	-14	SER	-	expression tag	UNP G0S8H7
A	-13	HIS	-	expression tag	UNP G0S8H7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	PRO	-	expression tag	UNP G0S8H7
A	-11	GLN	-	expression tag	UNP G0S8H7
A	-10	PHE	-	expression tag	UNP G0S8H7
A	-9	GLU	-	expression tag	UNP G0S8H7
A	-8	LYS	-	expression tag	UNP G0S8H7
A	-7	LEU	-	expression tag	UNP G0S8H7
A	-6	GLU	-	expression tag	UNP G0S8H7
A	-5	VAL	-	expression tag	UNP G0S8H7
A	-4	LEU	-	expression tag	UNP G0S8H7
A	-3	PHE	-	expression tag	UNP G0S8H7
A	-2	GLN	-	expression tag	UNP G0S8H7
A	-1	GLY	-	expression tag	UNP G0S8H7
A	0	PRO	-	expression tag	UNP G0S8H7

- Molecule 2 is a protein called putative polycomb protein Ezh2, putative polycomb protein Suz12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	797	Total	C	N	O	S	0	0	0
			6426	4058	1155	1173	40			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	182	SER	-	expression tag	UNP G0SDW4
B	183	ASN	-	expression tag	UNP G0SDW4
B	184	HIS	-	expression tag	UNP G0SDW4
B	185	HIS	-	expression tag	UNP G0SDW4
B	186	HIS	-	expression tag	UNP G0SDW4
B	187	HIS	-	expression tag	UNP G0SDW4
B	188	HIS	-	expression tag	UNP G0SDW4
B	189	HIS	-	expression tag	UNP G0SDW4
B	190	ALA	-	expression tag	UNP G0SDW4
B	2524	LEU	-	linker	UNP G0SDW4
B	2525	VAL	-	linker	UNP G0SDW4
B	2526	PRO	-	linker	UNP G0SDW4
B	2527	ARG	-	linker	UNP G0SDW4
B	2528	GLY	-	linker	UNP G0SDW4
B	2529	SER	-	linker	UNP G0SDW4

- Molecule 3 is a protein called 11-Mer peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	8	Total	C	N	O	0	0	0
			60	38	13	9			


- Molecule 4 is a protein called 20-Mer peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	E	9	Total	C	N	O	0	1	0
			59	35	12	12			

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	8	Total	Zn	0	0
			8	8		

- Molecule 6 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	244	Total	O	0	0
			244	244		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	277	Total 277	O 277	0	0
7	D	4	Total 4	O 4	0	0
7	E	3	Total 3	O 3	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: putative polycomb protein Eed




- Molecule 2: putative polycomb protein Ezh2, putative polycomb protein Suz12







● Molecule 3: 11-Mer peptide



● Molecule 4: 20-Mer peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.55Å 74.61Å 128.51Å 90.00° 107.70° 90.00°	Depositor
Resolution (Å)	47.32 – 2.27 45.84 – 2.27	Depositor EDS
% Data completeness (in resolution range)	90.7 (47.32-2.27) 90.7 (45.84-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.27Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.168 , 0.212 0.168 , 0.213	Depositor DCC
R_{free} test set	3147 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.9	EDS
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 61840 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10889	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SAH, M3L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/3894	0.68	0/5306
2	B	0.49	0/6570	0.65	0/8870
3	D	0.53	0/48	0.61	0/63
4	E	0.47	0/61	0.74	0/81
All	All	0.51	0/10573	0.66	0/14320

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3782	0	3634	20	0
2	B	6426	0	6304	41	0
3	D	60	0	70	0	0
4	E	59	0	57	5	0
5	B	8	0	0	0	0
6	B	26	0	19	1	0
7	A	244	0	0	0	0
7	B	277	0	0	0	0
7	D	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	E	3	0	0	0	0
All	All	10889	0	10084	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:ASP:OD1	1:A:258:ARG:HD3	1.80	0.81
2:B:866:TRP:HZ3	4:E:25:ALA:CB	2.04	0.70
2:B:738:GLN:HA	2:B:749:VAL:HG11	1.77	0.66
2:B:2622:GLU:HB2	2:B:2662:VAL:HG13	1.79	0.65
2:B:763:CYS:H	2:B:764:GLY:HA2	1.61	0.64
2:B:318:ASN:HB3	2:B:838:ARG:HH12	1.62	0.64
2:B:866:TRP:CZ3	4:E:25:ALA:HB3	2.34	0.63
2:B:253:ASP:HB3	2:B:255:ASN:OD1	1.99	0.62
1:A:189:ASP:HB2	1:A:207:HIS:HD2	1.66	0.61
2:B:866:TRP:CZ3	4:E:25:ALA:CB	2.85	0.60
2:B:866:TRP:HZ3	4:E:25:ALA:HB2	1.71	0.56
2:B:763:CYS:N	2:B:764:GLY:HA2	2.21	0.55
2:B:574:CYS:HA	2:B:625:CYS:HB3	1.88	0.54
2:B:601:MET:O	2:B:605:ILE:HG12	2.09	0.52
1:A:186:HIS:CG	1:A:190:LEU:HD21	2.45	0.51
2:B:614:GLU:OE2	2:B:629:HIS:HD2	1.93	0.51
2:B:611:LEU:HD13	2:B:2669:ILE:HG13	1.92	0.51
1:A:120:VAL:HB	1:A:134:LEU:HB2	1.94	0.50
1:A:323:CYS:O	1:A:326:GLN:HB2	2.13	0.49
2:B:866:TRP:HZ3	4:E:25:ALA:HB3	1.67	0.48
1:A:320:THR:O	1:A:320:THR:CG2	2.61	0.48
1:A:203:LEU:HG	1:A:250:PHE:CE2	2.49	0.47
2:B:789:LEU:HD11	2:B:901:LYS:HB3	1.96	0.47
1:A:217:PRO:HD3	1:A:230:ILE:HD11	1.96	0.47
2:B:756:ASP:HB3	2:B:759:LEU:HB2	1.97	0.47
1:A:316:ALA:HB1	1:A:511:ILE:HD11	1.97	0.46
2:B:2625:LEU:HD21	2:B:2657:LEU:HD13	1.98	0.45
2:B:632:LEU:HD12	2:B:637:LEU:HD12	1.98	0.45
2:B:2594:GLN:HA	2:B:2604:LYS:HD3	1.97	0.45
2:B:893:TYR:CE2	2:B:896:HIS:HA	2.52	0.45
2:B:505:THR:HA	2:B:533:GLU:HG2	1.99	0.45
1:A:189:ASP:HB2	1:A:207:HIS:CD2	2.49	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ILE:HB	1:A:85:GLN:HB3	1.98	0.45
1:A:220:PRO:HB2	1:A:222:GLU:HG2	1.99	0.44
2:B:597:THR:HA	2:B:600:TRP:CE2	2.52	0.44
2:B:2586:LEU:HD11	2:B:2616:ARG:HD3	1.98	0.44
2:B:614:GLU:HB3	2:B:625:CYS:SG	2.58	0.44
2:B:309:GLU:HG2	2:B:2580:ILE:HG12	2.00	0.43
2:B:343:TRP:O	2:B:347:LEU:HG	2.18	0.43
2:B:259:PRO:HB2	2:B:262:VAL:HG13	1.99	0.43
2:B:620:ILE:HD12	2:B:2675:LEU:HD13	2.01	0.43
1:A:334:TYR:O	1:A:341:PRO:HA	2.19	0.43
2:B:311:ARG:HH22	2:B:792:GLY:HA3	1.84	0.43
1:A:320:THR:HG23	1:A:323:CYS:SG	2.59	0.42
1:A:186:HIS:CD2	1:A:190:LEU:HD21	2.55	0.42
2:B:598:LEU:HA	2:B:617:VAL:HG21	2.00	0.42
2:B:853:VAL:HG13	6:B:8009:SAH:H4'	2.01	0.42
1:A:19:ILE:HG12	1:A:545:ARG:HG2	2.02	0.42
1:A:165:ILE:HB	1:A:179:CYS:HB3	2.03	0.41
2:B:720:CYS:HA	2:B:721:PRO:HD3	1.93	0.41
1:A:83:ILE:HB	2:B:288:TYR:CE2	2.55	0.41
2:B:590:TRP:CG	2:B:631:LYS:HD3	2.55	0.41
1:A:252:GLY:HA3	2:B:228:THR:HG23	2.01	0.41
2:B:2637:ALA:HB1	2:B:2678:ALA:HB2	2.03	0.41
2:B:887:ILE:HG13	2:B:916:PHE:HB3	2.03	0.41
2:B:321:LEU:HD13	2:B:842:ARG:HD2	2.03	0.41
2:B:406:PRO:HA	2:B:407:GLU:HA	1.85	0.40
1:A:98:SER:HA	1:A:112:CYS:O	2.22	0.40
2:B:512:PHE:CE1	2:B:2662:VAL:HG11	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/605 (78%)	463 (98%)	10 (2%)	0	100	100
2	B	765/937 (82%)	733 (96%)	29 (4%)	3 (0%)	39	47
3	D	5/11 (46%)	5 (100%)	0	0	100	100
4	E	8/20 (40%)	8 (100%)	0	0	100	100
All	All	1251/1573 (80%)	1209 (97%)	39 (3%)	3 (0%)	52	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	579	GLY
2	B	923	PRO
2	B	425	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/495 (82%)	397 (98%)	9 (2%)	60	75
2	B	699/816 (86%)	680 (97%)	19 (3%)	52	68
3	D	4/6 (67%)	4 (100%)	0	100	100
4	E	4/13 (31%)	4 (100%)	0	100	100
All	All	1113/1330 (84%)	1085 (98%)	28 (2%)	55	71

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	35	GLN
1	A	90	ASP
1	A	175	GLN
1	A	188	TYR
1	A	258	ARG
1	A	298	TYR
1	A	320	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	365	ARG
2	B	262	VAL
2	B	356	PHE
2	B	396	SER
2	B	493	MET
2	B	568	PRO
2	B	628	VAL
2	B	631	LYS
2	B	638	ARG
2	B	659	ARG
2	B	738	GLN
2	B	805	GLU
2	B	857	PHE
2	B	922	PHE
2	B	929	LEU
2	B	2567	VAL
2	B	2595	ASP
2	B	2644	ARG
2	B	2679	ARG
2	B	2683	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	175	GLN
2	B	629	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	M3L	D	27	3	9,11,12	0.38	0	12,14,16	0.70	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	M3L	D	27	3	-	0/8/10/12	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	27	M3L	O-C-CA	-2.21	119.78	125.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SAH	B	8009	-	22,28,28	0.56	0	18,40,40	0.81	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SAH	B	8009	-	-	0/7/31/31	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	8009	SAH	CB-CA-N	2.39	117.25	110.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	8009	SAH	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	481/605 (79%)	-0.04	7 (1%) 76 81	15, 28, 62, 82	0
2	B	797/937 (85%)	0.26	58 (7%) 18 24	18, 45, 88, 122	0
3	D	7/11 (63%)	0.37	1 (14%) 4 5	31, 37, 51, 76	0
4	E	9/20 (45%)	0.94	2 (22%) 1 1	33, 46, 76, 76	0
All	All	1294/1573 (82%)	0.15	68 (5%) 30 38	15, 38, 79, 122	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	628	VAL	5.0
2	B	491	ALA	4.9
2	B	451	GLN	4.5
2	B	473	PHE	4.4
2	B	262	VAL	4.3
2	B	577	ILE	4.1
2	B	401	PHE	4.1
2	B	276	TYR	4.0
2	B	926	THR	3.7
2	B	647	ILE	3.7
2	B	450	ASP	3.7
2	B	197	TRP	3.6
2	B	406	PRO	3.5
2	B	389	ASN	3.4
2	B	568	PRO	3.4
2	B	490	GLU	3.4
1	A	492	ARG	3.2
2	B	569	PRO	3.1
2	B	924	ASN	3.1
2	B	439	MET	3.0
2	B	405	GLN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	435	ARG	3.0
2	B	626	TRP	3.0
2	B	578	HIS	3.0
2	B	778	VAL	2.9
2	B	928	LYS	2.9
2	B	391	GLU	2.8
2	B	636	ASP	2.8
2	B	202	ILE	2.8
2	B	2680	GLY	2.8
2	B	443	ALA	2.8
2	B	630	ARG	2.8
2	B	493	MET	2.8
2	B	574	CYS	2.8
1	A	556	ARG	2.7
2	B	413	THR	2.7
2	B	199	VAL	2.6
2	B	338	GLN	2.6
1	A	35	GLN	2.6
2	B	356	PHE	2.5
2	B	275	GLU	2.5
2	B	2681	ARG	2.5
2	B	404	SER	2.4
2	B	649	LYS	2.4
2	B	2673	GLN	2.4
2	B	198	THR	2.4
2	B	777	GLU	2.4
2	B	2551	LEU	2.4
4	E	21	ALA	2.3
2	B	923	PRO	2.3
1	A	97	ALA	2.3
1	A	482	PRO	2.3
4	E	22	THR	2.2
2	B	648	PRO	2.2
2	B	2644	ARG	2.2
2	B	495	LYS	2.2
2	B	397	ASN	2.2
2	B	550	ILE	2.1
2	B	730	HIS	2.1
2	B	728	ALA	2.1
2	B	388	LEU	2.1
2	B	349	ASN	2.1
2	B	363	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	390	ILE	2.0
2	B	548	ALA	2.0
2	B	414	PRO	2.0
1	A	193	VAL	2.0
3	D	23	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	M3L	D	27	12/13	0.96	0.16	-	19,22,29,30	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	ZN	B	8001	1/1	1.00	0.12	0.27	33,33,33,33	0
6	SAH	B	8009	26/26	0.94	0.17	0.26	14,25,31,36	26
5	ZN	B	8002	1/1	1.00	0.12	-0.14	37,37,37,37	0
5	ZN	B	8006	1/1	1.00	0.10	-0.16	29,29,29,29	0
5	ZN	B	8007	1/1	1.00	0.13	-0.19	26,26,26,26	0
5	ZN	B	8003	1/1	0.99	0.11	-0.27	36,36,36,36	0
5	ZN	B	8004	1/1	0.99	0.10	-0.42	31,31,31,31	0
5	ZN	B	8005	1/1	1.00	0.09	-0.46	32,32,32,32	0
5	ZN	B	8008	1/1	0.98	0.04	-1.99	80,80,80,80	0

6.5 Other polymers

There are no such residues in this entry.