



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:35 PM GMT

PDB ID : 4CJ8
Title : monoclinic crystal form of Bogt6a E192Q in complex with UDP-GalNAc, UDP and GalNAc
Authors : Pham, T.; Stinson, B.; Thiagarajan, N.; Lizotte-Waniewski, M.; Brew, K.; Acharya, K.R.
Deposited on : 2013-12-19
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1051 (3.60-3.40)
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)
RSRZ outliers	91569	1058 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	<div> <div>73%</div> <div>22%</div> <div>• •</div> </div>
1	B	248	<div> <div>76%</div> <div>19%</div> <div>• •</div> </div>
1	C	248	<div> <div>72%</div> <div>23%</div> <div>•</div> </div>
1	D	248	<div> <div>75%</div> <div>19%</div> <div>• •</div> </div>
1	E	248	<div> <div>77%</div> <div>21%</div> <div>••</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	248	 79%18% ..
1	G	248	 78%19% ..
1	H	248	 79%21%
1	I	248	 77%17% • 5%
1	J	248	 72%20% • 5%
1	K	248	 73%21% • •
1	L	248	 72%21% • 5%
1	M	248	 77%21% ..
1	N	248	 %75%20% • •
1	O	248	 2%71%23% • •
1	P	248	 82%15% • •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	UDP	N	1244	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 32877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOSYLTRANSFERASE FAMILY 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	238	Total	C	N	O	S	10	0	0
			1989	1300	324	358	7			
1	B	238	Total	C	N	O	S	10	0	0
			1989	1300	324	358	7			
1	C	238	Total	C	N	O	S	10	0	0
			1989	1300	324	358	7			
1	D	237	Total	C	N	O	S	10	0	0
			1983	1297	323	356	7			
1	E	246	Total	C	N	O	S	10	0	0
			2057	1342	341	367	7			
1	F	246	Total	C	N	O	S	10	0	0
			2057	1342	341	367	7			
1	G	246	Total	C	N	O	S	10	0	0
			2057	1342	341	367	7			
1	H	247	Total	C	N	O	S	10	0	0
			2063	1345	342	369	7			
1	I	236	Total	C	N	O	S	10	0	0
			1973	1291	320	355	7			
1	J	236	Total	C	N	O	S	10	0	0
			1973	1291	320	355	7			
1	K	237	Total	C	N	O	S	10	0	0
			1983	1297	323	356	7			
1	L	236	Total	C	N	O	S	10	0	0
			1973	1291	320	355	7			
1	M	245	Total	C	N	O	S	10	0	0
			2047	1336	338	366	7			
1	N	243	Total	C	N	O	S	10	0	0
			2027	1324	332	364	7			
1	O	240	Total	C	N	O	S	10	0	0
			2000	1306	326	361	7			
1	P	243	Total	C	N	O	S	10	0	0
			2027	1324	332	364	7			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
A	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
A	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
B	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
B	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
B	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
C	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
C	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
C	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
D	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
D	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
D	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
E	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
E	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
E	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
F	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
F	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
F	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
G	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
G	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
G	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
H	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
H	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
H	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
I	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
I	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
I	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
J	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
J	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
J	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
K	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
K	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
K	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
L	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
L	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
L	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
M	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
M	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
M	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
N	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
N	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
N	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
O	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
O	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
O	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2
P	-1	SER	-	EXPRESSION TAG	UNP A7LVT2
P	0	HIS	-	EXPRESSION TAG	UNP A7LVT2
P	192	GLN	GLU	ENGINEERED MUTATION	UNP A7LVT2

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	G	1	Total C O 6 3 3	0	0
2	G	1	Total C O 6 3 3	0	0

- Molecule 3 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	B	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	C	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	D	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	I	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	J	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	K	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	L	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	M	1	Total	C	N	O	P	0	0
			25	9	2	12	2		
3	N	1	Total	C	N	O	P	0	0
			25	9	2	12	2		

- Molecule 4 is SUGAR (N-ACETYL-2-DEOXY-2-AMINO-GALACTOSE) (three-letter code: A2G) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			15	8	1	6		
4	B	1	Total	C	N	O	0	0
			15	8	1	6		
4	C	1	Total	C	N	O	0	0
			15	8	1	6		
4	D	1	Total	C	N	O	0	0
			15	8	1	6		
4	I	1	Total	C	N	O	0	0
			15	8	1	6		
4	J	1	Total	C	N	O	0	0
			15	8	1	6		
4	K	1	Total	C	N	O	0	0
			15	8	1	6		
4	L	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O_4S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is URIDINE-DIPHOSPHATE-N-ACETYLGALACTOSAMINE (three-letter code: UD2) (formula: $C_{17}H_{27}N_3O_{17}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	E	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
7	F	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
7	G	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
7	H	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
7	O	1	Total	C	N	O	P	0	0
			39	17	3	17	2		
7	P	1	Total	C	N	O	P	0	0
			39	17	3	17	2		

- Molecule 8 is SUGAR (N-ACETYL-D-GALACTOSAMINE) (three-letter code: NGA) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	M	1	Total	C	N	O	0	0
			14	8	1	5		
8	N	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	2	Total	O	0	0
			2	2		
9	C	1	Total	O	0	0
			1	1		
9	D	1	Total	O	0	0
			1	1		
9	E	1	Total	O	0	0
			1	1		
9	L	2	Total	O	0	0
			2	2		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




• Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




• Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




• Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6



- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain J:  72% 20% 5%




- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain K:  73% 21% 5%





- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain L:  72% 21% 5%





- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain M:  77% 21% 5%



- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain N:  75% 20% 5%



- Molecule 1: GLYCOSYLTRANSFERASE FAMILY 6

Chain O:  71% 23% 5%

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	176.69Å 79.74Å 179.08Å 90.00° 95.23° 90.00°	Depositor
Resolution (Å)	87.98 – 3.50 87.98 – 3.50	Depositor EDS
% Data completeness (in resolution range)	97.7 (87.98-3.50) 87.0 (87.98-3.50)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 3.49Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.4_1496)	Depositor
R, R_{free}	0.225 , 0.249 0.237 , 0.264	Depositor DCC
R_{free} test set	3137 reflections (5.77%)	DCC
Wilson B-factor (Å ²)	61.6	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 18.6	EDS
Estimated twinning fraction	0.087 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 61938 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	32877	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.90 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.9103e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: UD2, GOL, UDP, NGA, 1PE, SO4, A2G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/2055	0.35	0/2791
1	B	0.21	0/2055	0.35	0/2791
1	C	0.21	0/2055	0.35	0/2791
1	D	0.21	0/2049	0.36	0/2783
1	E	0.21	0/2124	0.36	0/2881
1	F	0.21	0/2124	0.37	0/2881
1	G	0.21	0/2124	0.36	0/2881
1	H	0.21	0/2130	0.35	0/2889
1	I	0.21	0/2038	0.36	0/2768
1	J	0.21	0/2038	0.36	0/2768
1	K	0.21	0/2049	0.35	0/2783
1	L	0.21	0/2038	0.35	0/2768
1	M	0.21	0/2113	0.36	0/2866
1	N	0.21	0/2093	0.35	0/2841
1	O	0.25	0/2066	0.37	0/2805
1	P	0.21	0/2093	0.36	0/2841
All	All	0.21	0/33244	0.36	0/45128

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1989	0	1926	37	0
1	B	1989	0	1926	34	0
1	C	1989	0	1926	33	0
1	D	1983	0	1921	28	0
1	E	2057	0	2001	31	0
1	F	2057	0	2001	29	0
1	G	2057	0	2001	37	0
1	H	2063	0	2006	32	0
1	I	1973	0	1914	28	0
1	J	1973	0	1914	38	0
1	K	1983	0	1921	40	0
1	L	1973	0	1914	35	1
1	M	2047	0	1992	33	0
1	N	2027	0	1966	44	0
1	O	2000	0	1933	64	0
1	P	2027	0	1968	34	0
2	A	12	0	16	0	0
2	D	6	0	8	0	0
2	G	12	0	16	1	0
3	A	25	0	11	4	0
3	B	25	0	11	5	0
3	C	25	0	11	3	1
3	D	25	0	11	6	0
3	I	25	0	11	2	0
3	J	25	0	11	2	0
3	K	25	0	11	4	0
3	L	25	0	11	2	0
3	M	25	0	11	4	0
3	N	25	0	11	18	0
4	A	15	0	15	2	0
4	B	15	0	15	0	0
4	C	15	0	15	0	0
4	D	15	0	15	0	0
4	I	15	0	15	0	0
4	J	15	0	15	0	0
4	K	15	0	15	2	0
4	L	15	0	15	5	0
5	A	16	0	22	1	0
6	B	5	0	0	0	0
7	E	39	0	25	5	0
7	F	39	0	25	5	0
7	G	39	0	25	7	0
7	H	39	0	25	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	O	39	0	25	20	0
7	P	39	0	25	3	0
8	M	14	0	13	2	0
8	N	14	0	13	2	0
9	B	2	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
9	E	1	0	0	0	0
9	L	2	0	0	0	0
All	All	32877	0	31698	576	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 576 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:N:95:ASN:ND2	3:N:1244:UDP:H5'2	1.74	1.02
1:D:191:ASP:OD1	1:D:192:GLN:HG3	1.63	0.98
1:C:231:LYS:NZ	3:C:1237:UDP:O2A	1.96	0.97
1:G:95:ASN:ND2	1:G:96:ALA:H	1.63	0.96
1:F:95:ASN:HD22	1:F:96:ALA:H	0.99	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:184:HIS:CE1	3:C:1237:UDP:O3B[1_654]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/248 (95%)	219 (93%)	17 (7%)	0	100	100
1	B	236/248 (95%)	221 (94%)	15 (6%)	0	100	100
1	C	236/248 (95%)	219 (93%)	17 (7%)	0	100	100
1	D	235/248 (95%)	220 (94%)	14 (6%)	1 (0%)	39	81
1	E	244/248 (98%)	231 (95%)	13 (5%)	0	100	100
1	F	244/248 (98%)	227 (93%)	17 (7%)	0	100	100
1	G	244/248 (98%)	231 (95%)	12 (5%)	1 (0%)	39	81
1	H	245/248 (99%)	230 (94%)	15 (6%)	0	100	100
1	I	234/248 (94%)	221 (94%)	13 (6%)	0	100	100
1	J	234/248 (94%)	218 (93%)	16 (7%)	0	100	100
1	K	235/248 (95%)	216 (92%)	19 (8%)	0	100	100
1	L	234/248 (94%)	218 (93%)	16 (7%)	0	100	100
1	M	243/248 (98%)	233 (96%)	10 (4%)	0	100	100
1	N	241/248 (97%)	227 (94%)	13 (5%)	1 (0%)	39	81
1	O	238/248 (96%)	222 (93%)	15 (6%)	1 (0%)	39	81
1	P	241/248 (97%)	224 (93%)	17 (7%)	0	100	100
All	All	3820/3968 (96%)	3577 (94%)	239 (6%)	4 (0%)	56	90

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	237	GLY
1	N	146	PRO
1	D	224	PRO
1	O	45	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/223 (96%)	209 (97%)	6 (3%)	51	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	215/223 (96%)	211 (98%)	4 (2%)	65	87
1	C	215/223 (96%)	209 (97%)	6 (3%)	51	82
1	D	214/223 (96%)	209 (98%)	5 (2%)	58	85
1	E	221/223 (99%)	212 (96%)	9 (4%)	37	74
1	F	221/223 (99%)	209 (95%)	12 (5%)	27	67
1	G	221/223 (99%)	213 (96%)	8 (4%)	42	77
1	H	222/223 (100%)	217 (98%)	5 (2%)	58	85
1	I	213/223 (96%)	208 (98%)	5 (2%)	58	85
1	J	213/223 (96%)	205 (96%)	8 (4%)	40	76
1	K	214/223 (96%)	206 (96%)	8 (4%)	41	76
1	L	213/223 (96%)	203 (95%)	10 (5%)	32	72
1	M	220/223 (99%)	211 (96%)	9 (4%)	37	74
1	N	218/223 (98%)	208 (95%)	10 (5%)	33	72
1	O	215/223 (96%)	205 (95%)	10 (5%)	32	72
1	P	218/223 (98%)	209 (96%)	9 (4%)	37	74
All	All	3468/3568 (97%)	3344 (96%)	124 (4%)	42	77

5 of 124 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	95	ASN
1	K	62	LYS
1	O	233	LYS
1	I	125	PHE
1	J	95	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	192	GLN
1	I	95	ASN
1	O	95	ASN
1	H	95	ASN
1	H	192	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

33 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	1236	-	5,5,5	0.36	0	5,5,5	0.19	0
3	UDP	A	1237	-	18,26,26	0.72	0	26,40,40	1.76	2 (7%)
4	A2G	A	1238	-	15,15,15	0.38	0	17,21,21	0.57	0
2	GOL	A	1239	-	5,5,5	0.35	0	5,5,5	0.23	0
5	1PE	A	1240	-	15,15,15	0.54	0	14,14,14	0.47	0
3	UDP	B	1237	-	18,26,26	0.72	0	26,40,40	1.69	2 (7%)
4	A2G	B	1238	-	15,15,15	0.38	0	17,21,21	0.57	0
6	SO4	B	1239	-	4,4,4	0.24	0	6,6,6	0.08	0
3	UDP	C	1237	-	18,26,26	0.72	0	26,40,40	1.75	2 (7%)
4	A2G	C	1238	-	15,15,15	0.38	0	17,21,21	0.55	0
3	UDP	D	1237	-	18,26,26	0.73	0	26,40,40	1.76	2 (7%)
4	A2G	D	1238	-	15,15,15	0.39	0	17,21,21	0.48	0
2	GOL	D	1239	-	5,5,5	0.35	0	5,5,5	0.23	0
7	UD2	E	1246	-	32,41,41	0.55	0	46,62,62	1.36	2 (4%)
7	UD2	F	1246	-	32,41,41	0.54	0	46,62,62	1.36	2 (4%)
7	UD2	G	1246	-	32,41,41	0.53	0	46,62,62	1.37	2 (4%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	G	1247	-	5,5,5	0.34	0	5,5,5	0.20	0
2	GOL	G	1248	-	5,5,5	0.35	0	5,5,5	0.21	0
7	UD2	H	1246	-	32,41,41	0.55	0	46,62,62	1.37	3 (6%)
3	UDP	I	1237	-	18,26,26	0.71	0	26,40,40	1.71	2 (7%)
4	A2G	I	1238	-	15,15,15	0.39	0	17,21,21	0.51	0
3	UDP	J	1237	-	18,26,26	0.71	0	26,40,40	1.63	2 (7%)
4	A2G	J	1238	-	15,15,15	0.39	0	17,21,21	0.48	0
3	UDP	K	1237	-	18,26,26	0.72	0	26,40,40	1.77	2 (7%)
4	A2G	K	1238	-	15,15,15	0.39	0	17,21,21	0.46	0
3	UDP	L	1237	-	18,26,26	0.72	0	26,40,40	1.65	2 (7%)
4	A2G	L	1238	-	15,15,15	0.39	0	17,21,21	0.49	0
3	UDP	M	1246	-	18,26,26	0.72	0	26,40,40	1.75	2 (7%)
8	NGA	M	1247	1	14,14,15	0.52	0	15,19,21	0.64	0
3	UDP	N	1244	-	18,26,26	0.72	0	26,40,40	1.76	2 (7%)
8	NGA	N	1245	1	14,14,15	0.51	0	15,19,21	0.64	0
7	UD2	O	1241	-	32,41,41	0.56	0	46,62,62	1.40	2 (4%)
7	UD2	P	1244	-	32,41,41	0.54	0	46,62,62	1.38	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1236	-	-	0/4/4/4	0/0/0/0
3	UDP	A	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	A	1238	-	-	0/6/26/26	0/1/1/1
2	GOL	A	1239	-	-	0/4/4/4	0/0/0/0
5	1PE	A	1240	-	-	0/13/13/13	0/0/0/0
3	UDP	B	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	B	1238	-	-	0/6/26/26	0/1/1/1
6	SO4	B	1239	-	-	0/0/0/0	0/0/0/0
3	UDP	C	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	C	1238	-	-	0/6/26/26	0/1/1/1
3	UDP	D	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	D	1238	-	-	0/6/26/26	0/1/1/1
2	GOL	D	1239	-	-	0/4/4/4	0/0/0/0
7	UD2	E	1246	-	-	0/22/63/63	0/3/3/3
7	UD2	F	1246	-	-	0/22/63/63	0/3/3/3
7	UD2	G	1246	-	-	0/22/63/63	0/3/3/3
2	GOL	G	1247	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	G	1248	-	-	0/4/4/4	0/0/0/0
7	UD2	H	1246	-	-	0/22/63/63	0/3/3/3
3	UDP	I	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	I	1238	-	-	1/6/26/26	0/1/1/1
3	UDP	J	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	J	1238	-	-	0/6/26/26	0/1/1/1
3	UDP	K	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	K	1238	-	-	0/6/26/26	0/1/1/1
3	UDP	L	1237	-	-	0/12/32/32	0/2/2/2
4	A2G	L	1238	-	-	1/6/26/26	0/1/1/1
3	UDP	M	1246	-	-	0/12/32/32	0/2/2/2
8	NGA	M	1247	1	-	0/6/23/26	0/1/1/1
3	UDP	N	1244	-	-	0/12/32/32	0/2/2/2
8	NGA	N	1245	1	-	0/6/23/26	0/1/1/1
7	UD2	O	1241	-	-	0/22/63/63	0/3/3/3
7	UD2	P	1244	-	-	0/22/63/63	0/3/3/3

There are no bond length outliers.

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	O	1241	UD2	PB-O3A-PA	-4.41	120.33	132.73
3	A	1237	UDP	PA-O3A-PB	-3.80	119.92	132.67
3	D	1237	UDP	PA-O3A-PB	-3.70	120.27	132.67
3	N	1244	UDP	PA-O3A-PB	-3.68	120.33	132.67
3	C	1237	UDP	PA-O3A-PB	-3.67	120.36	132.67

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	1238	A2G	O7-C7-N2-C2
4	I	1238	A2G	O7-C7-N2-C2

There are no ring outliers.

23 monomers are involved in 106 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1237	UDP	4	0
4	A	1238	A2G	2	0
5	A	1240	1PE	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1237	UDP	5	0
3	C	1237	UDP	3	1
3	D	1237	UDP	6	0
7	E	1246	UD2	5	0
7	F	1246	UD2	5	0
7	G	1246	UD2	7	0
2	G	1247	GOL	1	0
7	H	1246	UD2	4	0
3	I	1237	UDP	2	0
3	J	1237	UDP	2	0
3	K	1237	UDP	4	0
4	K	1238	A2G	2	0
3	L	1237	UDP	2	0
4	L	1238	A2G	5	0
3	M	1246	UDP	4	0
8	M	1247	NGA	2	0
3	N	1244	UDP	18	0
8	N	1245	NGA	2	0
7	O	1241	UD2	20	0
7	P	1244	UD2	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/248 (95%)	-0.41	0 100 100	35, 52, 66, 92	3 (1%)
1	B	238/248 (95%)	-0.36	0 100 100	41, 63, 81, 102	3 (1%)
1	C	238/248 (95%)	-0.32	0 100 100	46, 63, 82, 100	3 (1%)
1	D	237/248 (95%)	-0.30	0 100 100	46, 64, 82, 89	3 (1%)
1	E	246/248 (99%)	-0.39	0 100 100	38, 56, 84, 94	3 (1%)
1	F	246/248 (99%)	-0.34	0 100 100	47, 62, 88, 99	3 (1%)
1	G	246/248 (99%)	-0.39	0 100 100	43, 59, 79, 92	3 (1%)
1	H	247/248 (99%)	-0.29	0 100 100	43, 66, 94, 104	3 (1%)
1	I	236/248 (95%)	-0.24	0 100 100	41, 66, 88, 109	3 (1%)
1	J	236/248 (95%)	-0.10	0 100 100	59, 84, 99, 109	3 (1%)
1	K	237/248 (95%)	-0.27	0 100 100	41, 70, 85, 111	3 (1%)
1	L	236/248 (95%)	-0.19	1 (0%) 93 90	51, 79, 102, 111	3 (1%)
1	M	245/248 (98%)	-0.21	0 100 100	42, 66, 85, 102	3 (1%)
1	N	243/248 (97%)	0.01	3 (1%) 81 72	53, 94, 110, 120	3 (1%)
1	O	240/248 (96%)	0.30	6 (2%) 61 50	66, 106, 126, 140	3 (1%)
1	P	243/248 (97%)	-0.07	1 (0%) 93 90	50, 81, 98, 113	3 (1%)
All	All	3852/3968 (97%)	-0.22	11 (0%) 94 91	35, 68, 105, 140	48 (1%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	O	160	GLY	5.6
1	O	186	ILE	4.1
1	O	187	PRO	3.1
1	L	32	GLN	2.6
1	P	117	LEU	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	1PE	A	1240	16/16	0.83	0.27	1.42	68,77,84,86	0
4	A2G	L	1238	15/15	0.88	0.29	0.95	60,67,74,75	0
8	NGA	M	1247	14/15	0.90	0.27	0.89	84,93,100,104	0
4	A2G	I	1238	15/15	0.82	0.25	0.18	56,76,86,94	0
3	UDP	B	1237	25/25	0.89	0.26	0.16	64,78,111,133	0
4	A2G	A	1238	15/15	0.86	0.24	0.11	44,53,68,73	0
3	UDP	C	1237	25/25	0.89	0.26	0.01	61,78,107,131	0
3	UDP	A	1237	25/25	0.89	0.23	-0.10	51,57,90,100	0
3	UDP	D	1237	25/25	0.94	0.23	-0.16	66,73,101,104	0
6	SO4	B	1239	5/5	0.76	0.24	-0.28	98,98,105,119	0
7	UD2	P	1244	39/39	0.92	0.25	-0.31	70,80,93,94	0
8	NGA	N	1245	14/15	0.91	0.25	-0.34	84,93,100,104	0
7	UD2	F	1246	39/39	0.93	0.23	-0.34	48,61,77,79	0
7	UD2	H	1246	39/39	0.93	0.24	-0.39	52,61,72,84	0
3	UDP	L	1237	25/25	0.91	0.20	-0.65	73,86,101,108	0
7	UD2	O	1241	39/39	0.85	0.25	-0.72	87,114,125,131	0
4	A2G	C	1238	15/15	0.86	0.22	-0.73	52,68,77,78	0
7	UD2	E	1246	39/39	0.93	0.20	-0.80	48,59,75,79	0
4	A2G	B	1238	15/15	0.88	0.22	-0.83	42,55,65,66	0
3	UDP	N	1244	25/25	0.88	0.21	-0.85	94,104,109,113	0
3	UDP	M	1246	25/25	0.94	0.21	-0.88	66,71,77,82	0
4	A2G	J	1238	15/15	0.89	0.20	-0.93	50,70,76,78	0
4	A2G	K	1238	15/15	0.85	0.24	-0.97	69,81,85,89	0
7	UD2	G	1246	39/39	0.94	0.21	-1.18	50,58,70,78	0
3	UDP	I	1237	25/25	0.93	0.18	-1.36	63,70,91,112	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	UDP	J	1237	25/25	0.86	0.20	-1.41	86,102,113,122	0
4	A2G	D	1238	15/15	0.91	0.20	-1.49	69,75,81,81	0
2	GOL	A	1239	6/6	0.92	0.15	-1.65	53,57,59,61	0
3	UDP	K	1237	25/25	0.94	0.16	-1.93	56,74,91,101	0
2	GOL	G	1247	6/6	0.93	0.16	-	66,71,72,81	0
2	GOL	D	1239	6/6	0.90	0.19	-	55,65,68,76	0
2	GOL	G	1248	6/6	0.91	0.25	-	54,67,71,75	0
2	GOL	A	1236	6/6	0.90	0.20	-	55,65,67,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.