



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:17 AM GMT

PDB ID : 2CKP  
Title : CRYSTAL STRUCTURE OF HUMAN CHOLINE KINASE ALPHA-2 IN  
COMPLEX WITH ADP  
Authors : Malito, E.; Lavie, A.  
Deposited on : 2006-04-20  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	390	 2% 59% 19% • 19%
1	B	390	 2% 47% 20% • • 30%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4679 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CHOLINE KINASE ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2533	1650	419	448	16			
1	B	274	Total	C	N	O	S	0	0	0
			2093	1358	350	372	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	220	SER	GLY	CONFLICT	UNP P35790
A	422	LEU	GLN	CONFLICT	UNP P35790
B	220	SER	GLY	CONFLICT	UNP P35790
B	422	LEU	GLN	CONFLICT	UNP P35790

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		
3	B	11	Total	O	0	0
			11	11		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

Chain B:

2%

47%

20%

30%

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.27Å 127.14Å 158.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 3.10 14.99 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (15.00-3.10) 99.3 (14.99-3.10)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.65 (at 3.12Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.256 , 0.329 0.255 , 0.328	Depositor DCC
$R_{free}$ test set	1990 reflections (11.31%)	DCC
Wilson B-factor (Å <sup>2</sup> )	68.6	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 86.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 19580 reflections	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	4679	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	0/2602	0.66	2/3512 (0.1%)
1	B	0.46	0/2145	0.62	2/2906 (0.1%)
All	All	0.51	0/4747	0.64	4/6418 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	315	LEU	CA-CB-CG	6.27	129.71	115.30
1	B	130	PRO	N-CA-CB	5.79	110.25	103.30
1	A	214	LEU	CA-CB-CG	5.72	128.46	115.30
1	A	179	LEU	CA-CB-CG	5.49	127.92	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	282	LEU	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2533	0	2408	57	0
1	B	2093	0	1887	53	0
2	A	27	0	12	0	0
3	A	15	0	0	1	0
3	B	11	0	0	2	0
All	All	4679	0	4307	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:92:TYR:HA	1:B:93:LEU:C	1.83	0.97
1:A:308:GLN:HB3	1:A:349:GLU:HG3	1.48	0.92
1:A:308:GLN:HB3	1:A:349:GLU:CG	2.11	0.81
1:B:345:ASN:HD21	1:B:420:TRP:HE1	1.33	0.77
1:A:380:SER:HB3	1:A:402:LYS:NZ	2.03	0.74
1:A:380:SER:HB3	1:A:402:LYS:HZ1	1.54	0.73
1:B:400:ILE:O	1:B:404:GLU:HB2	1.89	0.71
1:B:282:LEU:O	1:B:284:LEU:N	2.24	0.70
1:B:398:LYS:O	1:B:402:LYS:HG2	1.92	0.69
1:B:146:ARG:HH11	1:B:204:ARG:HH21	1.41	0.68
1:A:129:LEU:H	1:A:130:PRO:HD2	1.60	0.67
1:A:129:LEU:N	1:A:130:PRO:HD2	2.13	0.64
1:A:276:LYS:HE2	1:A:442:GLN:HE22	1.61	0.64
1:A:245:GLU:O	1:A:247:LYS:HG3	1.98	0.63
1:B:308:GLN:HE21	1:B:310:GLY:H	1.46	0.63
1:B:451:GLN:O	1:B:455:LEU:HB2	1.99	0.63
1:A:214:LEU:O	1:A:309:GLU:HB2	1.99	0.62
1:A:391:GLU:OE1	1:A:391:GLU:HA	2.00	0.62
1:A:250:PHE:HB3	1:A:290:ARG:NH2	2.15	0.62
1:A:183:MET:SD	1:A:240:MET:SD	2.99	0.60
1:A:247:LYS:HD2	1:A:248:TRP:N	2.16	0.59
1:B:211:SER:CB	1:B:315:LEU:HA	2.33	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:211:SER:HB3	1:B:315:LEU:HA	1.85	0.58
1:A:292:LEU:HD21	1:A:411:ARG:HD3	1.86	0.58
1:B:347:PHE:CE2	1:B:375:GLN:HB2	2.38	0.58
1:A:356:TYR:HB3	1:A:361:PHE:HA	1.87	0.57
1:B:96:LYS:NZ	1:B:96:LYS:HB2	2.20	0.56
1:B:228:ILE:HD13	1:B:312:ILE:HD13	1.86	0.56
1:B:396:GLU:O	1:B:400:ILE:HG12	2.05	0.56
1:B:92:TYR:CB	1:B:95:CYS:HB2	2.35	0.56
1:A:250:PHE:HB3	1:A:290:ARG:HH21	1.70	0.55
1:A:241:PRO:HA	1:B:197:TYR:O	2.07	0.55
1:B:108:GLU:CB	3:B:2002:HOH:O	2.54	0.55
1:A:304:HIS:CD2	1:A:328:LEU:HD21	2.41	0.55
1:B:284:LEU:HA	1:B:287:GLU:HG2	1.90	0.54
1:A:125:PHE:HB2	1:A:145:LEU:HB3	1.90	0.53
1:A:423:TRP:O	1:A:427:GLN:HG2	2.07	0.53
1:B:116:ILE:HG23	1:B:124:LEU:HB3	1.91	0.53
1:A:276:LYS:O	1:A:276:LYS:HE3	2.08	0.52
1:A:293:LEU:HD11	1:A:415:ALA:HB2	1.91	0.52
1:A:292:LEU:CD2	1:A:411:ARG:HD3	2.39	0.52
1:A:247:LYS:HD2	1:A:248:TRP:H	1.73	0.52
1:B:345:ASN:ND2	1:B:420:TRP:HE1	2.06	0.52
1:A:274:LEU:HG	1:A:278:LEU:HD12	1.91	0.52
1:B:309:GLU:HG2	1:B:310:GLY:N	2.25	0.51
1:B:146:ARG:NH1	1:B:204:ARG:HH21	2.08	0.51
1:A:146:ARG:O	1:A:203:GLY:HA3	2.10	0.51
1:A:197:TYR:O	1:A:205:LEU:HB2	2.11	0.51
1:A:379:ILE:HD12	1:A:405:MET:SD	2.51	0.50
1:A:194:PRO:HB3	1:A:209:ILE:HD11	1.91	0.50
1:B:383:LEU:HD21	1:B:405:MET:HE1	1.95	0.49
1:B:236:HIS:HD1	1:B:382:TYR:HE1	1.60	0.48
1:B:125:PHE:HB2	1:B:145:LEU:HB3	1.95	0.48
1:A:184:PHE:CZ	1:A:194:PRO:HG2	2.48	0.48
1:A:383:LEU:HD21	1:A:405:MET:SD	2.53	0.48
1:B:403:GLU:HG3	3:B:2010:HOH:O	2.13	0.48
1:B:315:LEU:HD22	1:B:325:LYS:O	2.13	0.48
1:B:375:GLN:O	1:B:379:ILE:HG23	2.14	0.48
1:A:414:LEU:HD21	1:A:452:LYS:HB3	1.95	0.48
1:A:183:MET:SD	1:A:240:MET:CE	3.03	0.47
1:B:253:MET:HG3	1:B:419:LEU:HD13	1.96	0.47
1:B:365:ASN:HB3	1:B:368:LYS:HG2	1.97	0.47
1:A:343:ILE:O	1:A:346:HIS:N	2.47	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:HIS:HD2	1:A:328:LEU:CD2	2.28	0.47
1:A:276:LYS:HE2	1:A:442:GLN:NE2	2.29	0.46
1:B:253:MET:HG3	1:B:419:LEU:CD1	2.45	0.46
1:B:219:LEU:HD22	1:B:346:HIS:HE1	1.81	0.46
1:B:328:LEU:HD12	1:B:328:LEU:H	1.80	0.46
1:B:92:TYR:HA	1:B:93:LEU:O	2.12	0.45
1:A:183:MET:SD	1:A:240:MET:HE1	2.56	0.45
1:B:96:LYS:C	1:B:96:LYS:HD3	2.36	0.45
1:A:308:GLN:HB3	1:A:349:GLU:CD	2.36	0.45
1:B:195:LYS:H	1:B:207:GLN:HB2	1.82	0.45
1:A:186:ILE:HG12	1:B:189:GLU:HG3	1.99	0.45
1:A:390:PHE:HA	1:A:393:LEU:HD12	1.99	0.44
1:A:240:MET:HE2	1:A:242:PHE:CE2	2.52	0.44
1:A:100:PRO:HG3	1:B:244:LYS:HE2	1.99	0.44
1:A:304:HIS:CD2	1:A:328:LEU:CD2	3.00	0.44
1:B:449:PHE:O	1:B:453:ARG:HD3	2.17	0.44
1:B:195:LYS:N	1:B:207:GLN:HB2	2.33	0.44
1:B:227:GLU:O	1:B:230:GLU:HG3	2.18	0.44
1:B:369:TYR:CD1	1:B:370:PRO:HD2	2.53	0.43
1:A:351:MET:HE2	1:A:369:TYR:HB2	2.00	0.43
1:A:427:GLN:O	1:A:431:SER:HB3	2.18	0.43
1:B:92:TYR:CA	1:B:93:LEU:C	2.72	0.43
1:A:240:MET:HB3	1:A:241:PRO:HD2	1.99	0.43
1:A:99:LEU:O	1:A:103:TRP:HB2	2.19	0.43
1:A:93:LEU:O	1:A:97:GLU:HG3	2.19	0.42
1:A:416:SER:O	1:A:420:TRP:CD1	2.72	0.42
1:B:376:LEU:CD2	1:B:406:LEU:HD21	2.49	0.42
1:B:185:ALA:O	1:B:189:GLU:HG2	2.20	0.42
1:B:278:LEU:HB3	1:B:279:SER:H	1.56	0.42
1:B:92:TYR:HA	1:B:94:TRP:N	2.31	0.42
1:A:129:LEU:N	1:A:130:PRO:CD	2.83	0.42
1:B:369:TYR:CG	1:B:370:PRO:HD2	2.55	0.42
1:A:101:GLY:HA3	3:A:2001:HOH:O	2.20	0.42
1:B:372:LYS:O	1:B:376:LEU:HB2	2.20	0.41
1:B:405:MET:O	1:B:409:VAL:HG23	2.20	0.41
1:B:359:TYR:CD1	1:B:360:PRO:HA	2.56	0.41
1:A:129:LEU:O	1:A:130:PRO:C	2.59	0.41
1:A:416:SER:O	1:A:420:TRP:HD1	2.04	0.41
1:B:245:GLU:HA	1:B:246:PRO:HD2	1.97	0.41
1:A:233:ALA:HB1	1:A:386:PHE:HB2	2.03	0.41
1:A:337:ASN:OD1	1:A:338:TYR:N	2.53	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:LYS:HA	1:A:239:LYS:HD3	1.74	0.40
1:A:244:LYS:HG3	1:A:244:LYS:HZ2	1.81	0.40
1:B:216:THR:OG1	1:B:353:ASP:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/390 (78%)	261 (85%)	42 (14%)	3 (1%)	19	58
1	B	258/390 (66%)	210 (81%)	39 (15%)	9 (4%)	4	24
All	All	564/780 (72%)	471 (84%)	81 (14%)	12 (2%)	9	37

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	ALA
1	B	93	LEU
1	B	110	GLU
1	B	282	LEU
1	B	283	PRO
1	B	356	TYR
1	B	94	TRP
1	B	306	ASP
1	B	394	SER
1	B	201	PRO
1	A	129	LEU
1	A	433	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/349 (74%)	236 (91%)	23 (9%)	12	42
1	B	200/349 (57%)	177 (88%)	23 (12%)	7	27
All	All	459/698 (66%)	413 (90%)	46 (10%)	9	34

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	ARG
1	A	122	ASN
1	A	179	LEU
1	A	181	SER
1	A	214	LEU
1	A	231	LYS
1	A	245	GLU
1	A	247	LYS
1	A	257	LEU
1	A	275	HIS
1	A	276	LYS
1	A	284	LEU
1	A	292	LEU
1	A	308	GLN
1	A	356	TYR
1	A	358	LYS
1	A	380	SER
1	A	395	THR
1	A	396	GLU
1	A	400	ILE
1	A	406	LEU
1	A	419	LEU
1	A	453	ARG
1	B	93	LEU
1	B	96	LYS
1	B	112	HIS
1	B	116	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	124	LEU
1	B	230	GLU
1	B	296	THR
1	B	311	ASN
1	B	315	LEU
1	B	324	GLN
1	B	326	LEU
1	B	327	MET
1	B	328	LEU
1	B	331	PHE
1	B	334	SER
1	B	335	SER
1	B	342	ASP
1	B	353	ASP
1	B	376	LEU
1	B	387	GLN
1	B	406	LEU
1	B	453	ARG
1	B	457	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	122	ASN
1	A	442	GLN
1	B	288	ASN
1	B	308	GLN
1	B	311	ASN
1	B	345	ASN
1	B	346	HIS
1	B	387	GLN
1	B	410	ASN
1	B	417	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ADP	A	1458	-	22,29,29	1.08	1 (4%)	27,45,45	1.96	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1458	-	-	0/12/32/32	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1458	ADP	C5-C4	3.28	1.47	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1458	ADP	N3-C2-N1	-6.45	123.95	128.89
2	A	1458	ADP	C4-C5-N7	-4.49	105.35	109.48
2	A	1458	ADP	C2'-C1'-N9	-3.02	109.67	114.29

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	1458	ADP	O3B-PB-O2B	2.15	115.56	107.38
2	A	1458	ADP	O4'-C1'-N9	2.26	112.84	108.10
2	A	1458	ADP	C4'-O4'-C1'	2.65	112.63	109.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/390 (81%)	-0.28	9 (2%) 56 32	33, 46, 69, 76	0
1	B	274/390 (70%)	-0.10	8 (2%) 55 31	36, 54, 72, 75	0
All	All	590/780 (75%)	-0.20	17 (2%) 55 31	33, 52, 70, 76	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	121	SER	6.6
1	B	122	ASN	3.7
1	A	118	GLY	3.2
1	B	357	GLU	3.0
1	A	109	ASP	2.9
1	B	390	PHE	2.7
1	B	114	SER	2.6
1	A	432	SER	2.6
1	A	355	SER	2.6
1	A	87	THR	2.5
1	A	112	HIS	2.3
1	B	201	PRO	2.2
1	A	107	ARG	2.2
1	B	294	GLU	2.1
1	B	359	TYR	2.1
1	A	130	PRO	2.1
1	B	245	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ADP	A	1458	27/27	0.90	0.22	0.53	59,60,66,67	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.