



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:53 AM GMT

PDB ID : 3CKD
Title : Crystal structure of the C-terminal domain of the Shigella type III effector IpaH
Authors : Lam, R.; Singer, A.U.; Cuff, M.E.; Skarina, T.; Kagan, O.; DiLeo, R.; Edwards, A.M.; Joachimiak, A.; Savchenko, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2008-03-14
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	<div> <div>7%</div> <div>55%</div> <div>23%</div> <div>6%</div> <div>16%</div> </div>
1	B	312	<div> <div>9%</div> <div>61%</div> <div>12%</div> <div>5%</div> <div>22%</div> </div>
1	C	312	<div> <div>%</div> <div>61%</div> <div>22%</div> <div>• •</div> <div>13%</div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Invasion plasmid antigen, secreted by the Mxi-Spa secretion machinery.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	Se	0	0	0
			2114	1319	373	416	1	5			
1	B	244	Total	C	N	O	Se		0	0	0
			1970	1233	343	389	5				
1	C	272	Total	C	N	O	S	Se	0	0	0
			2184	1359	385	434	1	5			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	GLY	-	EXPRESSION TAG	UNP Q8VSA1
B	264	GLY	-	EXPRESSION TAG	UNP Q8VSA1
C	264	GLY	-	EXPRESSION TAG	UNP Q8VSA1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			1	1		
5	B	4	Total	O	0	0
			4	4		
5	C	19	Total	O	0	0
			19	19		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Invasion plasmid antigen, secreted by the Mxi-Spa secretion machinery



- Molecule 1: Invasion plasmid antigen, secreted by the Mxi-Spa secretion machinery



- Molecule 1: Invasion plasmid antigen, secreted by the Mxi-Spa secretion machinery



ASN	HIS	ILE	ALA
VAL	LYS	GLU	MSE
PHE	TYR	G457	V458
		S459	G460
		V461	T462
		A463	N464
		E477	
		F481	
		F485	
		G489	
		K496	
		W503	
		E507	
		Q520	
		K527	
		G533	D534
		A537	
		E540	
		T549	
		Y554	R555
		Q556	L557
		T558	
		V561	L562
		A563	L564
		R565	LEU
		SER	GLU
		ASN	GLY
		SER	
GLY	GLY	PRO	GLN
ILE	PHE	PHE	SER
MSE	GLY	ASN	SER
ALA	ALA	PRO	GLU
			W283
		V289	T290
		A291	W292
		N296	K297
		S302	
		W305	H306
		E309	H310
		E311	E312
		T316	F317
		S318	
		S325	D326
		T327	V328
		S329	ALA
		ARG	ASN
		T333	S334
		G335	
		Q339	
		Q437	A441
		L444	Q445
		L446	S447
		T448	A451
		E452	
Q355	V360	A361	A362
T365	E366	S367	C368
E369	D370	R371	
L374	T375	W376	
L379	R380	T382	
E390			
L401	R405		
F408	R409		
I412	A417	R418	D419
K420	V421	H425	F426
V427	D428	E429	I430
E431	V432		
Q437	A441	L444	Q445
L446	S447	T448	A451
E452			

4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	128.93Å 128.93Å 282.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 2.65 49.19 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.21-2.65) 98.9 (49.19-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.49 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.284 0.261 , 0.311	Depositor DCC
R_{free} test set	1743 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	58.5	Xtriage
Anisotropy	0.005	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 34706 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6327	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/2145	0.72	1/2889 (0.0%)
1	B	0.71	0/1996	0.75	3/2688 (0.1%)
1	C	1.04	7/2217 (0.3%)	0.97	8/2990 (0.3%)
All	All	0.83	7/6358 (0.1%)	0.82	12/8567 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	481	PHE	CE1-CZ	5.96	1.48	1.37
1	C	507	GLU	CG-CD	5.52	1.60	1.51
1	C	485	PHE	CE1-CZ	5.36	1.47	1.37
1	C	390	GLU	CB-CG	-5.33	1.42	1.52
1	C	540	GLU	CG-CD	5.30	1.59	1.51

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	409	ARG	NE-CZ-NH2	-8.23	116.19	120.30
1	C	565	ARG	NE-CZ-NH2	-7.70	116.45	120.30
1	C	409	ARG	NE-CZ-NH1	7.14	123.87	120.30
1	C	418	ARG	NE-CZ-NH1	6.39	123.49	120.30
1	C	374	LEU	CA-CB-CG	6.03	129.17	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	368	CYS	Peptide
1	C	533	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2114	0	2017	64	0
1	B	1970	0	1885	26	0
1	C	2184	0	2081	41	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	C	18	0	24	1	0
4	C	7	0	10	1	0
5	A	1	0	0	0	0
5	B	4	0	0	0	0
5	C	19	0	0	0	0
All	All	6327	0	6017	130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:555:ARG:CG	1:A:555:ARG:HH11	1.56	1.16
1:A:555:ARG:HH11	1:A:555:ARG:HG3	1.08	1.11
1:A:303:GLN:HE21	1:A:304:ILE:HG13	1.19	1.06
1:A:555:ARG:NH1	1:A:555:ARG:HG3	1.71	0.97
1:A:468:THR:HG22	1:A:472:MSE:HE2	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/312 (81%)	219 (86%)	31 (12%)	4 (2%)	12	26
1	B	232/312 (74%)	207 (89%)	20 (9%)	5 (2%)	8	19
1	C	266/312 (85%)	247 (93%)	16 (6%)	3 (1%)	17	38
All	All	752/936 (80%)	673 (90%)	67 (9%)	12 (2%)	12	26

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	HIS
1	B	394	ASP
1	A	370	ASP
1	B	314	ALA
1	B	564	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/252 (86%)	183 (84%)	35 (16%)	3	6
1	B	204/252 (81%)	177 (87%)	27 (13%)	5	10
1	C	228/252 (90%)	201 (88%)	27 (12%)	6	13
All	All	650/756 (86%)	561 (86%)	89 (14%)	4	9

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	318	SER
1	B	420	LYS
1	C	428	ASP
1	B	324	LEU
1	B	380	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	544	GLN
1	B	296	ASN
1	C	283	HIS
1	A	516	ASN
1	B	520	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	601	-	4,4,4	0.35	0	6,6,6	0.23	0
2	SO4	C	600	-	4,4,4	0.24	0	6,6,6	0.71	0
3	GOL	C	602	-	5,5,5	0.37	0	5,5,5	0.52	0
3	GOL	C	603	-	5,5,5	0.40	0	5,5,5	0.48	0
3	GOL	C	604	-	5,5,5	0.27	0	5,5,5	0.78	0
4	PEG	C	605	-	6,6,6	0.64	0	5,5,5	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	B	601	-	-	0/0/0/0	0/0/0/0
2	SO4	C	600	-	-	0/0/0/0	0/0/0/0
3	GOL	C	602	-	-	0/4/4/4	0/0/0/0
3	GOL	C	603	-	-	0/4/4/4	0/0/0/0
3	GOL	C	604	-	-	0/4/4/4	0/0/0/0
4	PEG	C	605	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	603	GOL	1	0
4	C	605	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/312 (82%)	0.78	23 (8%) 12 9	63, 81, 105, 113	22 (8%)
1	B	239/312 (76%)	0.73	27 (11%) 7 5	46, 62, 81, 89	9 (3%)
1	C	267/312 (85%)	0.20	3 (1%) 82 82	24, 36, 67, 79	9 (3%)
All	All	763/936 (81%)	0.56	53 (6%) 20 17	24, 63, 93, 113	40 (5%)

The worst 5 of 53 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	395	ASN	4.9
1	A	305	TRP	4.7
1	A	293	PHE	4.6
1	B	350	SER	4.2
1	B	285	LEU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	C	603	6/6	0.89	0.22	1.54	49,53,54,55	6
2	SO4	C	600	5/5	0.98	0.11	-	48,48,51,51	5
3	GOL	C	602	6/6	0.91	0.26	-	44,46,46,47	6
3	GOL	C	604	6/6	0.90	0.18	-	63,65,67,72	0
2	SO4	B	601	5/5	0.96	0.16	-	55,56,58,59	5
4	PEG	C	605	7/7	0.87	0.21	-	43,52,61,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.