Full wwPDB X-ray Structure Validation Report (i)

Feb 25, 2016 – 11:15 AM GMT

PDB ID	:	5CKK
Title	:	Crystal structure of 9DB1* deoxyribozyme
Authors	:	Ponce-Salvatierra, A.; Hoebartner, C.; Pena, V.
Deposited on	:	2015-07-15
Resolution	:	2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	FAILED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2827 (2.80-2.80)
RNA backbone	2183	1091 (3.20-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	Y	44	50% 50%
2	D	15	67% 27% 7%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (44-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	Y	44	909	431	166	268	44	0	0	0

- Molecule 2 is a RNA chain called RNA (5'-R(P*GP*CP*AP*CP*UP*AP*GP*AP*UP*CP *GP*GP*AP*UP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	D	15	323	144	60	104	15	0	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	Y	2	2	2	0	0
3	D	1	1	1	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

• Molecule 1: DNA (44-MER)

Chain Y:

• Molecule 2: RNA (5'-R(P*GP*CP*AP*CP*UP*AP*GP*AP*UP*CP*GP*GP*AP*UP*G)-3')

Chain D:



4 Data and refinement statistics (i)

EDS failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α, β, γ	83.16Å 83.16Å 55.99Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.45 – 2.80	Depositor
% Data completeness (in resolution range)	96.6 (46.45-2.80)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	0.92 (at 2.81Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.275 , 0.287	Depositor
Wilson B-factor (Å^2)	113.6	Xtriage
Anisotropy	0.639	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 5074 reflections	Xtriage
Total number of atoms	1235	wwPDB-VP
Average B, all atoms (Å^2)	148.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	Y	0.53	0/1019	0.92	0/1573
2	D	0.19	0/361	0.74	0/561
All	All	0.47	0/1380	0.88	0/2134

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Y	909	0	497	15	0
2	D	323	0	163	3	0
3	D	1	0	0	0	0
3	Y	2	0	0	0	0
All	All	1235	0	660	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Y:16:DC:H2'	1:Y:17:DG:C8	2.24	0.72
1:Y:25:DG:H2"	1:Y:26:DG:H5"	1.78	0.66
1:Y:32:DC:H2'	1:Y:33:DC:C6	2.35	0.61
1:Y:1:DA:H3'	1:Y:2:DT:H71	1.86	0.57
1:Y:10:DA:H2'	1:Y:10:DA:N3	2.24	0.52
1:Y:33:DC:H2'	1:Y:34:DG:C8	2.46	0.50
1:Y:8:DG:H2"	1:Y:24:DG:N7	2.27	0.49
1:Y:10:DA:H2"	1:Y:11:DT:C6	2.49	0.47
1:Y:7:DT:H2"	1:Y:8:DG:C8	2.50	0.47
1:Y:21:DG:H1'	1:Y:23:DA:N6	2.29	0.47
1:Y:38:DA:H4'	1:Y:39:DA:OP1	2.15	0.47
1:Y:10:DA:H4'	1:Y:11:DT:OP1	2.16	0.45
1:Y:27:DG:H3'	1:Y:28:DT:H5'	1.98	0.44
2:D:-5:C:H2'	2:D:-4:A:C8	2.53	0.43
2:D:8:U:H2'	2:D:9:G:H8	1.84	0.42
1:Y:31:DG:H2'	1:Y:32:DC:C6	2.55	0.42
2:D:8:U:H2'	2:D:9:G:C8	2.56	0.41
1:Y:11:DT:H1'	1:Y:12:DC:H5	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

There are no protein molecules in this entry.

5.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	14/15 (93%)	2 (14%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	-5	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	1	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS failed to run properly - this section will therefore be empty.

6.4 Ligands (i)

EDS failed to run properly - this section will therefore be empty.

6.5 Other polymers (i)

EDS failed to run properly - this section will therefore be empty.