



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:12 PM GMT

PDB ID : 5CKL  
Title : Fic protein from *Neisseria meningitidis* (NmFic) mutant E156R in dimeric form  
Authors : Stanger, F.V.; Schirmer, T.  
Deposited on : 2015-07-15  
Resolution : 0.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 0.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1235 (1.10-0.90)
Clashscore	102246	1333 (1.10-0.90)
Ramachandran outliers	100387	1247 (1.10-0.90)
Sidechain outliers	100360	1246 (1.10-0.90)
RSRZ outliers	91569	1239 (1.10-0.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	188	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	201	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Adenosine monophosphate-protein transferase NmFic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	181	1670	1053	301	310	6	6	19	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	MET	-	initiating methionine	UNP Q7DDR9
A	5	HIS	-	expression tag	UNP Q7DDR9
A	6	HIS	-	expression tag	UNP Q7DDR9
A	7	HIS	-	expression tag	UNP Q7DDR9
A	8	HIS	-	expression tag	UNP Q7DDR9
A	9	HIS	-	expression tag	UNP Q7DDR9
A	10	HIS	-	expression tag	UNP Q7DDR9
A	156	ARG	GLU	engineered mutation	UNP Q7DDR9

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Cl	0	0
			3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	355	Total	O	0	7
			362	362		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosine monophosphate-protein transferase NmFic



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.33Å 50.54Å 130.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.07 – 0.99 47.11 – 0.99	Depositor EDS
% Data completeness (in resolution range)	99.5 (65.07-0.99) 99.5 (47.11-0.99)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 0.99Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.112 , 0.125 0.112 , 0.125	Depositor DCC
$R_{free}$ test set	6534 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	8.4	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 129865 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	2041	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.36	22/1700 (1.3%)	1.61	36/2280 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	180	GLU	CG-CD	12.46	1.70	1.51
1	A	95	GLU	CD-OE1	-9.61	1.15	1.25
1	A	167	ASP	CB-CG	-8.81	1.33	1.51
1	A	173	GLU	CD-OE2	-8.52	1.16	1.25
1	A	48[A]	TYR	CB-CG	-8.29	1.39	1.51
1	A	48[B]	TYR	CB-CG	-8.29	1.39	1.51
1	A	71	ARG	CZ-NH1	-7.54	1.23	1.33
1	A	24[A]	ARG	CZ-NH1	7.51	1.42	1.33
1	A	24[B]	ARG	CZ-NH1	7.51	1.42	1.33
1	A	189	GLU	CG-CD	7.09	1.62	1.51
1	A	139	SER	CB-OG	-7.04	1.33	1.42
1	A	189	GLU	CD-OE1	6.75	1.33	1.25
1	A	91[A]	ARG	CZ-NH2	6.67	1.41	1.33
1	A	91[B]	ARG	CZ-NH2	6.67	1.41	1.33
1	A	173	GLU	CG-CD	6.62	1.61	1.51
1	A	180	GLU	CB-CG	-6.08	1.40	1.52
1	A	156	ARG	CD-NE	-6.06	1.36	1.46
1	A	184	TYR	CD1-CE1	5.59	1.47	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131	LYS	CE-NZ	-5.32	1.35	1.49
1	A	80[A]	GLU	CD-OE2	5.24	1.31	1.25
1	A	80[B]	GLU	CD-OE2	5.24	1.31	1.25
1	A	180	GLU	CD-OE2	5.02	1.31	1.25

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47[A]	ARG	NE-CZ-NH2	-20.72	109.94	120.30
1	A	47[B]	ARG	NE-CZ-NH2	-20.72	109.94	120.30
1	A	24[A]	ARG	NE-CZ-NH1	18.41	129.50	120.30
1	A	24[B]	ARG	NE-CZ-NH1	18.41	129.50	120.30
1	A	24[A]	ARG	NE-CZ-NH2	-14.75	112.93	120.30
1	A	24[B]	ARG	NE-CZ-NH2	-14.75	112.93	120.30
1	A	47[A]	ARG	CD-NE-CZ	11.75	140.05	123.60
1	A	47[B]	ARG	CD-NE-CZ	11.75	140.05	123.60
1	A	189	GLU	OE1-CD-OE2	11.44	137.03	123.30
1	A	184	TYR	CB-CG-CD2	-10.95	114.43	121.00
1	A	47[A]	ARG	NE-CZ-NH1	10.62	125.61	120.30
1	A	47[B]	ARG	NE-CZ-NH1	10.62	125.61	120.30
1	A	49	LEU	CB-CG-CD2	9.07	126.42	111.00
1	A	184	TYR	CB-CG-CD1	8.69	126.22	121.00
1	A	91[A]	ARG	NE-CZ-NH2	8.37	124.48	120.30
1	A	91[B]	ARG	NE-CZ-NH2	8.37	124.48	120.30
1	A	71	ARG	NE-CZ-NH2	-7.76	116.42	120.30
1	A	155	LEU	CB-CG-CD1	-6.95	99.18	111.00
1	A	156	ARG	NE-CZ-NH2	6.69	123.65	120.30
1	A	158[A]	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	A	158[B]	ARG	NE-CZ-NH1	6.66	123.63	120.30
1	A	32	ASP	CB-CG-OD2	6.64	124.28	118.30
1	A	70	PHE	CB-CG-CD2	6.62	125.44	120.80
1	A	55[A]	ASP	CB-CG-OD2	-6.37	112.56	118.30
1	A	55[B]	ASP	CB-CG-OD2	-6.37	112.56	118.30
1	A	149	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	68	GLY	CA-C-O	6.04	131.47	120.60
1	A	68	GLY	O-C-N	-5.97	113.06	123.20
1	A	170	ASP	CB-CG-OD2	-5.90	112.99	118.30
1	A	189	GLU	CG-CD-OE2	-5.84	106.62	118.30
1	A	159	PHE	CB-CG-CD1	5.70	124.79	120.80
1	A	167	ASP	CB-CG-OD2	-5.67	113.20	118.30
1	A	69	GLY	O-C-N	5.47	131.46	122.70
1	A	159	PHE	CB-CG-CD2	-5.36	117.05	120.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	180	GLU	OE1-CD-OE2	5.26	129.61	123.30
1	A	185	TYR	CB-CG-CD1	5.09	124.05	121.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	33	ARG	Mainchain
1	A	69	GLY	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1670	0	1634	16	0
2	A	6	0	7	0	0
3	A	3	0	0	0	0
4	A	362	0	0	9	1
All	All	2041	0	1641	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:145[B]:GLN:NE2	1:A:145[B]:GLN:CD	1.79	1.36
1:A:158[A]:ARG:HD3	4:A:337[A]:HOH:O	1.71	0.91
1:A:84[A]:LYS:NZ	4:A:301:HOH:O	2.06	0.88
1:A:140:LYS:NZ	4:A:303[A]:HOH:O	2.17	0.78
1:A:47[B]:ARG:NH2	4:A:304:HOH:O	2.17	0.77
1:A:44[B]:GLN:HG3	4:A:440:HOH:O	1.89	0.71
1:A:20[B]:HIS:HD2	4:A:532:HOH:O	1.73	0.70
1:A:34[B]:ILE:HG22	1:A:44[B]:GLN:CD	2.14	0.68
1:A:91[A]:ARG:O	1:A:91[A]:ARG:HG2	1.99	0.61

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91[B]:ARG:NH2	4:A:307:HOH:O	2.35	0.58
1:A:34[B]:ILE:HG22	1:A:44[B]:GLN:OE1	2.04	0.58
1:A:93:PHE:HE2	1:A:158[B]:ARG:HG2	1.71	0.56
1:A:20[B]:HIS:HE1	4:A:410:HOH:O	1.88	0.55
1:A:20[B]:HIS:CE1	4:A:410:HOH:O	2.65	0.49
1:A:47[B]:ARG:HG3	1:A:58:GLY:HA3	1.99	0.43
1:A:94[B]:GLU:HG3	1:A:158[B]:ARG:HD3	2.01	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:570:HOH:O	4:A:570:HOH:O[2_575]	1.35	0.85

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/188 (105%)	196 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/166 (108%)	178 (99%)	1 (1%)	90	65

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	107	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	201	-	5,5,5	2.38	3 (60%)	5,5,5	1.24	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	201	-	-	0/4/4/4	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	GOL	C3-C2	-2.95	1.40	1.52
2	A	201	GOL	O1-C1	-2.09	1.33	1.42
2	A	201	GOL	O3-C3	3.80	1.58	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	GOL	O2-C2-C1	2.16	118.84	108.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	33:ARG	C	34[B]:ILE	N	1.20

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/188 (96%)	-0.08	1 (0%) 90 82	5, 9, 22, 47	3 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	HIS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	201	6/6	0.94	0.10	6.10	11,14,21,22	0
3	CL	A	203	1/1	0.98	0.14	-	43,43,43,43	1
3	CL	A	202	1/1	0.86	0.17	-	63,63,63,63	0
3	CL	A	204	1/1	1.00	0.05	-	8,8,8,8	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.