



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:49 PM GMT

PDB ID : 1CMY
Title : THE MUTATION BETA99 ASP-TYR STABILIZES Y-A NEW, COMPOSITE QUATERNARY STATE OF HUMAN HEMOGLOBIN
Authors : Smith, F.R.; Lattman, E.E.; Carter Junior, C.W.
Deposited on : 1992-09-18
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	141	 45% 46% 7%
1	C	141	 36% 55% 9%
2	B	146	 38% 51% 10%
2	D	146	 39% 51% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4564 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN YPSILANTI (CARBONMONOXY) (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			
1	C	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			

- Molecule 2 is a protein called HEMOGLOBIN YPSILANTI (CARBONMONOXY) (BETA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1127	729	195	200	3			
2	D	146	Total	C	N	O	S	0	0	0
			1127	729	195	200	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	99	TYR	ASP	CONFLICT	UNP P68871
D	99	TYR	ASP	CONFLICT	UNP P68871

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: HEMOGLOBIN YPSILANTI (CARBONMONOXY) (ALPHA CHAIN)



• Molecule 1: HEMOGLOBIN YPSILANTI (CARBONMONOXY) (ALPHA CHAIN)



• Molecule 2: HEMOGLOBIN YPSILANTI (CARBONMONOXY) (BETA CHAIN)



• Molecule 2: HEMOGLOBIN YPSILANTI (CARBONMONOXY) (BETA CHAIN)





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.10Å 93.10Å 144.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.269 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4564	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/1097	1.23	4/1491 (0.3%)
1	C	0.70	0/1097	1.22	4/1491 (0.3%)
2	B	0.71	0/1158	1.20	2/1573 (0.1%)
2	D	0.71	0/1158	1.25	4/1573 (0.3%)
All	All	0.70	0/4510	1.22	14/6128 (0.2%)

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	77	HIS	CA-CB-CG	8.15	127.46	113.60
2	D	79	ASP	CB-CG-OD1	-8.07	111.04	118.30
1	C	141	ARG	NE-CZ-NH1	7.09	123.84	120.30
2	D	50	THR	N-CA-CB	6.76	123.15	110.30
1	A	50	HIS	CA-CB-CG	-5.55	104.16	113.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	137	11
1	C	1069	0	1073	138	11

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1127	0	1123	126	3
2	D	1127	0	1123	121	21
3	A	43	0	30	13	1
3	B	43	0	30	6	0
3	C	43	0	30	10	0
3	D	43	0	30	5	0
All	All	4564	0	4512	477	24

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 53.

The worst 5 of 477 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:145:TYR:CE1	2:D:146:HIS:CE1	1.87	1.60
2:D:145:TYR:CD1	2:D:146:HIS:CE1	1.93	1.55
2:D:145:TYR:CD1	2:D:146:HIS:ND1	2.01	1.28
2:B:145:TYR:CE1	2:B:146:HIS:CE1	2.33	1.16
1:C:1:VAL:CG1	1:C:2:LEU:H	1.57	1.14

The worst 5 of 24 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:LYS:CA	2:D:90:GLU:OE2[4_567]	0.76	1.44
1:C:90:LYS:CE	2:D:44:SER:CA[2_664]	1.03	1.17
1:C:90:LYS:CE	2:D:44:SER:C[2_664]	1.19	1.01
1:C:90:LYS:NZ	2:D:44:SER:CB[2_664]	1.25	0.95
1:A:15:GLY:O	2:D:90:GLU:OE1[4_567]	1.29	0.91

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	131 (94%)	5 (4%)	3 (2%)	8	38
1	C	139/141 (99%)	131 (94%)	5 (4%)	3 (2%)	8	38
2	B	144/146 (99%)	133 (92%)	8 (6%)	3 (2%)	9	40
2	D	144/146 (99%)	133 (92%)	9 (6%)	2 (1%)	14	51
All	All	566/574 (99%)	528 (93%)	27 (5%)	11 (2%)	10	43

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
2	B	3	LEU
2	B	77	HIS
1	C	72	HIS
2	D	3	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	102 (90%)	11 (10%)	10	37
1	C	113/113 (100%)	100 (88%)	13 (12%)	7	28
2	B	118/118 (100%)	98 (83%)	20 (17%)	2	13
2	D	118/118 (100%)	96 (81%)	22 (19%)	2	10
All	All	462/462 (100%)	396 (86%)	66 (14%)	4	19

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	145	TYR
1	C	109	LEU
2	D	109	VAL
2	B	146	HIS
1	C	85	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	117	HIS
2	B	146	HIS
2	D	77	HIS
2	B	108	ASN
2	D	102	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HEM	A	142	1	30,50,50	2.48	6 (20%)	24,82,82	2.70	12 (50%)
3	HEM	B	147	2	30,50,50	2.42	7 (23%)	24,82,82	3.01	11 (45%)
3	HEM	C	142	1	30,50,50	2.45	7 (23%)	24,82,82	2.74	12 (50%)
3	HEM	D	147	2	30,50,50	2.47	7 (23%)	24,82,82	2.68	13 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	142	1	-	0/10/54/54	0/0/8/8
3	HEM	B	147	2	-	0/10/54/54	0/0/8/8
3	HEM	C	142	1	-	0/10/54/54	0/0/8/8
3	HEM	D	147	2	-	0/10/54/54	0/0/8/8

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	142	HEM	C3B-C4B	-7.60	1.45	1.51
3	A	142	HEM	C3B-C4B	-7.57	1.45	1.51
3	D	147	HEM	C3B-C4B	-7.51	1.45	1.51
3	B	147	HEM	C3B-C4B	-7.46	1.45	1.51
3	A	142	HEM	C2D-C3D	-6.95	1.33	1.54

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	147	HEM	CAA-C2A-C1A	-4.49	122.13	127.01
3	A	142	HEM	C3C-CAC-CBC	-4.04	118.26	124.46
3	C	142	HEM	CAA-C2A-C1A	-3.53	123.18	127.01
3	C	142	HEM	CMA-C3A-C4A	-3.46	122.63	128.36
3	D	147	HEM	CMA-C3A-C4A	-3.34	122.84	128.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 35 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	142	HEM	13	1
3	B	147	HEM	6	0
3	C	142	HEM	10	0
3	D	147	HEM	5	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.