



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:52 PM GMT

PDB ID : 1COW
Title : BOVINE MITOCHONDRIAL F1-ATPASE COMPLEXED WITH AU-ROVERTIN B
Authors : Van Raaij, M.; Abrahams, J.P.; Leslie, A.G.W.; Walker, J.E.
Deposited on : 1996-05-08
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	510	<div> <div></div> <div> <div></div> <div>49%</div> <div>35%</div> <div>10%</div> <div>5%</div> </div> </div>
1	B	510	<div> <div>2%</div> <div> <div></div> <div>47%</div> <div>38%</div> <div>10%</div> <div>5%</div> </div> </div>
1	C	510	<div> <div></div> <div> <div></div> <div>53%</div> <div>37%</div> <div>5%</div> <div></div> </div> </div>
2	D	482	<div> <div>2%</div> <div> <div></div> <div>52%</div> <div>38%</div> <div>6%</div> <div></div> </div> </div>
2	E	482	<div> <div>4%</div> <div> <div></div> <div>43%</div> <div>40%</div> <div>11%</div> <div></div> </div> </div>
2	F	482	<div> <div></div> <div> <div></div> <div>55%</div> <div>34%</div> <div>7%</div> <div></div> </div> </div>
3	G	272	<div> <div>7%</div> <div> <div></div> <div>25%</div> <div>15%</div> <div>5%</div> <div></div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	D	601	-	-	-	X
4	MG	F	601	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 23462 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	0	0	0
			3715	2341	656	706	12			
1	B	487	Total	C	N	O	S	0	0	0
			3715	2341	656	706	12			
1	C	492	Total	C	N	O	S	0	0	0
			3748	2360	661	715	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	SER	CONFLICT	UNP P19483
B	481	GLY	SER	CONFLICT	UNP P19483
C	481	GLY	SER	CONFLICT	UNP P19483

- Molecule 2 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	467	Total	C	N	O	S	0	0	0
			3539	2243	601	684	11			
2	E	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			
2	F	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			


- Molecule 3 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	122	Total	C	N	O	S	0	0	0
			945	591	171	176	7			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	F	1	Total	Mg	0	0
			1	1		

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
5	B	1	Total 31	C 10	N 6	O 12	P 3	0	0
5	C	1	Total 31	C 10	N 6	O 12	P 3	0	0
5	F	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 7 is AUROVERTIN B (three-letter code: AUR) (formula: C₂₅H₃₂O₈).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			33	25	8		
7	E	1	Total	C	O	0	0
			33	25	8		


- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	80	Total 80	O 80	0	0
8	B	84	Total 84	O 84	0	0
8	C	98	Total 98	O 98	0	0
8	D	94	Total 94	O 94	0	0
8	E	47	Total 47	O 47	0	0
8	F	92	Total 92	O 92	0	0
8	G	23	Total 23	O 23	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BOVINE MITOCHONDRIAL F1-ATPASE








• Molecule 1: BOVINE MITOCHONDRIAL F1-ATPASE







• Molecule 2: BOVINE MITOCHONDRIAL F1-ATPASE







● Molecule 3: BOVINE MITOCHONDRIAL F1-ATPASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	283.40 Å 107.60 Å 140.20 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10 20.00 – 3.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-3.10) 99.4 (20.00-3.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.44 (at 3.09 Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	(Not available) , 0.280 0.216 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	53.8	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 91.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 77818 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	23462	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AUR, ANP, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	17/3766 (0.5%)	1.38	45/5080 (0.9%)
1	B	1.00	16/3766 (0.4%)	1.40	41/5080 (0.8%)
1	C	1.00	19/3799 (0.5%)	1.38	39/5126 (0.8%)
2	D	1.01	24/3596 (0.7%)	1.40	41/4879 (0.8%)
2	E	1.02	25/3587 (0.7%)	1.36	47/4867 (1.0%)
2	F	1.07	26/3587 (0.7%)	1.39	55/4867 (1.1%)
3	G	0.99	7/949 (0.7%)	1.42	18/1266 (1.4%)
All	All	1.01	134/23050 (0.6%)	1.39	286/31165 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	2
2	D	0	1
2	E	0	2
3	G	1	3
All	All	1	11

The worst 5 of 134 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	26	GLU	CD-OE1	8.28	1.34	1.25
2	F	42	GLU	CD-OE2	8.05	1.34	1.25
2	F	224	GLU	CD-OE2	7.94	1.34	1.25
2	D	436	GLU	CD-OE2	7.43	1.33	1.25
1	A	491	GLU	CD-OE1	7.19	1.33	1.25

The worst 5 of 286 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	408	ARG	NE-CZ-NH2	-16.74	111.93	120.30
2	D	408	ARG	NE-CZ-NH1	15.40	128.00	120.30
1	A	143	ARG	NE-CZ-NH2	-13.38	113.61	120.30
2	D	408	ARG	CD-NE-CZ	11.78	140.09	123.60
3	G	9	ARG	NE-CZ-NH1	11.37	125.99	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	G	209	LEU	CA

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	PHE	Mainchain
1	B	369	LEU	Mainchain
1	B	432	GLN	Mainchain
1	C	363	PRO	Mainchain
1	C	405	GLN	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3815	162	0
1	B	3715	0	3814	168	0
1	C	3748	0	3845	137	0
2	D	3539	0	3593	140	0
2	E	3530	0	3587	184	0
2	F	3530	0	3587	137	0
3	G	945	0	1019	39	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	31	0	13	1	0
5	B	31	0	13	8	0
5	C	31	0	13	4	0
5	F	31	0	12	2	0
6	D	27	0	12	3	0
7	E	33	0	32	14	0
7	F	33	0	32	6	0
8	A	80	0	0	6	0
8	B	84	0	0	12	0
8	C	98	0	0	11	0
8	D	94	0	0	11	0
8	E	47	0	0	9	0
8	F	92	0	0	11	0
8	G	23	0	0	2	0
All	All	23462	0	23387	914	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 914 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:215:GLN:HG3	2:F:356:ARG:HH22	1.07	1.15
1:C:127:ARG:HH12	1:C:255:GLU:HB2	0.97	1.07
1:C:30:ARG:HE	1:C:87:ILE:HD11	1.24	1.01
2:F:223:ASN:HD22	2:F:223:ASN:H	1.06	1.00
1:A:215:GLN:HG3	2:D:356:ARG:NH1	1.80	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/510 (95%)	430 (89%)	44 (9%)	11 (2%)	8	35
1	B	485/510 (95%)	430 (89%)	45 (9%)	10 (2%)	9	37
1	C	490/510 (96%)	438 (89%)	46 (9%)	6 (1%)	16	52
2	D	465/482 (96%)	415 (89%)	45 (10%)	5 (1%)	17	55
2	E	464/482 (96%)	401 (86%)	46 (10%)	17 (4%)	4	23
2	F	464/482 (96%)	422 (91%)	40 (9%)	2 (0%)	39	75
3	G	116/272 (43%)	98 (84%)	16 (14%)	2 (2%)	11	43
All	All	2969/3248 (91%)	2634 (89%)	282 (10%)	53 (2%)	11	42

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	25	LEU
1	A	57	SER
1	A	405	GLN
1	B	25	LEU
1	B	364	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/412 (95%)	341 (87%)	52 (13%)	5	20
1	B	393/412 (95%)	351 (89%)	42 (11%)	8	31
1	C	397/412 (96%)	365 (92%)	32 (8%)	15	47
2	D	377/386 (98%)	348 (92%)	29 (8%)	16	50
2	E	376/386 (97%)	339 (90%)	37 (10%)	10	36
2	F	376/386 (97%)	349 (93%)	27 (7%)	18	53
3	G	102/230 (44%)	94 (92%)	8 (8%)	16	49
All	All	2414/2624 (92%)	2187 (91%)	227 (9%)	11	39

5 of 227 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	101	GLU
2	D	56	SER
2	F	381	TYR
1	C	164	ARG
1	C	334	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	223	ASN
2	E	130	GLN
2	F	282	GLN
2	E	39	GLN
2	E	194	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 5 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ANP	A	600	4	27,33,33	1.53	5 (18%)	30,52,52	2.42	8 (26%)
5	ANP	B	600	4	27,33,33	1.42	7 (25%)	30,52,52	2.14	6 (20%)
5	ANP	C	600	4	27,33,33	1.74	6 (22%)	30,52,52	2.16	7 (23%)
6	ADP	D	600	4	22,29,29	1.09	1 (4%)	27,45,45	1.63	6 (22%)
7	AUR	E	479	-	31,35,35	1.90	6 (19%)	31,52,52	1.88	8 (25%)
5	ANP	F	600	4	27,33,33	1.55	8 (29%)	30,52,52	2.72	10 (33%)
7	AUR	F	602	-	31,35,35	1.54	5 (16%)	31,52,52	2.10	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ANP	A	600	4	-	1/12/38/38	0/3/3/3
5	ANP	B	600	4	-	1/12/38/38	0/3/3/3
5	ANP	C	600	4	-	1/12/38/38	0/3/3/3
6	ADP	D	600	4	-	0/12/32/32	0/3/3/3
7	AUR	E	479	-	-	0/16/55/55	0/1/3/3
5	ANP	F	600	4	-	0/12/38/38	0/3/3/3
7	AUR	F	602	-	-	0/16/55/55	0/1/3/3

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	E	479	AUR	C7-C8	-5.58	1.47	1.53
7	F	602	AUR	C7-C8	-4.30	1.48	1.53
7	E	479	AUR	O5-C5	-3.92	1.36	1.44
7	F	602	AUR	O5-C5	-3.79	1.36	1.44
5	F	600	ANP	O3'-C3'	-3.24	1.35	1.43

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	600	ANP	O1G-PG-N3B	-9.27	97.68	111.90
7	F	602	AUR	O17-C17-C16	-6.76	109.39	115.02
5	A	600	ANP	O1G-PG-N3B	-5.64	103.24	111.90
5	C	600	ANP	PA-O3A-PB	-5.01	115.87	132.67
5	B	600	ANP	O2G-PG-O1G	-4.74	100.89	113.49

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	600	ANP	O1G-PG-N3B-PB
5	A	600	ANP	O1G-PG-N3B-PB
5	C	600	ANP	O1G-PG-N3B-PB

There are no ring outliers.

7 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	600	ANP	1	0
5	B	600	ANP	8	0
5	C	600	ANP	4	0
6	D	600	ADP	3	0
7	E	479	AUR	14	0
5	F	600	ANP	2	0
7	F	602	AUR	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	487/510 (95%)	-0.48	4 (0%) 87 75	4, 40, 81, 100	0
1	B	487/510 (95%)	-0.42	10 (2%) 67 44	1, 35, 93, 100	0
1	C	492/510 (96%)	-0.61	4 (0%) 87 75	7, 30, 71, 100	0
2	D	467/482 (96%)	-0.42	9 (1%) 70 48	8, 37, 87, 100	0
2	E	466/482 (96%)	-0.18	17 (3%) 46 23	3, 53, 100, 100	0
2	F	466/482 (96%)	-0.57	3 (0%) 90 80	3, 30, 76, 91	0
3	G	122/272 (44%)	0.61	19 (15%) 3 1	7, 79, 100, 100	0
All	All	2987/3248 (91%)	-0.40	66 (2%) 65 42	1, 38, 92, 100	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	403	PHE	6.7
3	G	86	ALA	6.0
1	B	408	SER	5.1
2	E	389	ALA	4.5
3	G	25	MET	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	MG	D	601	1/1	0.99	0.36	10.37	16,16,16,16	0
4	MG	F	601	1/1	0.92	0.28	8.00	19,19,19,19	0
7	AUR	E	479	33/33	0.80	0.33	0.41	61,64,87,87	33
5	ANP	F	600	31/31	0.96	0.15	0.07	21,25,34,36	0
7	AUR	F	602	33/33	0.95	0.20	0.05	16,22,53,56	33
6	ADP	D	600	27/27	0.96	0.14	-0.30	16,30,34,36	0
5	ANP	A	600	31/31	0.96	0.13	-0.61	17,31,44,45	0
5	ANP	B	600	31/31	0.96	0.14	-0.71	18,37,44,47	0
5	ANP	C	600	31/31	0.97	0.12	-0.73	7,20,25,27	0
4	MG	B	601	1/1	0.88	0.17	-	35,35,35,35	0
4	MG	C	601	1/1	0.95	0.17	-	19,19,19,19	0
4	MG	A	601	1/1	0.97	0.12	-	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.