



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 07:05 PM BST

PDB ID : 2CPY
Title : solution structure of RNA binding domain 3 in RNA binding motif protein 12
Authors : Tsuda, K.; Muto, Y.; Inoue, M.; Kigawa, T.; Terada, T.; Shirouzu, M.;
Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2005-05-19

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	114	<div> <div style="width: 63%; background-color: green;"></div> <div style="width: 7%; background-color: yellow;"></div> <div style="width: 30%; background-color: cyan;"></div> </div> <div>63% 7% 30%</div>

2 Ensemble composition and analysis

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:542-A:577, A:586-A:629 (80)	0.24	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 4 single-model clusters were found.

Cluster number	Models
1	1, 2, 6, 9, 11, 12, 14, 15, 16, 17, 18, 19, 20
2	3, 7, 13
Single-model clusters	4; 5; 8; 10

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1704 atoms, of which 845 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called RNA-binding protein 12.

Mol	Chain	Residues	Atoms						Trace
1	A	114	Total	C	H	N	O	S	0
			1704	526	845	158	172	3	

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	529	GLY	-	CLONING ARTIFACT	UNP Q9NTZ6
A	530	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	531	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	532	GLY	-	CLONING ARTIFACT	UNP Q9NTZ6
A	533	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	534	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	535	GLY	-	CLONING ARTIFACT	UNP Q9NTZ6
A	637	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	638	GLY	-	CLONING ARTIFACT	UNP Q9NTZ6
A	639	PRO	-	CLONING ARTIFACT	UNP Q9NTZ6
A	640	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	641	SER	-	CLONING ARTIFACT	UNP Q9NTZ6
A	642	GLY	-	CLONING ARTIFACT	UNP Q9NTZ6

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA-binding protein 12




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: RNA-binding protein 12




4.2.2 Score per residue for model 2 (medoid)

- Molecule 1: RNA-binding protein 12




4.2.3 Score per residue for model 3

- Molecule 1: RNA-binding protein 12




4.2.4 Score per residue for model 4

- Molecule 1: RNA-binding protein 12




4.2.5 Score per residue for model 5

- Molecule 1: RNA-binding protein 12




4.2.6 Score per residue for model 6

- Molecule 1: RNA-binding protein 12




4.2.7 Score per residue for model 7

- Molecule 1: RNA-binding protein 12




4.2.8 Score per residue for model 8

- Molecule 1: RNA-binding protein 12




4.2.9 Score per residue for model 9

- Molecule 1: RNA-binding protein 12




4.2.10 Score per residue for model 10

- Molecule 1: RNA-binding protein 12




4.2.11 Score per residue for model 11

- Molecule 1: RNA-binding protein 12




4.2.12 Score per residue for model 12

- Molecule 1: RNA-binding protein 12




4.2.13 Score per residue for model 13

- Molecule 1: RNA-binding protein 12




4.2.14 Score per residue for model 14

- Molecule 1: RNA-binding protein 12




4.2.15 Score per residue for model 15

- Molecule 1: RNA-binding protein 12




4.2.16 Score per residue for model 16

- Molecule 1: RNA-binding protein 12




4.2.17 Score per residue for model 17

- Molecule 1: RNA-binding protein 12




4.2.18 Score per residue for model 18

- Molecule 1: RNA-binding protein 12




4.2.19 Score per residue for model 19

- Molecule 1: RNA-binding protein 12



4.2.20 Score per residue for model 20

- Molecule 1: RNA-binding protein 12



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *torsion angle dynamics, restrained molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy, structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.17
CYANA	refinement	2.0.17

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	643	656	656	10±4
All	All	12860	13120	13120	205

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:589:LEU:HD13	1:A:624:MET:CG	0.78	2.08	16	1
1:A:589:LEU:HD22	1:A:624:MET:HG3	0.76	1.58	4	1
1:A:547:HIS:CD2	1:A:589:LEU:CD2	0.69	2.75	12	11
1:A:589:LEU:HD13	1:A:624:MET:HG2	0.66	1.65	16	2
1:A:609:LEU:HD23	1:A:609:LEU:C	0.64	2.12	14	7
1:A:547:HIS:CE1	1:A:549:THR:CG2	0.64	2.81	19	10
1:A:577:LEU:HD12	1:A:587:GLN:HB2	0.62	1.70	15	11
1:A:567:ILE:HG22	1:A:597:ASP:OD2	0.59	1.97	11	1
1:A:567:ILE:O	1:A:569:VAL:HG12	0.59	1.97	3	2
1:A:549:THR:HG22	1:A:587:GLN:HG2	0.59	1.74	4	1
1:A:569:VAL:CG2	1:A:573:ALA:CB	0.58	2.81	7	2
1:A:547:HIS:CD2	1:A:589:LEU:HD23	0.57	2.33	12	8
1:A:590:VAL:HG11	1:A:592:PHE:CZ	0.57	2.35	2	3
1:A:589:LEU:HD12	1:A:628:GLU:OE1	0.56	2.01	10	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:589:LEU:HD13	1:A:624:MET:HG3	0.56	1.75	16	1
1:A:573:ALA:HB1	1:A:591:GLN:O	0.55	2.02	15	9
1:A:549:THR:HG22	1:A:587:GLN:OE1	0.55	2.02	17	1
1:A:590:VAL:CG1	1:A:592:PHE:CZ	0.55	2.89	19	3
1:A:609:LEU:CD2	1:A:609:LEU:C	0.54	2.76	14	6
1:A:609:LEU:C	1:A:609:LEU:HD23	0.54	2.22	13	6
1:A:569:VAL:HG22	1:A:573:ALA:HB3	0.53	1.80	7	3
1:A:561:LEU:HD13	1:A:571:GLU:OE2	0.52	2.03	3	1
1:A:577:LEU:HD12	1:A:587:GLN:CB	0.52	2.34	12	3
1:A:567:ILE:CG2	1:A:597:ASP:OD2	0.49	2.60	11	2
1:A:595:GLU:O	1:A:598:ALA:N	0.49	2.45	5	4
1:A:547:HIS:CE1	1:A:549:THR:HG23	0.48	2.43	14	2
1:A:596:ASP:OD1	1:A:597:ASP:N	0.48	2.46	6	1
1:A:556:THR:O	1:A:559:ASP:N	0.48	2.46	14	4
1:A:598:ALA:O	1:A:601:SER:N	0.48	2.45	2	2
1:A:602:GLU:CG	1:A:618:VAL:HG21	0.48	2.38	12	2
1:A:621:LEU:C	1:A:621:LEU:HD13	0.48	2.29	2	2
1:A:572:ASN:O	1:A:575:HIS:CD2	0.48	2.67	19	1
1:A:567:ILE:HG21	1:A:592:PHE:CE1	0.47	2.44	17	1
1:A:602:GLU:CD	1:A:603:ARG:N	0.47	2.67	5	1
1:A:598:ALA:O	1:A:600:LYS:N	0.47	2.47	2	2
1:A:562:GLN:O	1:A:565:GLU:N	0.47	2.47	18	1
1:A:609:LEU:HD23	1:A:610:ASN:N	0.47	2.24	17	4
1:A:604:LEU:O	1:A:607:LYS:CG	0.47	2.63	14	10
1:A:609:LEU:C	1:A:609:LEU:CD2	0.47	2.83	13	2
1:A:570:ASP:OD2	1:A:573:ALA:N	0.47	2.47	8	1
1:A:596:ASP:CG	1:A:597:ASP:N	0.47	2.69	6	1
1:A:557:LYS:HD2	1:A:576:VAL:HG23	0.46	1.87	13	1
1:A:563:PHE:CD1	1:A:563:PHE:C	0.46	2.89	16	1
1:A:561:LEU:HD11	1:A:571:GLU:HG2	0.46	1.87	2	1
1:A:543:LYS:HG2	1:A:618:VAL:HG12	0.46	1.88	17	1
1:A:600:LYS:O	1:A:603:ARG:N	0.45	2.49	5	1
1:A:595:GLU:O	1:A:597:ASP:N	0.45	2.50	15	2
1:A:607:LYS:O	1:A:614:ALA:HB3	0.44	2.13	8	1
1:A:595:GLU:N	1:A:595:GLU:OE1	0.44	2.50	20	1
1:A:624:MET:SD	1:A:624:MET:C	0.44	2.96	2	2
1:A:602:GLU:HB3	1:A:618:VAL:HG21	0.44	1.88	11	1
1:A:557:LYS:O	1:A:558:MET:C	0.44	2.56	7	2
1:A:558:MET:SD	1:A:558:MET:C	0.44	2.96	18	1
1:A:569:VAL:CG2	1:A:573:ALA:HB3	0.44	2.42	7	2
1:A:550:ASN:ND2	1:A:613:GLU:O	0.44	2.50	3	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:627:ILE:O	1:A:628:GLU:C	0.44	2.56	15	3
1:A:605:HIS:CD2	1:A:606:ARG:HG3	0.43	2.48	8	2
1:A:599:ARG:O	1:A:602:GLU:CG	0.43	2.66	6	1
1:A:560:VAL:O	1:A:563:PHE:N	0.43	2.51	12	1
1:A:561:LEU:HD21	1:A:571:GLU:HA	0.43	1.89	6	1
1:A:602:GLU:C	1:A:602:GLU:CD	0.43	2.77	10	1
1:A:556:THR:O	1:A:557:LYS:C	0.43	2.56	14	8
1:A:621:LEU:C	1:A:621:LEU:CD1	0.43	2.87	2	1
1:A:550:ASN:CG	1:A:613:GLU:O	0.42	2.57	14	1
1:A:552:PRO:HG3	1:A:612:ARG:CD	0.42	2.44	4	1
1:A:557:LYS:CD	1:A:576:VAL:HG23	0.42	2.44	13	1
1:A:615:PHE:CD1	1:A:615:PHE:N	0.42	2.87	20	1
1:A:592:PHE:CE2	1:A:601:SER:HB3	0.42	2.49	20	1
1:A:569:VAL:HG22	1:A:573:ALA:CB	0.42	2.41	7	2
1:A:626:GLU:CG	1:A:627:ILE:N	0.42	2.83	14	1
1:A:569:VAL:HG22	1:A:570:ASP:N	0.42	2.29	3	1
1:A:557:LYS:O	1:A:560:VAL:N	0.42	2.53	7	1
1:A:595:GLU:C	1:A:597:ASP:N	0.42	2.73	15	2
1:A:570:ASP:CG	1:A:573:ALA:HB2	0.41	2.35	8	1
1:A:598:ALA:C	1:A:600:LYS:N	0.41	2.73	2	3
1:A:602:GLU:OE2	1:A:618:VAL:HG21	0.41	2.15	17	1
1:A:598:ALA:O	1:A:599:ARG:C	0.41	2.59	17	2
1:A:586:GLY:O	1:A:587:GLN:CG	0.41	2.68	14	1
1:A:621:LEU:CD1	1:A:621:LEU:C	0.41	2.89	6	1
1:A:547:HIS:CE1	1:A:549:THR:HG22	0.41	2.50	11	1
1:A:594:ASN:OD1	1:A:594:ASN:N	0.41	2.52	11	1
1:A:543:LYS:HE2	1:A:543:LYS:N	0.41	2.31	17	1
1:A:620:THR:O	1:A:621:LEU:C	0.41	2.58	15	1
1:A:560:VAL:O	1:A:561:LEU:C	0.41	2.58	18	1
1:A:593:LYS:CD	1:A:593:LYS:N	0.41	2.84	5	1
1:A:572:ASN:C	1:A:574:VAL:N	0.40	2.75	11	1
1:A:547:HIS:CB	1:A:589:LEU:HD23	0.40	2.46	18	1
1:A:592:PHE:CZ	1:A:601:SER:HB3	0.40	2.52	14	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	80/114 (70%)	71±3 (89±4%)	9±3 (11±4%)	0±1 (0±1%)	38	79
All	All	1600/2280 (70%)	1417 (89%)	175 (11%)	8 (0%)	38	79

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	568	PRO	4
1	A	599	ARG	2
1	A	570	ASP	1
1	A	596	ASP	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	71/95 (75%)	68±2 (96±2%)	3±2 (4±2%)	41	84
All	All	1420/1900 (75%)	1360 (96%)	60 (4%)	41	84

All 25 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	625	ARG	6
1	A	628	GLU	4
1	A	557	LYS	4
1	A	558	MET	4
1	A	624	MET	4
1	A	593	LYS	4
1	A	599	ARG	3
1	A	602	GLU	3
1	A	571	GLU	3
1	A	559	ASP	3
1	A	562	GLN	2
1	A	609	LEU	2
1	A	623	ASP	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	597	ASP	2
1	A	596	ASP	2
1	A	543	LYS	2
1	A	626	GLU	2
1	A	553	PHE	1
1	A	607	LYS	1
1	A	570	ASP	1
1	A	594	ASN	1
1	A	621	LEU	1
1	A	572	ASN	1
1	A	622	GLU	1
1	A	603	ARG	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided