



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:38 PM GMT

PDB ID : 4CTY  
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with (R)-6-(2-Amino-2-(3-(2-(6-amino-4-methylpyridin-2-yl) ethyl)phenyl)ethyl)-4-methylpyridin-2-amine  
Authors : Chreifi, G.; Li, H.; Poulos, T.L.  
Deposited on : 2014-03-15  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	 4% 79% 11% 9%
1	B	443	 3% 82% 8% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X
6	GOL	A	881	-	-	-	X
6	GOL	B	880	-	-	-	X

## 2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 6966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	As	C	N	O				S
1	A	404	Total 3220	As 1	C 2048	N 567	O 588	S 16	0	1	0
1	B	403	Total 3212	As 1	C 2042	N 566	O 587	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	VARIANT	UNP P29473
B	100	ARG	CYS	VARIANT	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is (R)-6-(2-AMINO-2-(3-(2-(6-AMINO-4-METHYLPYRIDIN-2-YL)ETHYL)PHENYL)ETHYL)-4-METHYLPYRIDIN-2-AMINE (three-letter code: S5D) (formula:  $C_{22}H_{27}N_5$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C N	0	0
			27	22 5		
4	B	1	Total	C N	0	0
			27	22 5		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C O	0	0
			4	2 2		
5	B	1	Total	C O	0	0
			4	2 2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Zn 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	180	Total O 180 180	0	0
8	B	153	Total O 153 153	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.85Å 106.50Å 156.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.22 – 2.30 39.19 – 2.29	Depositor EDS
% Data completeness (in resolution range)	98.9 (39.22-2.30) 98.6 (39.19-2.29)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.94 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.161 , 0.219 0.169 , 0.225	Depositor DCC
$R_{free}$ test set	2174 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.7	Xtrriage
Anisotropy	0.443	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 37.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 43624 reflections	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6966	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, S5D, H4B, CAS, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.79	1/3303 (0.0%)	0.84	4/4497 (0.1%)
1	B	0.80	1/3291 (0.0%)	0.85	3/4480 (0.1%)
All	All	0.79	2/6594 (0.0%)	0.85	7/8977 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	76	TRP	CB-CG	-5.78	1.39	1.50
1	B	457	SER	CB-OG	-5.38	1.35	1.42

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331	ARG	NE-CZ-NH1	7.62	124.11	120.30
1	B	240	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	B	240	ARG	NE-CZ-NH1	5.87	123.23	120.30
1	A	328	LEU	CA-CB-CG	5.54	128.03	115.30
1	B	168	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	A	199	ASP	CB-CG-OD1	5.11	122.90	118.30
1	A	331	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3220	0	3127	26	0
1	B	3212	0	3117	16	0
2	A	43	0	30	4	0
2	B	43	0	30	5	0
3	A	17	0	15	1	0
3	B	17	0	15	3	0
4	A	27	0	27	1	0
4	B	27	0	27	1	0
5	A	4	0	3	1	0
5	B	4	0	3	0	0
6	A	12	0	16	0	0
6	B	6	0	8	2	0
7	A	1	0	0	0	0
8	A	180	0	0	2	0
8	B	153	0	0	3	0
All	All	6966	0	6418	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.72	0.70
2:B:500:HEM:HHC	2:B:500:HEM:HBB2	1.84	0.60
1:B:478:GLN:HB2	1:B:479:PRO:HD2	1.86	0.58
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.37	0.55
1:B:233:PHE:HB3	1:B:234:PRO:HD2	1.89	0.54
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.49	0.53
1:B:340:ASN:HD22	1:B:340:ASN:H	1.56	0.52
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.91	0.52
2:B:500:HEM:HBC2	2:B:500:HEM:HMC1	1.93	0.51
1:B:287:ARG:HD3	8:B:2087:HOH:O	2.11	0.51
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.41	0.50
1:B:378:LEU:HB2	8:B:2113:HOH:O	2.11	0.50
1:B:187:VAL:HG22	1:B:187:VAL:O	2.13	0.49
1:B:126:LEU:O	1:B:130:ARG:HG3	2.13	0.49
2:A:500:HEM:CMC	2:A:500:HEM:HBC2	2.44	0.48
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.95	0.48
1:A:378:LEU:HB2	8:A:2139:HOH:O	2.14	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:TRP:HA	3:B:600:H4B:N1	2.28	0.47
1:A:287:ARG:HD3	8:A:2106:HOH:O	2.14	0.47
6:B:880:GOL:H32	8:B:2152:HOH:O	2.15	0.47
1:A:125:LEU:O	1:A:125:LEU:HD23	2.16	0.46
1:A:158:GLU:OE2	1:A:166:HIS:HD2	1.96	0.46
3:B:600:H4B:O4	6:B:880:GOL:O2	2.31	0.46
1:B:246:TRP:CZ2	1:B:302:PRO:HG3	2.51	0.46
1:A:155:VAL:O	1:A:159:VAL:HG23	2.16	0.46
1:B:308:PRO:HD2	1:B:311:LEU:HD12	1.97	0.46
2:A:500:HEM:CBB	2:A:500:HEM:HHC	2.43	0.45
1:A:125:LEU:C	1:A:125:LEU:HD23	2.37	0.45
1:A:100:ARG:NH1	1:A:102:LEU:HD22	2.32	0.44
1:A:449:TRP:HA	3:A:600:H4B:N1	2.32	0.44
1:A:154:GLU:OE1	1:A:168:ARG:NH2	2.51	0.44
2:B:500:HEM:O1A	3:B:600:H4B:N3	2.46	0.43
1:A:455:SER:O	1:A:456:GLY:C	2.56	0.43
1:B:281:TRP:HB2	1:B:304:LEU:HD21	2.01	0.42
1:A:70:PHE:HB3	1:A:84:ASP:O	2.19	0.42
1:A:249:GLN:NE2	4:A:800:S5D:H3	2.34	0.42
1:A:125:LEU:HD21	1:A:165:TYR:CE1	2.54	0.42
1:A:398:ASP:O	1:A:402:VAL:HG23	2.20	0.42
2:B:500:HEM:O1D	4:B:800:S5D:N21	2.53	0.41
1:A:126:LEU:HA	1:A:126:LEU:HD12	1.90	0.41
1:A:97:THR:HB	1:A:98:PRO:HD2	2.02	0.41
1:A:247:ASN:OD1	1:A:247:ASN:N	2.54	0.41
1:A:126:LEU:HD11	1:A:156:GLU:HA	2.02	0.41
1:A:446:ASP:O	1:A:450:ILE:HG12	2.20	0.41
2:A:500:HEM:HMC1	2:A:500:HEM:HBC2	2.01	0.41
1:A:188:GLY:HA2	5:A:860:ACT:CH3	2.51	0.41
1:B:366:THR:O	1:B:370:CYS:HB2	2.20	0.41
1:A:121:PRO:O	1:A:122:ALA:HB3	2.21	0.41
1:A:465:GLU:HB3	1:B:105:LEU:HD22	2.03	0.40
1:A:472:SER:HA	1:A:473:PRO:C	2.42	0.40
1:A:319:HIS:CG	1:A:320:PRO:HD2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/443 (90%)	391 (98%)	8 (2%)	1 (0%)	46	57
1	B	398/443 (90%)	383 (96%)	13 (3%)	2 (0%)	34	41
All	All	798/886 (90%)	774 (97%)	21 (3%)	3 (0%)	39	48

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ALA
1	B	260	ASP
1	B	144	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/375 (92%)	329 (96%)	14 (4%)	37	50
1	B	342/375 (91%)	327 (96%)	15 (4%)	35	46
All	All	685/750 (91%)	656 (96%)	29 (4%)	36	49

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ARG
1	A	102	LEU
1	A	107	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	156	GLU
1	A	194	LYS
1	A	260	ASP
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	390	ARG
1	A	416	LYS
1	A	458	LEU
1	A	468	ASN
1	B	81	ILE
1	B	92	GLN
1	B	100	ARG
1	B	102	LEU
1	B	123	GLU
1	B	124	GLN
1	B	139	SER
1	B	202	ASP
1	B	218	LYS
1	B	259	GLN
1	B	296	GLN
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	389	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	153	GLN
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN
1	A	376	ASN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	405	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CAS	A	384	1	5,8,9	1.36	1 (20%)	2,9,11	1.33	0
1	CAS	B	384	1	5,8,9	1.21	0	2,9,11	1.66	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	0/0/0/0
1	CAS	B	384	1	-	0/0/7/9	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	384	CAS	AS-CE2	2.23	2.01	1.96

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	384	CAS	O-C-CA	-2.28	119.55	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	500	1	30,50,50	2.65	7 (23%)	24,82,82	2.54	11 (45%)
3	H4B	A	600	-	13,18,18	1.47	2 (15%)	11,26,26	3.25	6 (54%)
4	S5D	A	800	-	29,29,29	1.04	2 (6%)	36,40,40	2.00	12 (33%)
5	ACT	A	860	-	1,3,3	1.80	0	0,3,3	0.00	-
6	GOL	A	880	-	5,5,5	0.48	0	5,5,5	0.81	0
6	GOL	A	881	-	5,5,5	0.40	0	5,5,5	0.32	0
2	HEM	B	500	1	30,50,50	2.23	7 (23%)	24,82,82	2.58	10 (41%)
3	H4B	B	600	-	13,18,18	0.91	0	11,26,26	1.92	4 (36%)
4	S5D	B	800	-	29,29,29	1.05	3 (10%)	36,40,40	2.24	8 (22%)
5	ACT	B	860	-	1,3,3	0.16	0	0,3,3	0.00	-
6	GOL	B	880	-	5,5,5	0.50	0	5,5,5	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/10/54/54	0/0/8/8
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	S5D	A	800	-	-	0/13/13/13	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
6	GOL	A	880	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	881	-	-	0/4/4/4	0/0/0/0
2	HEM	B	500	1	-	0/10/54/54	0/0/8/8
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
4	S5D	B	800	-	-	0/13/13/13	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0
6	GOL	B	880	-	-	0/4/4/4	0/0/0/0

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C2D-C3D	-7.31	1.32	1.54
2	A	500	HEM	C3B-C4B	-6.94	1.45	1.51
2	B	500	HEM	C2D-C3D	-6.48	1.35	1.54
2	B	500	HEM	C2C-C1C	-6.15	1.40	1.52
2	A	500	HEM	C2C-C1C	-5.68	1.41	1.52
2	A	500	HEM	C3D-C4D	-4.43	1.45	1.51
2	A	500	HEM	C3C-CAC	-3.97	1.43	1.51
2	B	500	HEM	C3D-C4D	-3.97	1.46	1.51
4	A	800	S5D	C6-C29	-3.44	1.47	1.51
2	A	500	HEM	C2B-C1B	-3.36	1.40	1.51
4	B	800	S5D	C6-C29	-3.06	1.47	1.51
2	B	500	HEM	C3B-C4B	-2.98	1.49	1.51
3	A	600	H4B	C8A-N1	-2.35	1.30	1.34
2	B	500	HEM	C2B-C1B	-2.26	1.44	1.51
2	B	500	HEM	C3C-CAC	-2.13	1.47	1.51
4	A	800	S5D	C15-C16	2.09	1.42	1.38
4	B	800	S5D	C26-N21	2.20	1.38	1.34
2	A	500	HEM	FE-NC	2.35	2.05	1.95
3	A	600	H4B	C7-N8	2.37	1.49	1.46
4	B	800	S5D	C23-C24	2.45	1.43	1.38
2	B	500	HEM	FE-NC	3.11	2.08	1.95

All (51) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	S5D	C24-C25-C26	-5.91	116.58	120.28
2	B	500	HEM	CBA-CAA-C2A	-4.89	103.76	112.53
3	A	600	H4B	N3-C2-N1	-4.63	117.94	125.53
2	A	500	HEM	C1D-CHD-C4C	-4.08	119.01	125.82
4	B	800	S5D	C14-C15-C16	-3.76	117.92	120.28
4	B	800	S5D	C15-C16-N11	-3.52	118.98	122.96
2	B	500	HEM	C1D-CHD-C4C	-3.34	120.23	125.82

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	S5D	C28-C29-C6	-3.14	104.75	111.78
2	A	500	HEM	CBA-CAA-C2A	-2.96	107.22	112.53
4	A	800	S5D	C15-C16-N11	-2.34	120.31	122.96
2	A	500	HEM	C3C-CAC-CBC	-2.33	120.89	124.46
2	A	500	HEM	C3B-C4B-NB	-2.25	107.34	111.63
4	A	800	S5D	C6-C1-C2	-2.17	117.59	120.96
3	B	600	H4B	N3-C2-N1	-2.17	121.97	125.53
4	B	800	S5D	C27-C24-C25	-2.14	117.69	120.95
4	A	800	S5D	C27-C24-C23	-2.13	117.72	120.95
2	B	500	HEM	C3B-C4B-NB	-2.06	107.69	111.63
2	B	500	HEM	C3C-CAC-CBC	-2.01	121.37	124.46
4	A	800	S5D	N22-C22-N21	2.00	120.15	116.50
4	A	800	S5D	C25-C24-C23	2.11	120.78	118.08
4	B	800	S5D	N12-C12-N11	2.18	120.47	116.50
4	A	800	S5D	C5-C6-C1	2.22	121.46	118.79
2	A	500	HEM	CMD-C2D-C3D	2.26	124.35	114.35
4	B	800	S5D	C15-C14-C13	2.28	121.00	118.08
2	A	500	HEM	C2D-C3D-C4D	2.45	105.65	101.50
4	A	800	S5D	C12-N11-C16	2.51	120.01	118.23
2	B	500	HEM	CMD-C2D-C3D	2.55	125.61	114.35
4	A	800	S5D	C18-C16-C15	2.56	124.71	121.13
3	B	600	H4B	N2-C2-N3	2.78	121.80	117.20
4	A	800	S5D	C1-C6-C29	2.85	123.74	120.04
3	B	600	H4B	C2-N1-C8A	3.01	121.30	114.54
3	A	600	H4B	C2-N1-C8A	3.39	122.17	114.54
2	A	500	HEM	CAA-CBA-CGA	3.40	118.98	112.75
4	B	800	S5D	C22-N21-C26	3.51	120.72	118.23
2	B	500	HEM	CAA-CBA-CGA	3.53	119.21	112.75
3	B	600	H4B	C4-C4A-C8A	3.55	117.78	114.56
4	B	800	S5D	C18-C16-C15	3.66	126.26	121.13
2	B	500	HEM	CMB-C2B-C3B	3.77	125.93	116.53
2	A	500	HEM	CAD-C3D-C2D	3.78	124.09	113.22
3	A	600	H4B	C4-C4A-C8A	4.16	118.33	114.56
3	A	600	H4B	C4A-C8A-N8	4.21	123.39	118.43
2	B	500	HEM	CAD-C3D-C2D	4.26	125.46	113.22
2	A	500	HEM	CMB-C2B-C3B	4.37	127.43	116.53
2	B	500	HEM	CMC-C2C-C3C	4.42	127.56	116.53
3	A	600	H4B	N2-C2-N3	4.49	124.64	117.20
2	B	500	HEM	CAD-C3D-C4D	4.88	129.67	112.47
2	A	500	HEM	CMC-C2C-C3C	4.97	128.93	116.53
2	A	500	HEM	CAD-C3D-C4D	5.04	130.26	112.47
3	A	600	H4B	C4-N3-C2	5.07	122.97	115.94

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	S5D	C22-N21-C26	5.56	122.18	118.23
4	B	800	S5D	C12-N11-C16	9.51	124.99	118.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	4	0
3	A	600	H4B	1	0
4	A	800	S5D	1	0
5	A	860	ACT	1	0
2	B	500	HEM	5	0
3	B	600	H4B	3	0
4	B	800	S5D	1	0
6	B	880	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	403/443 (90%)	-0.04	16 (3%) 42 51	23, 36, 65, 93	0
1	B	402/443 (90%)	-0.09	12 (2%) 54 63	23, 39, 68, 103	0
All	All	805/886 (90%)	-0.07	28 (3%) 48 56	23, 37, 67, 103	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	GLN	7.6
1	A	67	GLY	5.1
1	A	160	ALA	5.1
1	A	239	GLY	4.7
1	A	259	GLN	4.2
1	B	121	PRO	3.9
1	A	156	GLU	3.3
1	B	109	ARG	3.3
1	A	69	LYS	3.2
1	A	153	GLN	3.1
1	A	123	GLU	3.1
1	A	91	GLN	2.9
1	B	141	LYS	2.9
1	B	142	ARG	2.9
1	B	260	ASP	2.9
1	B	144	GLY	2.8
1	A	126	LEU	2.7
1	A	124	GLN	2.6
1	A	122	ALA	2.6
1	A	121	PRO	2.4
1	B	123	GLU	2.4
1	B	239	GLY	2.3
1	A	338	VAL	2.3
1	A	163	GLY	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	146	GLN	2.1
1	B	143	SER	2.1
1	A	159	VAL	2.0
1	B	223	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
1	CAS	A	384	9/10	0.98	0.07	-	36,36,59,60	0
1	CAS	B	384	9/10	0.99	0.08	-	44,46,56,62	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
5	ACT	B	860	4/4	0.76	0.22	7.12	37,47,48,52	0
6	GOL	A	881	6/6	0.86	0.37	6.50	59,67,72,75	0
5	ACT	A	860	4/4	0.87	0.20	4.41	35,38,41,44	0
6	GOL	B	880	6/6	0.95	0.25	3.31	47,50,52,56	0
6	GOL	A	880	6/6	0.94	0.20	1.49	34,43,45,50	0
3	H4B	B	600	17/17	0.96	0.17	1.04	28,29,33,33	0
4	S5D	B	800	27/27	0.94	0.18	0.88	30,35,37,39	0
4	S5D	A	800	27/27	0.95	0.20	0.77	24,35,39,40	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	H4B	A	600	17/17	0.98	0.18	0.71	24,26,30,30	0
2	HEM	A	500	43/43	0.97	0.17	0.60	27,30,36,38	0
2	HEM	B	500	43/43	0.98	0.14	0.30	25,29,35,40	0
7	ZN	A	900	1/1	1.00	0.08	-0.93	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.