



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:43 PM GMT

PDB ID : 4CYI
Title : Chaetomium thermophilum Pan3
Authors : Wolf, J.; Valkov, E.; Allen, M.D.; Meineke, B.; Gordiyenko, Y.; McLaughlin, S.H.; Olsen, T.M.; Robinson, C.V.; Bycroft, M.; Stewart, M.; Passmore, L.A.
Deposited on : 2014-04-11
Resolution : 2.42 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3386 (2.44-2.40)
Clashscore	102246	3897 (2.44-2.40)
Ramachandran outliers	100387	3837 (2.44-2.40)
Sidechain outliers	100360	3838 (2.44-2.40)
RSRZ outliers	91569	3396 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	438	<div> <div>5%</div> <div>87%</div> <div>5% • 8%</div> </div>
1	B	438	<div> <div>6%</div> <div>84%</div> <div>8% • 7%</div> </div>
1	C	438	<div> <div>9%</div> <div>86%</div> <div>• 9%</div> </div>
1	D	438	<div> <div>6%</div> <div>87%</div> <div>5% 7%</div> </div>
1	E	438	<div> <div>13%</div> <div>85%</div> <div>• • 10%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	438	
1	G	438	
1	H	438	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 53391 atoms, of which 26174 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	405	Total	C	H	N	O	S	Se	0	0	0
			6562	2080	3289	581	598	5	9			
1	B	406	Total	C	H	N	O	S	Se	0	0	0
			6565	2082	3290	577	602	5	9			
1	C	397	Total	C	H	N	O	S	Se	0	0	0
			6421	2036	3219	564	588	5	9			
1	D	407	Total	C	H	N	O	S	Se	0	0	0
			6560	2082	3284	575	605	5	9			
1	E	394	Total	C	H	N	O	S	Se	0	0	0
			6386	2024	3202	561	585	5	9			
1	F	408	Total	C	H	N	O	S	Se	0	0	0
			6584	2088	3297	579	606	5	9			
1	G	397	Total	C	H	N	O	S	Se	0	0	0
			6429	2035	3223	568	589	5	9			
1	H	407	Total	C	H	N	O	S	Se	0	0	0
			6559	2083	3282	575	605	5	9			

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	B	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	C	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	D	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	E	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	F	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	G	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		
2	H	1	Total	C	H	N	O	P	0	0
			42	10	11	5	13	3		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	E	1	Total	Mg	0	0
			1	1		
3	H	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	C	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0
3	F	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	183	Total 183	O 183	0	0
4	B	129	Total 129	O 129	0	0
4	C	110	Total 110	O 110	0	0
4	D	149	Total 149	O 149	0	0
4	E	79	Total 79	O 79	0	0
4	F	92	Total 92	O 92	0	0
4	G	112	Total 112	O 112	0	0
4	H	127	Total 127	O 127	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN




- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN




- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN











- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN




- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN




- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN



- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN



- Molecule 1: PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-LIKE PROTEIN, PAB-DEPENDENT POLY(A)-SPECIFIC RIBONUCLEASE SUBUNIT PAN3-L PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.52Å 146.04Å 149.93Å 89.81° 81.13° 82.77°	Depositor
Resolution (Å)	52.21 – 2.42 52.21 – 2.42	Depositor EDS
% Data completeness (in resolution range)	98.4 (52.21-2.42) 97.1 (52.21-2.42)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.78 (at 2.42Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.190 , 0.234 0.200 , 0.240	Depositor DCC
R_{free} test set	8073 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.894	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 63.3	EDS
Estimated twinning fraction	0.014 for -h,-k,-h+l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 159442 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	53391	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/3334	0.46	0/4495
1	B	0.26	0/3334	0.45	0/4494
1	C	0.25	0/3259	0.43	0/4391
1	D	0.26	0/3335	0.44	0/4496
1	E	0.25	0/3239	0.43	0/4362
1	F	0.25	0/3346	0.44	0/4510
1	G	0.25	0/3260	0.44	0/4389
1	H	0.26	0/3338	0.46	0/4502
All	All	0.26	0/26445	0.44	0/35639

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3273	3289	3275	17	0
1	B	3275	3290	3277	20	0
1	C	3202	3219	3206	10	0
1	D	3276	3284	3271	13	0
1	E	3184	3202	3189	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3287	3297	3284	15	0
1	G	3206	3223	3210	9	0
1	H	3277	3282	3269	14	0
2	A	31	11	12	1	0
2	B	31	11	12	0	0
2	C	31	11	12	1	0
2	D	31	11	12	2	0
2	E	31	11	12	0	0
2	F	31	11	12	0	0
2	G	31	11	12	0	0
2	H	31	11	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	A	183	0	0	2	0
4	B	129	0	0	3	0
4	C	110	0	0	6	0
4	D	149	0	0	2	0
4	E	79	0	0	2	0
4	F	92	0	0	1	0
4	G	112	0	0	1	0
4	H	127	0	0	1	0
All	All	27217	26174	26077	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:227:ASP:OD2	1:B:231:LYS:NZ	2.20	0.75
1:F:260:ASN:ND2	1:F:260:ASN:O	2.20	0.75
1:F:225:ARG:NH1	4:F:2005:HOH:O	2.26	0.69
1:A:398:LYS:NZ	2:A:900:ATP:O1G	2.27	0.67
4:C:2072:HOH:O	1:D:513:ASP:OD1	2.12	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/438 (92%)	396 (99%)	5 (1%)	0	100	100
1	B	400/438 (91%)	397 (99%)	3 (1%)	0	100	100
1	C	391/438 (89%)	389 (100%)	2 (0%)	0	100	100
1	D	401/438 (92%)	397 (99%)	4 (1%)	0	100	100
1	E	388/438 (89%)	386 (100%)	2 (0%)	0	100	100
1	F	402/438 (92%)	398 (99%)	4 (1%)	0	100	100
1	G	391/438 (89%)	386 (99%)	5 (1%)	0	100	100
1	H	401/438 (92%)	398 (99%)	3 (1%)	0	100	100
All	All	3175/3504 (91%)	3147 (99%)	28 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/376 (96%)	349 (97%)	11 (3%)	47	68
1	B	361/376 (96%)	351 (97%)	10 (3%)	51	71
1	C	353/376 (94%)	346 (98%)	7 (2%)	63	80
1	D	361/376 (96%)	356 (99%)	5 (1%)	74	88

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	351/376 (93%)	341 (97%)	10 (3%)	51	71
1	F	362/376 (96%)	351 (97%)	11 (3%)	48	69
1	G	352/376 (94%)	344 (98%)	8 (2%)	58	77
1	H	362/376 (96%)	351 (97%)	11 (3%)	48	69
All	All	2862/3008 (95%)	2789 (97%)	73 (3%)	54	74

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	259	SER
1	E	604	ASP
1	H	388	ARG
1	E	540	GLU
1	E	629	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	605	ASN
1	F	237	GLN
1	F	284	HIS
1	E	568	HIS
1	F	246	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	A	900	3	24,33,33	0.86	1 (4%)	31,52,52	1.99	6 (19%)
2	ATP	B	900	3	24,33,33	0.84	1 (4%)	31,52,52	1.99	5 (16%)
2	ATP	C	900	3	24,33,33	0.87	1 (4%)	31,52,52	2.01	5 (16%)
2	ATP	D	900	3	24,33,33	0.89	1 (4%)	31,52,52	2.04	6 (19%)
2	ATP	E	900	3	24,33,33	0.95	1 (4%)	31,52,52	1.93	5 (16%)
2	ATP	F	900	3	24,33,33	0.84	1 (4%)	31,52,52	2.12	6 (19%)
2	ATP	G	900	3	24,33,33	0.91	1 (4%)	31,52,52	2.04	5 (16%)
2	ATP	H	900	3	24,33,33	0.87	1 (4%)	31,52,52	2.05	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	900	3	-	0/18/38/38	0/3/3/3
2	ATP	B	900	3	-	0/18/38/38	0/3/3/3
2	ATP	C	900	3	-	0/18/38/38	0/3/3/3
2	ATP	D	900	3	-	0/18/38/38	0/3/3/3
2	ATP	E	900	3	-	0/18/38/38	0/3/3/3
2	ATP	F	900	3	-	0/18/38/38	0/3/3/3
2	ATP	G	900	3	-	0/18/38/38	0/3/3/3
2	ATP	H	900	3	-	0/18/38/38	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	ATP	C5-C4	2.45	1.46	1.40
2	B	900	ATP	C5-C4	2.51	1.46	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	900	ATP	C5-C4	2.62	1.46	1.40
2	F	900	ATP	C5-C4	2.62	1.46	1.40
2	H	900	ATP	C5-C4	2.62	1.46	1.40

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	900	ATP	N3-C2-N1	-8.67	122.25	128.89
2	F	900	ATP	N3-C2-N1	-8.29	122.55	128.89
2	H	900	ATP	N3-C2-N1	-8.08	122.71	128.89
2	G	900	ATP	N3-C2-N1	-7.97	122.79	128.89
2	C	900	ATP	N3-C2-N1	-7.94	122.81	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	900	ATP	1	0
2	C	900	ATP	1	0
2	D	900	ATP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	396/438 (90%)	0.64	23 (5%) 26 26	23, 52, 110, 169	0
1	B	397/438 (90%)	0.63	27 (6%) 20 20	21, 60, 126, 190	0
1	C	388/438 (88%)	0.83	39 (10%) 9 8	36, 74, 128, 167	0
1	D	398/438 (90%)	0.60	27 (6%) 20 20	27, 58, 115, 167	0
1	E	385/438 (87%)	1.04	57 (14%) 3 3	42, 85, 149, 205	0
1	F	399/438 (91%)	0.68	30 (7%) 17 16	33, 62, 135, 191	0
1	G	388/438 (88%)	0.87	41 (10%) 8 7	30, 75, 131, 180	0
1	H	398/438 (90%)	0.54	20 (5%) 32 32	23, 54, 119, 169	0
All	All	3149/3504 (89%)	0.73	264 (8%) 14 13	21, 64, 130, 205	0

The worst 5 of 264 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	260	ASN	11.9
1	E	363	PRO	7.9
1	H	296	THR	7.5
1	C	261	ARG	7.2
1	H	295	LEU	7.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ATP	A	900	31/31	0.96	0.26	1.36	22,35,69,135	0
3	MG	H	1000	1/1	0.79	0.21	1.28	42,42,42,42	0
2	ATP	D	900	31/31	0.98	0.19	0.26	19,36,43,50	0
2	ATP	G	900	31/31	0.95	0.20	0.05	35,51,73,158	0
3	MG	D	1000	1/1	0.95	0.20	0.01	29,29,29,29	0
2	ATP	C	900	31/31	0.94	0.20	-0.09	32,50,71,128	0
2	ATP	H	900	31/31	0.98	0.17	-0.19	25,39,50,60	0
2	ATP	E	900	31/31	0.92	0.18	-0.27	39,60,79,122	0
2	ATP	B	900	31/31	0.97	0.17	-0.31	30,42,62,76	0
2	ATP	F	900	31/31	0.97	0.16	-0.46	34,46,59,67	0
3	MG	B	1000	1/1	0.80	0.15	-1.06	38,38,38,38	0
3	MG	E	1000	1/1	0.87	0.22	-	54,54,54,54	0
3	MG	C	1000	1/1	0.86	0.24	-	56,56,56,56	0
3	MG	A	1000	1/1	0.93	0.35	-	41,41,41,41	0
3	MG	F	1000	1/1	0.94	0.14	-	52,52,52,52	0
3	MG	G	1000	1/1	0.96	0.28	-	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.