



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:25 AM GMT

PDB ID : 2CZ0
Title : photo-activation state of Fe-type NHase in aerobic condition
Authors : Kawano, Y.; Hashimoto, K.; Odaka, M.; Nakayama, H.; Takio, K.; Endo, I.; Kamiya, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2005-07-09
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	206	 82% 12% . .
2	B	212	 83% 16% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3603 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitrile hydratase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	S	0	0	0
			1548	985	260	297	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	112	CSD	CYS	MODIFIED RESIDUE	UNP P13448
A	114	CSD	CYS	MODIFIED RESIDUE	UNP P13448


- Molecule 2 is a protein called Nitrile hydratase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	211	Total	C	N	O	S	0	0	0
			1651	1047	282	314	8			

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Fe	0	0
			1	1		

- Molecule 4 is BUTANOIC ACID (three-letter code: BUA) (formula: C₄H₈O₂).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			12	8	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	185	Total	O	0	0
			185	185		
5	B	206	Total	O	0	0
			206	206		

Note EDS was not executed.

- Chain A:  82% 12% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.02Å 60.18Å 81.57Å 90.00° 125.15° 90.00°	Depositor
Resolution (Å)	50.00 – 1.50	Depositor
% Data completeness (in resolution range)	79.5 (50.00-1.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.173 , 0.214	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3603	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, BUA, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/1571	1.19	9/2147 (0.4%)
2	B	0.51	0/1699	1.22	12/2311 (0.5%)
All	All	0.49	0/3270	1.20	21/4458 (0.5%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	138	ARG	NE-CZ-NH2	-8.19	116.21	120.30
2	B	61	ARG	NE-CZ-NH1	7.91	124.26	120.30
1	A	135	ARG	NE-CZ-NH2	-7.89	116.36	120.30
2	B	76	TYR	CB-CG-CD2	-7.71	116.37	121.00
2	B	37	TYR	CB-CG-CD2	-7.64	116.41	121.00
2	B	107	ARG	NE-CZ-NH1	7.14	123.87	120.30
2	B	76	TYR	CB-CG-CD1	6.74	125.05	121.00
1	A	174	ARG	NE-CZ-NH2	-6.48	117.06	120.30
2	B	67	TYR	CB-CG-CD1	6.30	124.78	121.00
2	B	207	TYR	CB-CG-CD2	-6.01	117.39	121.00
1	A	62	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	A	99	ASP	CB-CG-OD2	-5.89	113.00	118.30
2	B	145	TYR	CD1-CE1-CZ	-5.73	114.64	119.80
1	A	161	ASP	CB-CG-OD2	-5.57	113.28	118.30
1	A	55	ARG	NE-CZ-NH1	-5.49	117.55	120.30
1	A	160	TYR	CB-CG-CD2	-5.43	117.74	121.00
2	B	114	ARG	NE-CZ-NH1	-5.35	117.62	120.30
2	B	207	TYR	CB-CG-CD1	5.25	124.15	121.00
1	A	138	ARG	NE-CZ-NH1	5.23	122.92	120.30
2	B	65	ARG	NE-CZ-NH2	-5.10	117.75	120.30
2	B	6	ASP	CB-CG-OD2	-5.05	113.76	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1548	0	1514	15	0
2	B	1651	0	1567	18	0
3	A	1	0	0	0	0
4	A	12	0	14	1	0
5	A	185	0	0	5	0
5	B	206	0	0	0	0
All	All	3603	0	3095	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:ALA:HA	5:A:1479:HOH:O	1.86	0.73
2:B:129:VAL:HG21	2:B:208:LEU:HB3	1.69	0.73
1:A:137:VAL:HG23	5:A:1367:HOH:O	1.88	0.72
4:A:1301[B]:BUA:H31	2:B:40:MET:SD	2.31	0.71
1:A:146:GLU:HB3	2:B:15:LYS:NZ	2.07	0.69
2:B:129:VAL:HG23	2:B:209:GLU:O	1.94	0.67
1:A:183:GLN:O	1:A:187:GLN:HG3	2.03	0.58
1:A:184:GLU:HB2	5:A:1357:HOH:O	2.01	0.58
2:B:129:VAL:CG2	2:B:208:LEU:HB3	2.34	0.58
1:A:10:ASN:N	1:A:10:ASN:HD22	2.02	0.58
1:A:146:GLU:HB3	2:B:15:LYS:HZ3	1.70	0.55
1:A:93:TYR:OH	2:B:155:HIS:HE1	1.93	0.51
1:A:184:GLU:O	1:A:188:GLU:OE2	2.30	0.50
5:A:1342:HOH:O	2:B:34:HIS:HE1	1.94	0.49
1:A:82:ALA:HB2	1:A:87:LEU:HD11	1.94	0.48
1:A:124:PRO:HD3	1:A:192:LYS:HE3	1.95	0.48
2:B:117:PRO:HG2	2:B:206:GLY:HA3	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:49:ALA:O	2:B:87:LYS:HE3	2.14	0.47
2:B:34:HIS:CD2	2:B:34:HIS:H	2.33	0.47
1:A:141:ARG:NH1	5:A:1438:HOH:O	2.49	0.45
1:A:188:GLU:N	1:A:188:GLU:OE2	2.50	0.44
2:B:163:PHE:HA	2:B:164:PRO:HD3	1.92	0.42
2:B:186:ALA:HB1	2:B:194:THR:HG21	2.00	0.42
2:B:165:ASP:OD1	2:B:165:ASP:N	2.53	0.42
1:A:9:GLU:OE2	1:A:10:ASN:N	2.52	0.42
2:B:65:ARG:O	2:B:69:MET:HG3	2.19	0.41
2:B:161:TRP:HB3	2:B:162:PRO:HD2	2.02	0.41
2:B:121:THR:HB	2:B:123:PHE:CZ	2.56	0.41
1:A:108:VAL:HG22	1:A:109:CYS:N	2.35	0.41
2:B:141:ARG:HA	2:B:141:ARG:HD2	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/206 (94%)	189 (98%)	3 (2%)	1 (0%)	34	10
2	B	209/212 (99%)	205 (98%)	4 (2%)	0	100	100
All	All	402/418 (96%)	394 (98%)	7 (2%)	1 (0%)	52	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/170 (95%)	156 (97%)	5 (3%)	47	14
2	B	173/173 (100%)	168 (97%)	5 (3%)	50	16
All	All	334/343 (97%)	324 (97%)	10 (3%)	48	15

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLU
1	A	153	SER
1	A	154	ASP
1	A	184	GLU
1	A	188	GLU
2	B	103	PHE
2	B	165	ASP
2	B	167	ILE
2	B	172	ASN
2	B	177	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	GLN
1	A	105	ASN
1	A	202	GLN
2	B	21	ASN
2	B	34	HIS
2	B	127	GLN
2	B	155	HIS
2	B	172	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	A	112	1,3	3,7,8	0.62	0	3,8,10	3.10	2 (66%)
1	CSD	A	114	1,3	3,7,8	0.81	0	3,8,10	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	112	1,3	-	0/2/6/8	0/0/0/0
1	CSD	A	114	1,3	-	0/2/6/8	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	112	CSD	O-C-CA	-3.69	115.89	125.49
1	A	112	CSD	OD1-SG-CB	3.87	111.84	105.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BUA	A	1301[A]	-	2,5,5	0.36	0	1,5,5	0.82	0
4	BUA	A	1301[B]	-	2,5,5	0.28	0	1,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BUA	A	1301[A]	-	-	0/1/3/3	0/0/0/0
4	BUA	A	1301[B]	-	-	0/1/3/3	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1301[B]	BUA	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.