



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:40 PM GMT

PDB ID : 4D06
Title : Bacterial chalcone isomerase complexed with naringenin
Authors : Thomsen, M.; Palm, G.J.; Hinrichs, W.
Deposited on : 2014-04-24
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	283	<div> <div>6%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	B	283	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>7%</div> <div>8%</div> </div> </div>
1	C	283	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>6%</div> </div> </div>
1	D	283	<div> <div></div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>
1	E	283	<div> <div>7%</div> <div> <div></div> <div>93%</div> <div>7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	283	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAR	D	301	-	-	-	X
2	NAR	F	301	-	-	-	X
2	NAR	F	302[A]	-	-	-	X
2	NAR	F	302[B]	-	-	-	X
4	X8W	B	301	-	-	-	X
4	X8W	D	303[A]	-	-	-	X
5	GOL	B	302	-	-	-	X
5	GOL	F	303	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 15609 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHALCONE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	5	0
			2312	1495	380	423	14			
1	B	259	Total	C	N	O	S	0	8	0
			2172	1421	349	387	15			
1	C	282	Total	C	N	O	S	0	7	0
			2326	1509	380	423	14			
1	D	259	Total	C	N	O	S	0	8	0
			2170	1422	349	384	15			
1	E	282	Total	C	N	O	S	0	8	0
			2331	1509	381	426	15			
1	F	259	Total	C	N	O	S	0	6	0
			2161	1417	348	382	14			

- Molecule 2 is NARINGENIN (three-letter code: NAR) (formula: C₁₅H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			20	15	5		
2	C	1	Total	C	O	0	0
			20	15	5		
2	D	1	Total	C	O	0	0
			20	15	5		
2	D	1	Total	C	O	0	0
			20	15	5		
2	E	1	Total	C	O	0	0
			20	15	5		
2	F	1	Total	C	O	0	0
			20	15	5		
2	F	1	Total	C	O	0	1
			19	12	7		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	3	Total	Cl	0	0
			3	3		
3	E	4	Total	Cl	0	0
			4	4		
3	B	2	Total	Cl	0	0
			2	2		
3	C	3	Total	Cl	0	0
			3	3		
3	A	2	Total	Cl	0	0
			2	2		
3	F	3	Total	Cl	0	0
			3	3		

- Molecule 4 is (2E)-3-(4-HYDROXYPHENYL)-1-(2,4,6-TRIHYDROXYPHENYL)PROP-2-EN-1-ONE (three-letter code: X8W) (formula: C₁₅H₁₂O₅).



- GOL
-
- The diagram shows a skeletal structure of 1,2,3-propanetriol (glycerol). The carbon atoms are labeled C1, C2, and C3 in green. The hydroxyl groups are shown as HO (red) and OH (red). The labels O1, O2, and O3 are in green below the respective oxygen atoms. The structure is drawn with black lines for the carbon backbone and red lines for the C-O and O-H bonds.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 6 3 3	0	0
5	F	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	357	Total 361	O 361	0	4
6	B	325	Total 329	O 329	0	4
6	C	345	Total 347	O 347	0	2
6	D	294	Total 297	O 297	0	3
6	E	310	Total 312	O 312	0	2
6	F	264	Total 267	O 267	0	3


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CHALCONE ISOMERASE




- Molecule 1: CHALCONE ISOMERASE



- Molecule 1: CHALCONE ISOMERASE




- Molecule 1: CHALCONE ISOMERASE





- Molecule 1: CHALCONE ISOMERASE





● Molecule 1: CHALCONE ISOMERASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	181.16Å 187.99Å 196.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.18 – 2.00 49.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.18-2.00) 99.8 (49.18-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.140 , 0.163 0.151 , 0.170	Depositor DCC
R_{free} test set	11199 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.4	EDS
Estimated twinning fraction	0.012 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 224291 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	15609	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAR, GOL, X8W, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.92	1/2400 (0.0%)	0.96	6/3262 (0.2%)
1	B	0.95	3/2265 (0.1%)	1.00	7/3078 (0.2%)
1	C	1.04	4/2421 (0.2%)	0.97	7/3291 (0.2%)
1	D	1.01	1/2266 (0.0%)	1.05	14/3077 (0.5%)
1	E	0.96	1/2425 (0.0%)	1.00	8/3295 (0.2%)
1	F	0.99	2/2254 (0.1%)	0.98	5/3064 (0.2%)
All	All	0.98	12/14031 (0.1%)	0.99	47/19067 (0.2%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	207	ASP	CB-CG	8.12	1.68	1.51
1	C	177	GLU	CD-OE2	7.09	1.33	1.25
1	C	179	GLU	CD-OE1	-7.07	1.17	1.25
1	B	207	ASP	CB-CG	7.04	1.66	1.51
1	C	19	GLU	CD-OE2	5.83	1.32	1.25

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	207	ASP	CB-CG-OD1	9.96	127.27	118.30
1	D	9	ARG	NE-CZ-NH1	9.35	124.97	120.30
1	D	162	ARG	NE-CZ-NH1	9.25	124.92	120.30
1	A	162	ARG	NE-CZ-NH2	-9.07	115.76	120.30
1	A	162	ARG	NE-CZ-NH1	8.82	124.71	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2312	0	2229	8	0
1	B	2172	0	2113	11	0
1	C	2326	0	2251	5	0
1	D	2170	0	2123	11	0
1	E	2331	0	2254	7	0
1	F	2161	0	2113	8	0
2	A	20	0	9	3	0
2	C	20	0	9	1	0
2	D	40	0	22	5	0
2	E	20	0	9	1	0
2	F	39	0	15	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
3	E	4	0	0	0	0
3	F	3	0	0	0	0
4	B	20	0	9	5	0
4	D	36	0	18	8	0
5	B	6	0	8	2	0
5	F	6	0	8	0	0
6	A	361	0	0	4	0
6	B	329	0	0	12	0
6	C	347	0	0	1	0
6	D	297	0	0	7	0
6	E	312	0	0	4	0
6	F	267	0	0	1	0
All	All	15609	0	13190	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:303[B]:X8W:OAB	6:D:2240:HOH:O	1.73	1.04
1:B:33[A]:HIS:NE2	6:B:2070[A]:HOH:O	1.58	0.93
4:B:301:X8W:HAM	5:B:302:GOL:O1	1.70	0.91
4:B:301:X8W:OAD	6:B:2161:HOH:O	1.92	0.88
2:A:301:NAR:C15	6:A:2090:HOH:O	2.27	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/283 (101%)	280 (98%)	5 (2%)	0	100	100
1	B	263/283 (93%)	259 (98%)	4 (2%)	0	100	100
1	C	287/283 (101%)	281 (98%)	5 (2%)	1 (0%)	46	41
1	D	263/283 (93%)	260 (99%)	3 (1%)	0	100	100
1	E	288/283 (102%)	283 (98%)	5 (2%)	0	100	100
1	F	262/283 (93%)	259 (99%)	3 (1%)	0	100	100
All	All	1648/1698 (97%)	1622 (98%)	25 (2%)	1 (0%)	56	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	207	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/241 (102%)	242 (99%)	3 (1%)	78	81
1	B	233/241 (97%)	230 (99%)	3 (1%)	76	79
1	C	247/241 (102%)	242 (98%)	5 (2%)	63	65
1	D	233/241 (97%)	231 (99%)	2 (1%)	84	88
1	E	248/241 (103%)	246 (99%)	2 (1%)	86	89
1	F	232/241 (96%)	231 (100%)	1 (0%)	93	95
All	All	1438/1446 (99%)	1422 (99%)	16 (1%)	80	83

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	93[A]	PHE
1	C	93[B]	PHE
1	D	137	PHE
1	C	41	PHE
1	E	41	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	101	GLN
1	C	133	ASN
1	D	133	ASN
1	B	191	GLN
1	D	101	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 30 ligands modelled in this entry, 17 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAR	A	301	-	22,22,22	1.57	3 (13%)	32,32,32	1.96	9 (28%)
4	X8W	B	301	-	21,21,21	2.40	6 (28%)	29,29,29	1.49	7 (24%)
5	GOL	B	302	-	5,5,5	0.57	0	5,5,5	1.73	1 (20%)
2	NAR	C	301	-	22,22,22	1.92	4 (18%)	32,32,32	2.24	11 (34%)
2	NAR	D	301	-	22,22,22	2.76	8 (36%)	32,32,32	2.53	10 (31%)
2	NAR	D	302	-	22,22,22	3.27	6 (27%)	32,32,32	5.92	15 (46%)
4	X8W	D	303[A]	-	18,18,21	1.18	2 (11%)	23,23,29	1.77	4 (17%)
4	X8W	D	303[B]	-	18,18,21	1.16	2 (11%)	23,23,29	1.73	2 (8%)
2	NAR	E	301	-	22,22,22	1.66	4 (18%)	32,32,32	2.50	14 (43%)
2	NAR	F	301	-	22,22,22	1.21	2 (9%)	32,32,32	2.24	7 (21%)
2	NAR	F	302[A]	-	9,9,22	2.69	2 (22%)	12,12,32	1.11	1 (8%)
2	NAR	F	302[B]	-	9,9,22	3.20	2 (22%)	12,12,32	1.02	1 (8%)
5	GOL	F	303	-	5,5,5	1.04	0	5,5,5	1.80	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAR	A	301	-	-	0/4/16/16	0/3/3/3
4	X8W	B	301	-	-	0/9/9/9	0/2/2/2
5	GOL	B	302	-	-	0/4/4/4	0/0/0/0
2	NAR	C	301	-	-	0/4/16/16	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAR	D	301	-	-	0/4/16/16	0/3/3/3
2	NAR	D	302	-	-	0/4/16/16	0/3/3/3
4	X8W	D	303[A]	-	-	0/2/2/9	0/2/2/2
4	X8W	D	303[B]	-	-	0/2/2/9	0/2/2/2
2	NAR	E	301	-	-	0/4/16/16	0/3/3/3
2	NAR	F	301	-	-	0/4/16/16	0/3/3/3
2	NAR	F	302[A]	-	-	0/0/0/16	0/1/1/3
2	NAR	F	302[B]	-	-	0/0/0/16	0/1/1/3
5	GOL	F	303	-	-	0/4/4/4	0/0/0/0

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	302	NAR	C14-C13	-10.63	1.17	1.38
4	D	303[A]	X8W	CAT-CAN	-3.83	1.37	1.46
4	B	301	X8W	CAQ-CAG	-3.09	1.38	1.47
4	B	301	X8W	CAT-CAN	-3.00	1.42	1.49
4	D	303[B]	X8W	CAT-CAS	-2.51	1.38	1.41

The worst 5 of 84 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	302	NAR	C15-C10-C11	-15.42	98.65	118.31
2	D	302	NAR	O1-C6-C5	-6.79	113.80	121.55
2	D	302	NAR	C14-C13-C12	-6.56	110.45	119.79
2	F	301	NAR	O1-C6-C5	-6.34	114.32	121.55
2	D	301	NAR	C6-C5-C7	-6.09	114.79	120.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

10 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	NAR	3	0
4	B	301	X8W	5	0
5	B	302	GOL	2	0
2	C	301	NAR	1	0
2	D	301	NAR	4	0
2	D	302	NAR	1	0
4	D	303[A]	X8W	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	303[B]	X8W	5	0
2	E	301	NAR	1	0
2	F	301	NAR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/283 (99%)	-0.30	16 (5%) 27 29	18, 24, 46, 70	0
1	B	259/283 (91%)	-0.76	3 (1%) 81 81	17, 21, 32, 77	0
1	C	282/283 (99%)	-0.34	10 (3%) 48 49	17, 22, 40, 64	0
1	D	259/283 (91%)	-0.75	0 100 100	18, 20, 29, 59	0
1	E	282/283 (99%)	-0.18	20 (7%) 19 20	18, 22, 46, 69	0
1	F	259/283 (91%)	-0.68	1 (0%) 93 93	18, 20, 33, 62	0
All	All	1623/1698 (95%)	-0.49	50 (3%) 52 53	17, 22, 41, 77	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	130	ALA	5.9
1	E	107	ALA	5.5
1	E	130	ALA	5.2
1	E	115	ASP	4.8
1	F	106	ALA	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAR	D	301	20/20	0.74	0.23	8.34	35,49,61,62	11
2	NAR	F	302[B]	10/20	0.86	0.20	6.54	44,54,59,74	9
5	GOL	B	302	6/6	0.96	0.17	5.63	26,27,27,27	6
2	NAR	F	302[A]	10/20	0.86	0.20	5.52	47,57,59,74	9
4	X8W	D	303[A]	18/20	0.92	0.16	4.52	36,62,67,70	18
5	GOL	F	303	6/6	0.90	0.14	2.81	34,36,39,44	0
4	X8W	B	301	20/20	0.80	0.20	2.54	49,58,69,78	8
2	NAR	F	301	20/20	0.94	0.11	2.00	28,30,32,32	20
4	X8W	D	303[B]	18/20	0.92	0.16	1.60	37,39,76,77	18
2	NAR	C	301	20/20	0.93	0.14	1.37	26,30,32,32	20
2	NAR	A	301	20/20	0.92	0.13	1.27	26,30,33,33	20
2	NAR	E	301	20/20	0.93	0.14	1.22	27,32,36,36	20
2	NAR	D	302	20/20	0.95	0.10	1.07	26,28,35,44	20
3	CL	E	403	1/1	0.96	0.03	-	63,63,63,63	1
3	CL	D	403	1/1	0.97	0.05	-	49,49,49,49	0
3	CL	B	402	1/1	0.89	0.09	-	54,54,54,54	0
3	CL	E	404	1/1	0.74	0.15	-	60,60,60,60	0
3	CL	D	402	1/1	0.95	0.05	-	56,56,56,56	0
3	CL	F	402	1/1	0.97	0.05	-	51,51,51,51	0
3	CL	E	401	1/1	0.97	0.07	-	50,50,50,50	0
3	CL	D	401	1/1	0.97	0.04	-	48,48,48,48	0
3	CL	C	402	1/1	0.92	0.20	-	60,60,60,60	0
3	CL	A	402	1/1	0.95	0.10	-	53,53,53,53	0
3	CL	C	403	1/1	0.89	0.12	-	54,54,54,54	0
3	CL	F	401	1/1	0.90	0.09	-	48,48,48,48	0
3	CL	F	403	1/1	0.98	0.08	-	45,45,45,45	0
3	CL	C	401	1/1	0.92	0.08	-	52,52,52,52	0
3	CL	E	402	1/1	0.93	0.08	-	59,59,59,59	0
3	CL	B	401	1/1	0.97	0.09	-	53,53,53,53	0
3	CL	A	401	1/1	0.94	0.07	-	59,59,59,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.