



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 11, 2016 – 05:12 PM EDT

PDB ID : 5D13
Title : Third PDZ domain (PDZ3) of PSD-95 complexed with CFMOC-KKETEV peptide
Authors : De, S.; Spaller, M.R.; Olson, R.
Deposited on : 2015-08-03
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	<div> <div>3%</div> <div>84%</div> <div>16%</div> </div>
1	B	119	<div> <div>%</div> <div>82%</div> <div>16%</div> </div>
1	C	119	<div> <div>%</div> <div>82%</div> <div>15%</div> </div>
1	D	119	<div> <div>19%</div> <div>72%</div> <div>12%</div> <div>16%</div> </div>
2	E	7	<div> <div>100%</div> </div>
2	F	7	<div> <div>14%</div> <div>86%</div> <div>14%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	7	 86% 14%
2	H	7	 100%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5887 atoms, of which 2739 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Disks large homolog 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	H	N	O	0	0	0
			1357	440	644	134	139			
1	B	100	Total	C	H	N	O	0	0	0
			1447	459	710	136	142			
1	C	101	Total	C	H	N	O	0	0	0
			1398	450	666	137	145			
1	D	100	Total	C	H	N	O	0	0	0
			1159	399	517	117	126			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	297	GLY	-	expression tag	UNP P31016
A	298	SER	-	expression tag	UNP P31016
A	299	PRO	-	expression tag	UNP P31016
A	300	GLU	-	expression tag	UNP P31016
A	301	PHE	-	expression tag	UNP P31016
A	403	ASN	-	expression tag	UNP P31016
A	404	SER	-	expression tag	UNP P31016
A	405	ARG	-	expression tag	UNP P31016
A	406	VAL	-	expression tag	UNP P31016
A	407	ASP	-	expression tag	UNP P31016
A	408	SER	-	expression tag	UNP P31016
A	409	SER	-	expression tag	UNP P31016
A	410	GLY	-	expression tag	UNP P31016
A	411	ARG	-	expression tag	UNP P31016
A	412	ILE	-	expression tag	UNP P31016
A	413	VAL	-	expression tag	UNP P31016
A	414	THR	-	expression tag	UNP P31016
A	415	ASP	-	expression tag	UNP P31016
B	297	GLY	-	expression tag	UNP P31016
B	298	SER	-	expression tag	UNP P31016
B	299	PRO	-	expression tag	UNP P31016

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	300	GLU	-	expression tag	UNP P31016
B	301	PHE	-	expression tag	UNP P31016
B	403	ASN	-	expression tag	UNP P31016
B	404	SER	-	expression tag	UNP P31016
B	405	ARG	-	expression tag	UNP P31016
B	406	VAL	-	expression tag	UNP P31016
B	407	ASP	-	expression tag	UNP P31016
B	408	SER	-	expression tag	UNP P31016
B	409	SER	-	expression tag	UNP P31016
B	410	GLY	-	expression tag	UNP P31016
B	411	ARG	-	expression tag	UNP P31016
B	412	ILE	-	expression tag	UNP P31016
B	413	VAL	-	expression tag	UNP P31016
B	414	THR	-	expression tag	UNP P31016
B	415	ASP	-	expression tag	UNP P31016
C	297	GLY	-	expression tag	UNP P31016
C	298	SER	-	expression tag	UNP P31016
C	299	PRO	-	expression tag	UNP P31016
C	300	GLU	-	expression tag	UNP P31016
C	301	PHE	-	expression tag	UNP P31016
C	403	ASN	-	expression tag	UNP P31016
C	404	SER	-	expression tag	UNP P31016
C	405	ARG	-	expression tag	UNP P31016
C	406	VAL	-	expression tag	UNP P31016
C	407	ASP	-	expression tag	UNP P31016
C	408	SER	-	expression tag	UNP P31016
C	409	SER	-	expression tag	UNP P31016
C	410	GLY	-	expression tag	UNP P31016
C	411	ARG	-	expression tag	UNP P31016
C	412	ILE	-	expression tag	UNP P31016
C	413	VAL	-	expression tag	UNP P31016
C	414	THR	-	expression tag	UNP P31016
C	415	ASP	-	expression tag	UNP P31016
D	297	GLY	-	expression tag	UNP P31016
D	298	SER	-	expression tag	UNP P31016
D	299	PRO	-	expression tag	UNP P31016
D	300	GLU	-	expression tag	UNP P31016
D	301	PHE	-	expression tag	UNP P31016
D	403	ASN	-	expression tag	UNP P31016
D	404	SER	-	expression tag	UNP P31016
D	405	ARG	-	expression tag	UNP P31016
D	406	VAL	-	expression tag	UNP P31016

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	407	ASP	-	expression tag	UNP P31016
D	408	SER	-	expression tag	UNP P31016
D	409	SER	-	expression tag	UNP P31016
D	410	GLY	-	expression tag	UNP P31016
D	411	ARG	-	expression tag	UNP P31016
D	412	ILE	-	expression tag	UNP P31016
D	413	VAL	-	expression tag	UNP P31016
D	414	THR	-	expression tag	UNP P31016
D	415	ASP	-	expression tag	UNP P31016

- Molecule 2 is a protein called CFMOC-KKETEV peptide.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	7	Total	C	H	N	O	0	0	0
			120	44	56	7	13			
2	E	7	Total	C	H	N	O	0	0	0
			105	41	45	6	13			
2	H	7	Total	C	H	N	O	0	0	0
			105	41	45	6	13			
2	F	7	Total	C	H	N	O	0	0	0
			97	39	41	6	11			

- Molecule 3 is 3-(9H-fluoren-9-yl)propanal (three-letter code: 56J) (formula: C₁₆H₁₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	H	O	0	0
			30	16	13	1		

- Molecule 4 is LYSINE (three-letter code: LYS) (formula: $C_6H_{15}N_2O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	H	N	O	0	0
			11	6	2	2	1		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	6	Total	O	0	0
			6	6		
5	B	24	Total	O	0	0
			24	24		
5	C	19	Total	O	0	0
			19	19		
5	D	2	Total	O	0	0
			2	2		
5	G	5	Total	O	0	0
			5	5		
5	E	2	Total	O	0	0
			2	2		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Disks large homolog 4





- Molecule 1: Disks large homolog 4




- Molecule 1: Disks large homolog 4



- Molecule 1: Disks large homolog 4



- Molecule 2: CFMOC-KKETEV peptide



- Molecule 2: CFMOC-KKETEV peptide

Chain E:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: CFMOC-KKETEV peptide

Chain H:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: CFMOC-KKETEV peptide

Chain F:  14% 86% 14%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	84.94Å 84.94Å 210.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	36.50 – 2.15 36.50 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.50-2.15) 100.0 (36.50-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.16Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.210 , 0.257 0.210 , 0.256	Depositor DCC
R_{free} test set	1530 reflections (4.99%)	DCC
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.728	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 68.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/3*k+1/3*l 0.012 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+1/3*l,-4/3*h+4/3*k+1/3*l 0.000 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k+1/3*l 0.000 for -h,2/3*h+1/3*k+1/3*l,4/3*h+8/3*k-1/3*l 0.000 for 1/3*h+2/3*k-1/3*l,-k,-8/3*h-4/3*k-1/3*l 0.011 for -1/3*h-2/3*k+1/3*l,-2/3*h-1/3*k-1/3*l,4/3*h-4/3*k-1/3*l 0.039 for -h-k,k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5887	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.21 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.5769e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 56J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/724	0.59	0/981
1	B	0.60	0/748	0.69	0/1011
1	C	0.46	0/743	0.63	0/1004
1	D	0.31	0/652	0.54	0/891
2	E	0.44	0/42	0.58	0/55
2	F	0.32	0/38	0.44	0/50
2	G	0.69	0/46	0.76	0/59
2	H	0.29	0/42	0.56	0/55
All	All	0.47	0/3035	0.62	0/4106

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	713	644	648	0	0
1	B	737	710	711	2	0
1	C	732	666	674	2	0
1	D	642	517	529	12	0
2	E	60	45	31	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	56	41	27	2	0
2	G	64	56	42	1	0
2	H	60	45	31	0	0
3	C	17	13	0	0	0
4	C	9	2	12	0	0
5	A	6	0	0	0	0
5	B	24	0	0	1	0
5	C	19	0	0	1	0
5	D	2	0	0	2	0
5	E	2	0	0	0	0
5	G	5	0	0	1	0
All	All	3148	2739	2705	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:380:LYS:O	5:D:501:HOH:O	1.94	0.86
1:D:318:ARG:NH1	1:D:383:GLY:O	2.35	0.59
2:G:424:GLU:O	5:G:501:HOH:O	2.17	0.57
1:D:374:GLN:O	1:D:378:ALA:N	2.30	0.52
1:D:371:SER:OG	1:D:374:GLN:HG2	2.12	0.50

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	98/119 (82%)	98 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	98/119 (82%)	97 (99%)	1 (1%)	0	100	100
1	C	99/119 (83%)	98 (99%)	1 (1%)	0	100	100
1	D	98/119 (82%)	98 (100%)	0	0	100	100
2	E	4/7 (57%)	4 (100%)	0	0	100	100
2	F	4/7 (57%)	4 (100%)	0	0	100	100
2	G	4/7 (57%)	4 (100%)	0	0	100	100
2	H	4/7 (57%)	4 (100%)	0	0	100	100
All	All	409/504 (81%)	407 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/94 (68%)	64 (100%)	0	100	100
1	B	72/94 (77%)	72 (100%)	0	100	100
1	C	68/94 (72%)	68 (100%)	0	100	100
1	D	47/94 (50%)	47 (100%)	0	100	100
2	E	4/6 (67%)	4 (100%)	0	100	100
2	F	3/6 (50%)	3 (100%)	0	100	100
2	G	5/6 (83%)	5 (100%)	0	100	100
2	H	4/6 (67%)	4 (100%)	0	100	100
All	All	267/400 (67%)	267 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	56J	C	501	4	19,19,19	0.88	0	25,26,26	1.04	2 (8%)
4	LYS	C	502	3	6,8,9	0.49	0	6,8,10	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	56J	C	501	4	-	0/3/16/16	0/3/3/3
4	LYS	C	502	3	-	0/5/7/9	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	C	501	56J	C03-C04-C05	-2.54	116.74	120.30
3	C	501	56J	C12-C13-C08	-2.10	117.35	120.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/119 (84%)	0.27	3 (3%) 54 64	49, 70, 103, 145	0
1	B	100/119 (84%)	0.31	1 (1%) 84 88	39, 57, 87, 106	0
1	C	101/119 (84%)	0.37	1 (0%) 84 88	42, 69, 99, 110	0
1	D	100/119 (84%)	0.99	23 (23%) 1 2	68, 105, 136, 154	0
2	E	6/7 (85%)	0.06	0 100 100	64, 72, 84, 101	0
2	F	6/7 (85%)	0.38	1 (16%) 2 4	86, 99, 108, 134	0
2	G	6/7 (85%)	0.09	0 100 100	47, 55, 74, 88	0
2	H	6/7 (85%)	0.28	0 100 100	70, 84, 104, 119	0
All	All	425/504 (84%)	0.47	29 (6%) 20 28	39, 74, 126, 154	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	404	SER	5.9
1	D	390	ALA	5.4
1	A	402	ALA	3.8
1	B	402	ALA	3.8
1	D	308	PRO	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	LYS	C	502	9/10	0.89	0.13	0.85	54,61,79,94	0
3	56J	C	501	17/17	0.94	0.13	-0.13	36,55,66,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.