



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:04 AM GMT

PDB ID : 3D4V  
Title : Crystal Structure of an AlkA Host/Guest Complex N7MethylGuanine:Cytosine Base Pair  
Authors : Lee, S.; Bowman, B.R.; Wang, S.; Verdine, G.L.  
Deposited on : 2008-05-15  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	<div> <div></div> <div>62% 35% .</div> </div>
1	B	282	<div> <div></div> <div>56% 40% .</div> </div>
1	C	282	<div> <div></div> <div>52% 44% .</div> </div>
1	D	282	<div> <div></div> <div>48% 47% 5%</div> </div>
2	E	12	<div> <div>8%</div> <div>17% 83%</div> </div>
2	G	12	<div> <div>8%</div> <div>8% 83% 8%</div> </div>
3	F	12	<div> <div></div> <div>33% 67%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain	
			8%	83%
3	H	12		

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-3-methyladenine glycosylase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2215	1427	387	389	12			
1	B	282	Total	C	N	O	S	0	0	0
			2215	1427	387	389	12			
1	C	281	Total	C	N	O	S	0	0	0
			2209	1424	386	387	12			
1	D	281	Total	C	N	O	S	0	0	0
			2209	1424	386	387	12			

- Molecule 2 is a DNA chain called 5'-D(\*DGP\*DAP\*DCP\*DAP\*DTP\*DGP\*DAP\*(FMG)P\*DTP\*DGP\*DCP\*DC)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	12	Total	C	F	N	O	P	0	0
			247	118	1	48	69	11		
2	G	12	Total	C	F	N	O	P	0	0
			247	118	1	48	69	11		

- Molecule 3 is a DNA chain called 5'-D(\*DGP\*DGP\*DCP\*DAP\*DCP\*DTP\*DCP\*DAP\*DT\*P\*DGP\*DTP\*DC)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	12	Total	C	N	O	P	0	0	0
			241	116	43	71	11			
3	H	12	Total	C	N	O	P	0	0	0
			241	116	43	71	11			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	7	Total 7	O 7	0	0
4	C	6	Total 6	O 6	0	0
4	D	4	Total 4	O 4	0	0
4	E	1	Total 1	O 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA-3-methyladenine glycosylase 2



#### • Molecule 1: DNA-3-methyladenine glycosylase 2




#### • Molecule 1: DNA-3-methyladenine glycosylase 2






- Molecule 1: DNA-3-methyladenine glycosylase 2

Chain D: 48% 47% 5%




- Molecule 2: 5'-D(\*DGP\*DAP\*DCP\*DAP\*DTP\*DGP\*DAP\*(FMG)P\*DTP\*DGP\*DCP\*DC)-3'

Chain E: 8% 17% 83%




- Molecule 2: 5'-D(\*DGP\*DAP\*DCP\*DAP\*DTP\*DGP\*DAP\*(FMG)P\*DTP\*DGP\*DCP\*DC)-3'

Chain G: 8% 83% 8%



- Molecule 3: 5'-D(\*DGP\*DGP\*DCP\*DAP\*DCP\*DTP\*DCP\*DAP\*DTP\*DGP\*DTP\*DC)-3'

Chain F: 33% 67%



- Molecule 3: 5'-D(\*DGP\*DGP\*DCP\*DAP\*DCP\*DTP\*DCP\*DAP\*DTP\*DGP\*DTP\*DC)-3'

Chain H: 8% 17% 83%

G14	G15	G16	A17	G18	T19	C20	A21	T22	G23	T24	G25
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.10Å 100.36Å 102.28Å 90.00° 94.45° 90.00°	Depositor
Resolution (Å)	50.00 – 2.90 31.79 – 2.90	Depositor EDS
% Data completeness (in resolution range)	91.5 (50.00-2.90) 91.3 (31.79-2.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.57 (at 2.90Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.208 , 0.275 0.207 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	54.4	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 59.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 32280 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9845	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2277	0.62	0/3104
1	B	0.41	0/2277	0.66	0/3104
1	C	0.40	0/2271	0.63	0/3097
1	D	0.40	0/2271	0.62	0/3097
2	E	0.55	0/249	0.73	0/380
2	G	0.68	1/249 (0.4%)	0.70	0/380
3	F	0.50	0/269	0.81	0/413
3	H	0.59	0/269	0.73	0/413
All	All	0.42	1/10132 (0.0%)	0.65	0/13988

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	7	DA	C3'-O3'	-5.52	1.36	1.44

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	0	2209	101	0
1	B	2215	0	2209	116	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2209	0	2204	120	0
1	D	2209	0	2204	150	0
2	E	247	0	138	10	0
2	G	247	0	138	19	0
3	F	241	0	137	13	0
3	H	241	0	137	16	0
4	A	3	0	0	0	0
4	B	7	0	0	0	0
4	C	6	0	0	0	0
4	D	4	0	0	0	0
4	E	1	0	0	0	0
All	All	9845	0	9376	528	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 528 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:21:DA:H2''	3:H:22:DT:H5'	1.20	1.12
1:C:178:ARG:HH11	1:C:178:ARG:HB2	1.17	1.05
1:B:77:LYS:HB3	1:B:111:VAL:HG23	1.40	1.02
1:A:124:GLN:HE21	1:A:178:ARG:HH11	1.05	1.00
3:H:15:DG:H2''	3:H:16:DC:H5'	1.47	0.95

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	280/282 (99%)	238 (85%)	35 (12%)	7 (2%)	<b>7</b> <b>27</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	280/282 (99%)	247 (88%)	29 (10%)	4 (1%)	14	44
1	C	279/282 (99%)	244 (88%)	30 (11%)	5 (2%)	11	37
1	D	279/282 (99%)	239 (86%)	32 (12%)	8 (3%)	6	23
All	All	1118/1128 (99%)	968 (87%)	126 (11%)	24 (2%)	9	32

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	257	ALA
1	A	275	GLU
1	B	175	PRO
1	B	275	GLU
1	C	250	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/222 (100%)	213 (96%)	9 (4%)	37	73
1	B	222/222 (100%)	210 (95%)	12 (5%)	27	62
1	C	222/222 (100%)	208 (94%)	14 (6%)	22	54
1	D	222/222 (100%)	204 (92%)	18 (8%)	15	39
All	All	888/888 (100%)	835 (94%)	53 (6%)	24	57

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	105	LEU
1	C	167	GLN
1	D	160	ARG
1	C	111	VAL
1	C	147	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	159	GLN
1	B	221	ASN
1	D	140	GLN
1	B	167	GLN
1	B	184	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FMG	E	8	3,2	18,26,27	1.59	3 (16%)	20,39,42	2.63	6 (30%)
2	FMG	G	8	3,2	18,26,27	1.72	5 (27%)	20,39,42	2.78	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMG	E	8	3,2	-	0/3/25/26	0/3/3/3
2	FMG	G	8	3,2	-	0/3/25/26	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	8	FMG	C8-N7	-2.48	1.28	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	8	FMG	F1'-C2'	-2.22	1.34	1.40
2	G	8	FMG	C8-N9	2.15	1.37	1.33
2	E	8	FMG	C2-N1	2.52	1.39	1.35
2	E	8	FMG	C8-N9	2.79	1.38	1.33

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	8	FMG	C5-C6-N1	-8.15	112.44	123.59
2	G	8	FMG	C5-C6-N1	-8.05	112.59	123.59
2	G	8	FMG	C4'-O4'-C1'	-4.42	104.86	109.72
2	E	8	FMG	N3-C2-N1	-2.66	123.40	127.44
2	G	8	FMG	N3-C2-N1	-2.55	123.56	127.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	8	FMG	2	0
2	G	8	FMG	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	282/282 (100%)	-0.18	4 (1%) 78 76	23, 56, 105, 141	0
1	B	282/282 (100%)	-0.40	1 (0%) 93 92	23, 45, 86, 141	0
1	C	281/282 (99%)	-0.34	2 (0%) 89 88	17, 49, 100, 138	0
1	D	281/282 (99%)	-0.28	1 (0%) 93 92	27, 54, 99, 180	0
2	E	11/12 (91%)	0.43	1 (9%) 11 7	64, 83, 130, 139	0
2	G	11/12 (91%)	0.28	1 (9%) 11 7	69, 92, 111, 123	0
3	F	12/12 (100%)	0.52	0 100 100	74, 123, 131, 133	0
3	H	12/12 (100%)	0.25	1 (8%) 14 9	49, 82, 116, 117	0
All	All	1172/1176 (99%)	-0.27	11 (0%) 85 84	17, 52, 105, 180	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	282	ALA	4.5
2	E	12	DC	3.0
1	A	282	ALA	2.9
1	A	110	CYS	2.9
1	C	281	GLU	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMG	G	8	24/25	0.84	0.27	-	82,89,103,107	0
2	FMG	E	8	24/25	0.90	0.26	-	82,89,103,107	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.