



Full wwPDB X-ray Structure Validation Report ⓘ

May 31, 2016 – 06:05 PM EDT

PDB ID : 5D41
Title : EGFR kinase domain in complex with mutant selective allosteric inhibitor
Authors : Yun, C.-H.; Park, E.; Eck, M.J.
Deposited on : 2015-08-07
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>5%</div> <div> <div></div> <div>79%</div> <div>12%</div> <div>9%</div> </div> </div>
1	B	331	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>13%</div> <div>9%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5313 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	0	1	0
			2408	1543	408	437	20			
1	B	302	Total	C	N	O	S	0	1	0
			2418	1549	408	442	19			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	692	GLY	-	expression tag	UNP P00533
A	693	SER	THR	expression tag	UNP P00533
A	694	THR	PRO	expression tag	UNP P00533
A	790	MET	THR	engineered mutation	UNP P00533
A	948	ARG	VAL	engineered mutation	UNP P00533
B	692	GLY	-	expression tag	UNP P00533
B	693	SER	THR	expression tag	UNP P00533
B	694	THR	PRO	expression tag	UNP P00533
B	790	MET	THR	engineered mutation	UNP P00533
B	948	ARG	VAL	engineered mutation	UNP P00533

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
3	B	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 4 is (2R)-2-(1-oxo-1,3-dihydro-2H-isoinol-2-yl)-2-phenyl-N-(1,3-thiazol-2-yl)acetamide (three-letter code: 57N) (formula: C₁₉H₁₅N₃O₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			25	19	3	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	215	Total 215	O 215	0	0
5	B	183	Total 183	O 183	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Epidermal growth factor receptor



• Molecule 1: Epidermal growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.13Å 72.50Å 76.00Å 90.00° 113.24° 90.00°	Depositor
Resolution (Å)	42.24 – 2.31 43.69 – 2.31	Depositor EDS
% Data completeness (in resolution range)	97.7 (42.24-2.31) 97.6 (43.69-2.31)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.32Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.174 , 0.207 0.174 , 0.206	Depositor DCC
R_{free} test set	1681 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	29.0	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5313	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.99 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.8822e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP, 57N

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2464	0.55	0/3336
1	B	0.44	0/2472	0.54	0/3345
All	All	0.42	0/4936	0.54	0/6681

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2408	0	2430	31	0
1	B	2418	0	2435	32	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	12	0	0
3	B	31	0	12	2	0
4	A	25	0	0	1	0
5	A	215	0	0	10	2
5	B	183	0	0	14	2
All	All	5313	0	4889	62	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:806:LYS:HG3	1:A:807:ASP:OD1	1.59	1.01
1:A:986:ARG:NH1	5:A:1201:HOH:O	1.95	0.99
3:B:2001:ANP:O1B	5:B:2101:HOH:O	1.85	0.92
1:B:766:MET:SD	5:B:2278:HOH:O	2.35	0.83
1:A:962:ARG:NH2	5:A:1203:HOH:O	2.14	0.80
1:B:733:PRO:O	5:B:2102:HOH:O	2.00	0.79
1:A:922:GLU:OE1	5:A:1202:HOH:O	2.02	0.78
1:B:715:ILE:HD11	1:B:730:LEU:HG	1.67	0.77
1:B:985:GLU:OE2	5:B:2103:HOH:O	2.06	0.72
1:A:876:VAL:N	5:A:1207:HOH:O	2.23	0.71
1:B:759:ILE:HD13	1:B:786:VAL:HG21	1.72	0.70
1:B:1006:ASP:OD2	5:B:2104:HOH:O	2.11	0.69
1:A:962:ARG:NH1	5:A:1208:HOH:O	2.26	0.68
1:B:1004:GLU:OE1	5:B:2106:HOH:O	2.14	0.66
1:A:701:GLN:OE1	5:A:1204:HOH:O	2.14	0.66
1:B:749:GLU:OE2	5:B:2107:HOH:O	2.14	0.65
1:B:999:ARG:NH2	1:B:1005:GLU:O	2.29	0.63
1:A:806:LYS:CG	1:A:807:ASP:OD1	2.45	0.60
1:B:719:GLY:HA3	3:B:2001:ANP:H4'	1.83	0.59
1:B:804:GLU:HB3	5:B:2155:HOH:O	2.03	0.58
1:A:783:THR:O	5:A:1205:HOH:O	2.18	0.57
1:A:715:ILE:HD11	1:A:728:LYS:HE2	1.87	0.56
1:B:949:LYS:HA	1:B:952:MET:HE1	1.87	0.56
1:A:905:TRP:HD1	1:A:947:MET:HE1	1.69	0.55
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.42	0.54
1:B:882:ALA:HA	1:B:898:TRP:CD2	2.43	0.54
1:A:1012:ASP:OD1	1:A:1013:ALA:N	2.41	0.54
1:B:949:LYS:HG2	1:B:959:PRO:HD3	1.92	0.52
1:B:832:ARG:NH2	5:B:2123:HOH:O	2.42	0.52
1:B:986:ARG:NH2	5:B:2116:HOH:O	2.36	0.52
1:B:760:LEU:HD13	1:B:782:LEU:HD11	1.92	0.52
1:A:835:HIS:CD2	1:A:856:PHE:HB3	2.46	0.50
1:A:1004:GLU:OE1	1:B:1005:GLU:HG3	2.11	0.50
1:B:982:GLN:NE2	5:B:2125:HOH:O	2.47	0.47
1:B:905:TRP:HB2	1:B:947:MET:HE1	1.97	0.47
1:B:769:VAL:HG21	1:B:856:PHE:CZ	2.50	0.47
1:B:809:ILE:O	5:B:2108:HOH:O	2.20	0.47
1:A:804:GLU:OE1	1:B:737:LYS:HD2	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:707:LEU:HD13	1:B:789:ILE:HD13	1.97	0.46
1:B:724:GLY:HA2	1:B:748:ARG:HG2	1.97	0.46
1:B:943:VAL:O	1:B:947:MET:HG3	2.17	0.45
1:A:802:VAL:HA	1:A:809:ILE:HD11	1.99	0.45
1:A:824:GLY:HA3	1:A:853:ILE:HD12	1.97	0.45
1:A:707:LEU:HD12	1:A:789:ILE:HD13	1.98	0.44
1:B:985:GLU:CD	5:B:2103:HOH:O	2.51	0.44
1:A:1001:LEU:HA	1:A:1001:LEU:HD12	1.68	0.43
1:A:698:ALA:HA	1:A:699:PRO:HD3	1.87	0.43
1:A:1007:MET:HG2	1:A:1008:ASP:H	1.83	0.43
1:A:1007:MET:C	5:A:1209:HOH:O	2.56	0.43
1:A:806:LYS:HB2	1:A:910:PHE:HB3	2.00	0.43
1:A:835:HIS:O	1:A:836:ARG:HB2	2.19	0.43
1:B:800:ASP:O	1:B:804:GLU:HG3	2.19	0.42
1:B:709:GLU:CD	1:B:783:THR:HG21	2.39	0.42
1:A:740:ILE:HG21	1:B:1002:MET:HE2	2.01	0.41
1:A:808:ASN:HA	1:A:987:MET:HE2	2.02	0.41
4:A:1103:57N:S18	4:A:1103:57N:O13	2.79	0.41
1:B:991:SER:HB2	5:B:2167:HOH:O	2.21	0.41
1:B:793:MET:HA	1:B:794:PRO:HD2	1.94	0.40
1:A:811:SER:OG	1:A:975:PRO:HB2	2.21	0.40
1:A:999:ARG:HA	1:A:1003:ASP:HB3	2.03	0.40
1:A:698:ALA:N	5:A:1233:HOH:O	2.54	0.40
1:A:962:ARG:NH1	5:A:1228:HOH:O	2.52	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1378:HOH:O	5:B:2190:HOH:O[4_455]	1.85	0.35
5:A:1355:HOH:O	5:B:2203:HOH:O[4_445]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/331 (90%)	288 (97%)	9 (3%)	0	100	100
1	B	299/331 (90%)	290 (97%)	9 (3%)	0	100	100
All	All	596/662 (90%)	578 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/290 (92%)	266 (100%)	0	100	100
1	B	265/290 (91%)	264 (100%)	1 (0%)	93	97
All	All	531/580 (92%)	530 (100%)	1 (0%)	95	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	998	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ANP	A	1102	2	29,33,33	4.82	16 (55%)	26,52,52	3.00	6 (23%)
4	57N	A	1103	-	26,28,28	5.47	21 (80%)	29,39,39	3.82	6 (20%)
3	ANP	B	2001	2	29,33,33	5.11	15 (51%)	26,52,52	3.07	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	1102	2	-	0/13/38/38	0/3/3/3
4	57N	A	1103	-	-	0/14/28/28	0/4/4/4
3	ANP	B	2001	2	-	0/13/38/38	0/3/3/3

All (52) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2001	ANP	C2'-C1'	-14.05	1.31	1.53
3	A	1102	ANP	C2'-C1'	-12.95	1.33	1.53
3	A	1102	ANP	O3'-C3'	-5.12	1.30	1.43
3	B	2001	ANP	O3'-C3'	-5.08	1.31	1.43
3	B	2001	ANP	O4'-C4'	-4.86	1.33	1.45
3	A	1102	ANP	O4'-C4'	-4.56	1.34	1.45
3	B	2001	ANP	C5-C4	-3.82	1.31	1.40
4	A	1103	57N	C17-S18	-3.78	1.51	1.71
3	B	2001	ANP	PG-O3G	-3.76	1.46	1.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1102	ANP	C5-C4	-3.49	1.32	1.40
3	B	2001	ANP	O5'-C5'	-3.45	1.31	1.44
3	A	1102	ANP	O5'-C5'	-3.41	1.31	1.44
3	A	1102	ANP	PG-O3G	-3.05	1.48	1.56
4	A	1103	57N	C09-N25	-2.65	1.43	1.46
4	A	1103	57N	O13-C11	-2.60	1.18	1.23
4	A	1103	57N	C09-C11	-2.10	1.51	1.54
4	A	1103	57N	C03-C07	2.17	1.52	1.48
3	B	2001	ANP	PA-O5'	2.33	1.69	1.59
3	A	1102	ANP	PA-O5'	2.33	1.69	1.59
3	A	1102	ANP	C6-N6	2.46	1.44	1.34
3	B	2001	ANP	C6-N6	2.47	1.44	1.34
3	A	1102	ANP	PB-N3B	2.59	1.70	1.63
3	A	1102	ANP	O2'-C2'	2.65	1.49	1.43
3	B	2001	ANP	O2'-C2'	2.67	1.49	1.43
3	A	1102	ANP	C4-N3	2.73	1.39	1.35
3	B	2001	ANP	PB-N3B	2.83	1.71	1.63
4	A	1103	57N	C16-N15	4.01	1.56	1.37
4	A	1103	57N	C14-N12	4.39	1.43	1.36
4	A	1103	57N	C11-N12	4.63	1.44	1.35
3	B	2001	ANP	C2-N3	4.75	1.40	1.32
4	A	1103	57N	C03-C04	4.82	1.45	1.39
4	A	1103	57N	C01-C06	5.16	1.50	1.38
3	B	2001	ANP	PG-N3B	5.17	1.77	1.63
3	A	1102	ANP	PG-N3B	5.33	1.78	1.63
3	A	1102	ANP	C2-N3	5.44	1.41	1.32
4	A	1103	57N	C21-C20	5.68	1.52	1.38
4	A	1103	57N	C22-C21	6.03	1.53	1.38
4	A	1103	57N	C06-C05	6.22	1.51	1.38
4	A	1103	57N	C01-C02	6.76	1.52	1.38
4	A	1103	57N	C20-C19	6.84	1.52	1.38
4	A	1103	57N	C22-C23	6.96	1.52	1.38
3	A	1102	ANP	PB-O1B	7.11	1.53	1.46
4	A	1103	57N	C19-C10	8.26	1.52	1.39
4	A	1103	57N	C23-C10	8.54	1.53	1.39
4	A	1103	57N	C05-C04	8.68	1.54	1.39
4	A	1103	57N	C07-N25	8.83	1.45	1.36
4	A	1103	57N	C02-C03	9.12	1.54	1.39
3	B	2001	ANP	PB-O1B	9.25	1.56	1.46
3	B	2001	ANP	PB-O3A	10.44	1.72	1.59
3	A	1102	ANP	PB-O3A	10.54	1.72	1.59
3	A	1102	ANP	O4'-C1'	12.83	1.59	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2001	ANP	O4'-C1'	13.88	1.61	1.41

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2001	ANP	N3-C2-N1	-12.13	119.34	128.87
3	A	1102	ANP	N3-C2-N1	-11.61	119.75	128.87
4	A	1103	57N	C24-N25-C07	-11.34	109.23	113.03
3	B	2001	ANP	N6-C6-N1	-7.43	106.05	118.52
3	A	1102	ANP	N6-C6-N1	-7.08	106.64	118.52
3	A	1102	ANP	C4'-O4'-C1'	-3.62	105.80	109.64
3	B	2001	ANP	O4'-C1'-N9	-3.48	101.52	108.11
3	B	2001	ANP	PA-O3A-PB	-2.69	122.96	132.71
4	A	1103	57N	C24-C04-C03	-2.68	107.98	109.71
3	A	1102	ANP	PA-O3A-PB	-2.68	123.00	132.71
4	A	1103	57N	O08-C07-N25	-2.23	123.61	125.17
3	A	1102	ANP	C1'-N9-C4	-2.12	124.44	126.81
3	A	1102	ANP	C5'-C4'-C3'	-2.07	107.21	115.20
4	A	1103	57N	C23-C10-C19	2.42	121.36	118.30
4	A	1103	57N	C09-C11-N12	4.52	119.25	114.50
4	A	1103	57N	C04-C24-N25	15.23	107.26	102.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1103	57N	1	0
3	B	2001	ANP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	300/331 (90%)	0.00	17 (5%)	27 36	17, 29, 61, 76	0
1	B	302/331 (91%)	0.07	13 (4%)	39 48	18, 31, 62, 99	0
All	All	602/662 (90%)	0.04	30 (4%)	32 41	17, 30, 61, 99	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	861	LEU	7.9
1	A	698	ALA	6.6
1	B	750	ALA	4.8
1	B	752	SER	4.6
1	B	859	ALA	4.0
1	B	753	PRO	3.7
1	B	751	THR	3.6
1	A	1013	ALA	3.6
1	A	753	PRO	3.6
1	B	723	PHE	3.4
1	A	989	LEU	3.3
1	A	807	ASP	3.1
1	B	784	SER	2.9
1	A	752	SER	2.8
1	A	1007	MET	2.8
1	A	988	HIS	2.7
1	B	755	ALA	2.7
1	B	862	LEU	2.6
1	B	749	GLU	2.6
1	A	760	LEU	2.5
1	A	986	ARG	2.4
1	A	701	GLN	2.3
1	A	764	TYR	2.3
1	A	784	SER	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	754	LYS	2.2
1	B	875	LYS	2.2
1	A	699	PRO	2.2
1	A	1014	ASP	2.1
1	A	783	THR	2.1
1	A	1012	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	57N	A	1103	25/25	0.96	0.17	1.08	15,26,36,42	0
3	ANP	B	2001	31/31	0.98	0.12	-0.18	20,28,43,46	0
3	ANP	A	1102	31/31	0.98	0.13	-0.58	16,20,25,28	0
2	MG	A	1101	1/1	0.98	0.06	-	20,20,20,20	0
2	MG	B	2000	1/1	0.96	0.08	-	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.