



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:29 AM GMT

PDB ID : 2D5H  
Title : Crystal Structure of Recombinant Soybean Proglycinin A3B4 subunit, its Comparison with Mature Glycinin A3B4 subunit, Responsible for Hexamer Assembly  
Authors : Itoh, T.; Adachi, M.; Masuda, T.; Mikami, B.; Utsumi, S.  
Deposited on : 2005-11-01  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865




# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	493	<div> <div>4%</div> <div>32% 40% • 24%</div> </div>
1	B	493	<div> <div>3%</div> <div>33% 39% • 25%</div> </div>
1	C	493	<div> <div>6%</div> <div>26% 44% 5% 25%</div> </div>
1	D	493	<div> <div>%</div> <div>36% 38% • 24%</div> </div>
1	E	493	<div> <div>%</div> <div>40% 34% • 23%</div> </div>

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	493	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CO3	B	502	-	-	-	X



## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 17703 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called glycinin A3B4 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	374	Total	C	N	O	S	0	0	0
			2937	1844	523	562	8			
1	B	372	Total	C	N	O	S	0	0	0
			2920	1835	520	557	8			
1	C	370	Total	C	N	O	S	0	0	0
			2905	1827	517	553	8			
1	D	377	Total	C	N	O	S	0	0	0
			2967	1861	529	569	8			
1	E	380	Total	C	N	O	S	0	0	0
			2981	1872	530	571	8			
1	F	377	Total	C	N	O	S	0	0	0
			2963	1860	528	567	8			

- Molecule 2 is CARBONATE ION (three-letter code: CO3) (formula: CO<sub>3</sub>).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 1 3	0	0
2	B	1	Total C O 4 1 3	0	0
2	C	1	Total C O 4 1 3	0	0
2	D	1	Total C O 4 1 3	0	0
2	E	1	Total C O 4 1 3	0	0
2	F	1	Total C O 4 1 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Mg 1 1	0	0
3	E	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0
3	C	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0
3	F	1	Total Mg 1 1	0	0




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: glycinin A3B4 subunit



- Molecule 1: glycinin A3B4 subunit











• Molecule 1: glycinin A3B4 subunit





Chain F:





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.38Å 223.74Å 88.66Å 90.00° 119.85° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 45.36 – 2.78	Depositor EDS
% Data completeness (in resolution range)	85.1 (15.00-2.80) 83.1 (45.36-2.78)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.79 (at 2.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.248 , 0.325 0.231 , 0.306	Depositor DCC
$R_{free}$ test set	5961 reflections (10.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.4	Xtriage
Anisotropy	0.250	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 35.3	EDS
Estimated twinning fraction	0.047 for -h-l,k,h 0.047 for l,k,-h-l 0.034 for -h-l,-k,l 0.035 for l,-k,h 0.034 for h,-k,-h-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 63184 reflections	Xtriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	17703	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, CO3, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/2995	0.49	0/4068
1	B	0.27	0/2978	0.49	0/4045
1	C	0.27	0/2963	0.50	0/4025
1	D	0.26	0/3025	0.51	0/4108
1	E	0.28	0/3039	0.52	0/4127
1	F	0.27	0/3021	0.51	0/4103
All	All	0.27	0/18021	0.50	0/24476

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2937	0	2841	248	0
1	B	2920	0	2830	251	0
1	C	2905	0	2817	265	0
1	D	2967	0	2867	193	0
1	E	2981	0	2897	183	0
1	F	2963	0	2868	211	0
2	A	4	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	4	0	0	1	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
2	F	4	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
All	All	17703	0	17120	1233	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 1233 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:334:ASN:HD22	1:C:337:ARG:HG2	1.17	1.08
1:F:159:ASN:HD22	1:F:176:THR:HG22	1.20	1.06
1:F:115:ARG:HH22	1:F:331:LEU:HD12	1.20	1.04
1:E:247:ILE:HG12	1:E:248:SER:H	1.19	1.04
1:B:159:ASN:HD22	1:B:176:THR:HG22	1.29	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	365/493 (74%)	302 (83%)	54 (15%)	9 (2%)	<b>7</b> <b>24</b>

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	363/493 (74%)	297 (82%)	54 (15%)	12 (3%)	5	16
1	C	361/493 (73%)	291 (81%)	51 (14%)	19 (5%)	2	7
1	D	368/493 (75%)	307 (83%)	54 (15%)	7 (2%)	10	32
1	E	371/493 (75%)	311 (84%)	54 (15%)	6 (2%)	12	38
1	F	368/493 (75%)	311 (84%)	44 (12%)	13 (4%)	4	15
All	All	2196/2958 (74%)	1819 (83%)	311 (14%)	66 (3%)	5	18

5 of 66 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	426	GLN
1	B	426	GLN
1	C	245	SER
1	C	418	ARG
1	D	171	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/435 (74%)	310 (96%)	14 (4%)	35	70
1	B	322/435 (74%)	313 (97%)	9 (3%)	51	84
1	C	320/435 (74%)	305 (95%)	15 (5%)	32	67
1	D	327/435 (75%)	314 (96%)	13 (4%)	38	73
1	E	330/435 (76%)	318 (96%)	12 (4%)	42	76
1	F	327/435 (75%)	316 (97%)	11 (3%)	44	78
All	All	1950/2610 (75%)	1876 (96%)	74 (4%)	40	74

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	371	TYR

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type
1	D	121	ASP
1	F	200	GLU
1	C	380	TYR
1	C	436	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 134 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	178	GLN
1	D	12	ASN
1	F	334	ASN
1	C	332	HIS
1	C	409	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSD	A	406	1	3,7,8	0.61	0	3,8,10	1.35	0
1	CSD	B	406	1	3,7,8	0.63	0	3,8,10	1.53	0
1	CSD	C	406	1	3,7,8	0.55	0	3,8,10	1.29	0
1	CSD	D	406	1	3,7,8	0.61	0	3,8,10	1.49	0
1	CSD	E	406	1	3,7,8	0.55	0	3,8,10	1.37	0
1	CSD	F	406	1	3,7,8	0.60	0	3,8,10	1.56	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical



component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	406	1	-	0/2/6/8	0/0/0/0
1	CSD	B	406	1	-	0/2/6/8	0/0/0/0
1	CSD	C	406	1	-	0/2/6/8	0/0/0/0
1	CSD	D	406	1	-	0/2/6/8	0/0/0/0
1	CSD	E	406	1	-	0/2/6/8	0/0/0/0
1	CSD	F	406	1	-	0/2/6/8	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	406	CSD	O-C-CA	-2.04	120.18	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	406	CSD	2	0
1	C	406	CSD	2	0
1	E	406	CSD	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CO3	A	501	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	B	502	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	C	503	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	D	504	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	E	505	-	0,3,3	0.00	-	0,3,3	0.00	-
2	CO3	F	506	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CO3	A	501	-	-	0/0/0/0	0/0/0/0
2	CO3	B	502	-	-	0/0/0/0	0/0/0/0
2	CO3	C	503	-	-	0/0/0/0	0/0/0/0
2	CO3	D	504	-	-	0/0/0/0	0/0/0/0
2	CO3	E	505	-	-	0/0/0/0	0/0/0/0
2	CO3	F	506	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	CO3	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	373/493 (75%)	0.08	18 (4%) 34 23	13, 37, 54, 62	0
1	B	371/493 (75%)	0.03	13 (3%) 48 35	6, 37, 56, 63	0
1	C	369/493 (74%)	0.28	28 (7%) 17 9	7, 39, 54, 66	0
1	D	376/493 (76%)	-0.38	7 (1%) 70 59	6, 19, 46, 57	0
1	E	379/493 (76%)	-0.43	3 (0%) 87 81	6, 13, 43, 58	0
1	F	376/493 (76%)	-0.41	3 (0%) 87 81	6, 18, 45, 57	0
All	All	2244/2958 (75%)	-0.14	72 (3%) 51 39	6, 28, 52, 66	0

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	88	THR	6.2
1	B	437	GLN	4.9
1	C	89	PHE	4.6
1	C	435	GLY	4.4
1	C	226	LYS	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSD	B	406	8/9	0.94	0.14	-	39,44,63,73	0
1	CSD	A	406	8/9	0.91	0.16	-	30,41,46,53	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSD	D	406	8/9	0.96	0.10	-	5,13,23,38	0
1	CSD	C	406	8/9	0.88	0.16	-	31,38,58,62	0
1	CSD	F	406	8/9	0.95	0.17	-	5,8,30,57	0
1	CSD	E	406	8/9	0.96	0.14	-	5,7,45,53	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CO3	B	502	4/4	0.88	0.22	4.20	12,22,39,45	0
2	CO3	A	501	4/4	0.88	0.24	1.57	14,35,39,41	0
3	MG	E	605	1/1	0.99	0.16	0.94	5,5,5,5	0
2	CO3	E	505	4/4	0.89	0.17	0.93	5,5,10,17	0
2	CO3	F	506	4/4	0.87	0.19	0.92	5,5,18,21	0
3	MG	D	604	1/1	0.96	0.16	-0.07	17,17,17,17	0
3	MG	B	602	1/1	0.91	0.21	-0.81	41,41,41,41	0
2	CO3	D	504	4/4	0.97	0.14	-0.82	5,19,20,21	0
2	CO3	C	503	4/4	0.91	0.15	-1.24	36,40,41,42	0
3	MG	F	606	1/1	0.97	0.14	-1.46	8,8,8,8	0
3	MG	C	603	1/1	0.93	0.13	-2.35	21,21,21,21	0
3	MG	A	601	1/1	0.94	0.08	-4.99	17,17,17,17	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.