



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:23 PM GMT

PDB ID : 5D5Z
Title : Structure of Chaetomium thermophilum Skn7 coiled-coil domain, crystal form II
Authors : Neudegger, T.; Verghese, J.; Hayer-Hartl, M.; Bracher, A.; Hartl, F.U.
Deposited on : 2015-08-11
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	50	<div> <div>2%</div> <div>84%</div> <div>16%</div> </div>
1	B	50	<div> <div>2%</div> <div>86%</div> <div>14%</div> </div>
1	C	50	<div> <div>8%</div> <div>86%</div> <div>14%</div> </div>
1	D	50	<div> <div>12%</div> <div>90%</div> <div>10%</div> </div>
1	E	50	<div> <div>18%</div> <div>70%</div> <div>22%</div> <div>6%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative transcription factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	50	Total	C	N	O	S	0	0	0
			397	243	70	81	3			
1	B	50	Total	C	N	O	S	0	0	0
			393	241	69	80	3			
1	C	50	Total	C	N	O	S	0	0	0
			393	240	69	81	3			
1	D	50	Total	C	N	O	S	0	0	0
			397	243	70	81	3			
1	E	49	Total	C	N	O	S	0	0	0
			387	238	68	78	3			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	37	Total	O	0	0
			37	37		
2	B	21	Total	O	0	0
			21	21		
2	C	19	Total	O	0	0
			19	19		
2	D	30	Total	O	0	0
			30	30		
2	E	14	Total	O	0	0
			14	14		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Putative transcription factor





- Molecule 1: Putative transcription factor




- Molecule 1: Putative transcription factor



- Molecule 1: Putative transcription factor



- Molecule 1: Putative transcription factor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	47.00Å 61.03Å 75.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.70 29.50 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.70) 100.0 (29.50-1.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.70Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.209 , 0.285 0.213 , 0.292	Depositor DCC
R_{free} test set	1259 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.354	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	3 of 24650 reflections (0.012%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2088	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	1/397 (0.3%)	1.13	1/531 (0.2%)
1	B	0.79	0/393	0.86	0/526
1	C	0.87	0/393	0.98	2/527 (0.4%)
1	D	0.95	0/397	0.88	0/531
1	E	1.01	0/387	1.13	2/519 (0.4%)
All	All	0.96	1/1967 (0.1%)	1.00	5/2634 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	197	SER	CB-OG	-7.60	1.32	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	193	ARG	NE-CZ-NH2	-8.37	116.12	120.30
1	C	193	ARG	NE-CZ-NH1	7.57	124.09	120.30
1	A	200	MET	CG-SD-CE	-7.14	88.78	100.20
1	E	178	LEU	CB-CG-CD2	-5.19	102.17	111.00
1	E	188	LEU	CB-CG-CD1	5.05	119.58	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	397	0	409	4	0
1	B	393	0	403	6	0
1	C	393	0	398	5	0
1	D	397	0	408	3	0
1	E	387	0	400	10	0
2	A	37	0	0	0	0
2	B	21	0	0	1	0
2	C	19	0	0	0	0
2	D	30	0	0	2	0
2	E	14	0	0	0	0
All	All	2088	0	2018	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:329:HOH:O	1:E:190:LYS:HD2	1.80	0.82
1:E:175:GLN:O	1:E:179:GLN:HG2	1.92	0.68
1:A:198:GLU:OE2	1:B:203:GLN:NE2	2.34	0.60
1:B:163:GLN:HE22	1:C:161:GLN:HE22	1.51	0.59
1:B:203:GLN:O	1:B:207:LYS:HG2	2.11	0.51
1:D:160:SER:N	2:D:302:HOH:O	2.43	0.51
1:D:206:VAL:HG11	1:E:206:VAL:N	2.27	0.50
1:E:160:SER:HA	1:E:163:GLN:NE2	2.28	0.49
1:A:186:TYR:CE2	1:A:190:LYS:HE3	2.49	0.48
1:E:201:THR:O	1:E:205:MET:HG3	2.13	0.48
1:E:163:GLN:O	1:E:166:ALA:HB3	2.16	0.46
1:E:160:SER:HA	1:E:163:GLN:CD	2.36	0.45
1:C:205:MET:O	1:C:209:GLN:NE2	2.50	0.45
1:E:166:ALA:O	1:E:169:GLU:HB3	2.17	0.45
1:B:190:LYS:HG2	1:B:193:ARG:NH2	2.34	0.43
1:D:207:LYS:HG3	1:E:205:MET:HE1	2.02	0.42
1:B:181:LEU:HD12	1:C:178:LEU:CD1	2.50	0.42
1:A:161:GLN:HB2	1:C:163:GLN:HE22	1.85	0.42
1:E:160:SER:HA	1:E:163:GLN:OE1	2.20	0.41
1:B:192:ASN:HA	2:B:302:HOH:O	2.21	0.41
1:A:207:LYS:HG3	1:C:205:MET:CE	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	48/50 (96%)	47 (98%)	1 (2%)	0	100	100
1	B	48/50 (96%)	48 (100%)	0	0	100	100
1	C	48/50 (96%)	48 (100%)	0	0	100	100
1	D	48/50 (96%)	48 (100%)	0	0	100	100
1	E	47/50 (94%)	43 (92%)	4 (8%)	0	100	100
All	All	239/250 (96%)	234 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	45/45 (100%)	44 (98%)	1 (2%)	60	39
1	B	44/45 (98%)	44 (100%)	0	100	100
1	C	44/45 (98%)	43 (98%)	1 (2%)	58	37
1	D	45/45 (100%)	43 (96%)	2 (4%)	35	13
1	E	44/45 (98%)	39 (89%)	5 (11%)	7	1
All	All	222/225 (99%)	213 (96%)	9 (4%)	37	15

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	176	GLN
1	C	162	GLN
1	D	168	SER
1	D	172	GLN
1	E	160	SER
1	E	161	GLN
1	E	163	GLN
1	E	177	GLN
1	E	190	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	184	GLN
1	B	163	GLN
1	B	172	GLN
1	B	184	GLN
1	C	162	GLN
1	C	163	GLN
1	C	175	GLN
1	D	161	GLN
1	D	179	GLN
1	E	161	GLN
1	E	162	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	50/50 (100%)	0.00	1 (2%) 68 73	19, 26, 45, 52	0
1	B	50/50 (100%)	0.17	1 (2%) 68 73	22, 32, 49, 61	0
1	C	50/50 (100%)	0.48	4 (8%) 15 17	20, 34, 51, 83	0
1	D	50/50 (100%)	0.65	6 (12%) 6 6	17, 28, 69, 75	0
1	E	49/50 (98%)	0.86	9 (18%) 2 2	17, 34, 65, 80	0
All	All	249/250 (99%)	0.43	21 (8%) 14 15	17, 31, 62, 83	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	164	ILE	4.0
1	C	209	GLN	3.8
1	E	165	ALA	3.8
1	E	163	GLN	3.7
1	C	207	LYS	3.6
1	D	208	ALA	3.3
1	A	160	SER	3.1
1	E	162	GLN	3.0
1	D	162	GLN	2.9
1	D	209	GLN	2.8
1	C	186	TYR	2.7
1	E	164	ILE	2.6
1	B	186	TYR	2.6
1	D	163	GLN	2.6
1	D	161	GLN	2.5
1	E	161	GLN	2.5
1	E	167	LEU	2.4
1	C	208	ALA	2.3
1	E	160	SER	2.3
1	E	172	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	171	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.