



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2016 – 04:49 AM EDT

PDB ID : 5D8Z  
Title : Structrue of a lucidum protein  
Authors : Guo, R.; Li, Q.; Shang, N.; Liu, G.; Ko, T.P.; Chen, C.C.; Liu, W.  
Deposited on : 2015-08-18  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790

**i**

## X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	<div> <div> <div></div> <div>2%</div> </div> <div> <div></div> <div>77%</div> </div> <div> <div></div> <div>14%</div> </div> <div> <div></div> <div>7%</div> </div> </div>
1	B	347	<div> <div> <div></div> <div>%</div> </div> <div> <div></div> <div>76%</div> </div> <div> <div></div> <div>16%</div> </div> <div> <div></div> <div>7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CBI	A	401	-	-	-	X
2	CBI	B	401	-	-	-	X

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called endoglucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2432	1523	407	490	12			
1	B	324	Total	C	N	O	S	0	0	0
			2432	1523	407	490	12			

- Molecule 2 is CELLOBIOSE (three-letter code: CBI) (formula:  $C_{12}H_{22}O_{11}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			23	12	11		
2	B	1	Total	C	O	0	0
			23	12	11		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total 53	O 53	0	0
3	B	20	Total 20	O 20	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: endoglucanase



#### • Molecule 1: endoglucanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.43 Å   236.40 Å   83.58 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	25.00 – 2.70 24.21 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.0 (25.00-2.70) 93.2 (24.21-2.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.34 (at 2.71 Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.179   ,   0.236 0.186   ,   0.243	Depositor DCC
$R_{free}$ test set	1129 reflections (5.45%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.2	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 31.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4983	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CBI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	0/2493	0.78	1/3411 (0.0%)
1	B	0.63	0/2493	0.76	2/3411 (0.1%)
All	All	0.65	0/4986	0.77	3/6822 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	53	GLN	N-CA-CB	5.78	121.01	110.60
1	B	186	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	B	53	GLN	N-CA-C	-5.42	96.36	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	309	TRP	Peptide
1	A	72	PHE	Peptide
1	B	309	TRP	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2216	32	0
1	B	2432	0	2216	34	0
2	A	23	0	22	0	0
2	B	23	0	22	0	0
3	A	53	0	0	0	0
3	B	20	0	0	0	1
All	All	4983	0	4476	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:GLN:O	1:B:55:TYR:N	2.05	0.89
1:B:52:THR:O	1:B:53:GLN:CG	2.38	0.72
1:B:52:THR:O	1:B:53:GLN:HG3	1.93	0.69
1:A:52:THR:O	1:A:53:GLN:HG3	1.93	0.68
1:B:199:GLY:H	1:B:203:THR:HG22	1.60	0.66
1:A:51:LEU:O	1:A:53:GLN:O	2.13	0.65
1:A:309:TRP:CD1	1:A:310:ALA:HB2	2.30	0.65
1:A:148:ALA:O	1:A:150:LYS:O	2.14	0.65
1:A:325:THR:O	1:A:331:TRP:HA	1.99	0.62
1:B:53:GLN:C	1:B:55:TYR:H	2.03	0.61
1:A:259:ALA:HB3	1:A:260:PRO:HD3	1.83	0.61
1:B:26:GLY:HA3	1:B:74:VAL:O	2.00	0.60
1:B:35:GLY:O	1:B:42:CYS:HA	2.02	0.59
1:A:53:GLN:O	1:A:55:TYR:N	2.35	0.58
1:B:237:TYR:C	1:B:238:LEU:HD12	2.24	0.58
1:A:253:ASN:HD22	1:A:292:GLN:NE2	2.03	0.57
1:B:186:ARG:HE	1:B:222:ASN:ND2	2.03	0.57
1:A:42:CYS:O	1:A:44:ALA:N	2.39	0.56
1:B:141:ALA:HB1	1:B:184:ALA:HB2	1.87	0.55
1:A:53:GLN:C	1:A:55:TYR:H	2.09	0.55
1:A:193:GLN:N	1:A:193:GLN:OE1	2.40	0.55

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:ARG:HE	1:A:222:ASN:ND2	2.06	0.54
1:A:224:ASP:OD1	1:A:226:SER:OG	2.17	0.54
1:B:286:GLU:O	1:B:290:CYS:SG	2.66	0.54
1:B:274:ASN:HD21	1:B:277:THR:HB	1.73	0.53
1:B:52:THR:O	1:B:53:GLN:CB	2.58	0.51
1:B:224:ASP:C	1:B:224:ASP:OD1	2.50	0.50
1:A:52:THR:O	1:A:53:GLN:CG	2.59	0.50
1:B:93:ILE:O	1:B:93:ILE:HD12	2.11	0.50
1:A:81:TRP:H	1:A:121:ASN:HD21	1.60	0.50
1:A:233:ASP:OD1	1:A:275:THR:OG1	2.25	0.50
1:B:258:TRP:HE1	1:B:292:GLN:NE2	2.11	0.49
1:B:238:LEU:HD12	1:B:238:LEU:N	2.27	0.49
1:B:199:GLY:N	1:B:203:THR:HG22	2.27	0.49
1:A:258:TRP:HE1	1:A:292:GLN:NE2	2.11	0.48
1:A:253:ASN:O	1:A:257:ALA:HB3	2.14	0.48
1:A:199:GLY:H	1:A:203:THR:HG22	1.78	0.48
1:B:222:ASN:HB3	1:B:223:PRO:CD	2.44	0.48
1:B:264:TRP:O	1:B:268:ASN:ND2	2.41	0.48
1:B:53:GLN:O	1:B:54:TYR:C	2.52	0.46
1:A:77:LEU:HD12	1:A:105:VAL:HG22	1.98	0.46
1:A:53:GLN:C	1:A:55:TYR:N	2.69	0.45
1:B:168:VAL:HG11	1:B:174:TRP:HB2	1.98	0.45
1:A:199:GLY:H	1:A:203:THR:CG2	2.30	0.45
1:A:74:VAL:HA	1:A:114:SER:O	2.17	0.45
1:B:259:ALA:HB3	1:B:260:PRO:CD	2.47	0.45
1:B:259:ALA:HB3	1:B:260:PRO:HD3	1.99	0.44
1:B:24:PHE:O	1:B:305:GLY:HA3	2.17	0.44
1:A:235:HIS:CG	1:A:276:GLU:HB2	2.52	0.44
1:B:208:PHE:O	1:B:213:SER:HB2	2.17	0.44
1:A:258:TRP:HE1	1:A:292:GLN:HE21	1.63	0.44
1:B:233:ASP:OD1	1:B:275:THR:OG1	2.24	0.44
1:B:325:THR:O	1:B:331:TRP:HA	2.18	0.43
1:B:231:ILE:HG21	1:B:304:LEU:HD12	2.01	0.43
1:A:58:ASP:O	1:A:62:GLN:HG3	2.19	0.42
1:A:81:TRP:H	1:A:121:ASN:ND2	2.17	0.42
1:A:58:ASP:OD1	1:A:61:GLY:N	2.46	0.42
1:B:53:GLN:C	1:B:55:TYR:N	2.63	0.42
1:B:138:GLN:OE1	1:B:138:GLN:N	2.53	0.42
1:B:186:ARG:HG3	1:B:223:PRO:HD3	2.01	0.41
1:A:153:ASP:N	1:A:153:ASP:OD1	2.53	0.41
1:B:166:HIS:CD2	1:B:202:TRP:CH2	3.07	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:ASN:HB3	1:A:223:PRO:HD2	2.02	0.41
1:A:222:ASN:HB3	1:A:223:PRO:CD	2.51	0.41
1:A:70:ASP:O	1:A:344:ASN:HB2	2.21	0.41
1:B:98:TRP:NE1	1:B:150:LYS:HD3	2.36	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:520:HOH:O	3:B:520:HOH:O[3_555]	1.78	0.42

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/347 (93%)	296 (92%)	23 (7%)	3 (1%)	21	49
1	B	322/347 (93%)	290 (90%)	30 (9%)	2 (1%)	30	59
All	All	644/694 (93%)	586 (91%)	53 (8%)	5 (1%)	24	51

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	54	TYR
1	A	310	ALA
1	B	54	TYR
1	B	310	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/265 (94%)	234 (94%)	15 (6%)	24	50
1	B	249/265 (94%)	236 (95%)	13 (5%)	29	58
All	All	498/530 (94%)	470 (94%)	28 (6%)	26	54

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	VAL
1	A	45	SER
1	A	53	GLN
1	A	64	LYS
1	A	92	ASP
1	A	150	LYS
1	A	155	ASP
1	A	203	THR
1	A	221	THR
1	A	236	LYS
1	A	273	PHE
1	A	285	CYS
1	A	287	THR
1	A	332	THR
1	A	344	ASN
1	B	27	VAL
1	B	128	GLU
1	B	150	LYS
1	B	153	ASP
1	B	215	ASP
1	B	228	THR
1	B	256	ASN
1	B	273	PHE
1	B	316	GLN
1	B	319	VAL
1	B	320	LEU
1	B	328	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	332	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	137	ASN
1	A	187	GLN
1	A	222	ASN
1	A	274	ASN
1	A	292	GLN
1	A	316	GLN
1	B	121	ASN
1	B	126	ASN
1	B	137	ASN
1	B	222	ASN
1	B	256	ASN
1	B	274	ASN
1	B	292	GLN
1	B	313	ASN
1	B	328	ASN
1	B	344	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CBI	A	401	-	24,24,24	0.72	0	35,35,35	1.29	6 (17%)
2	CBI	B	401	-	24,24,24	0.58	0	35,35,35	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CBI	A	401	-	-	0/8/48/48	0/2/2/2
2	CBI	B	401	-	-	0/8/48/48	0/2/2/2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	CBI	O3-C3-C4	-2.35	105.06	110.36
2	A	401	CBI	O5-C5-C4	2.05	113.57	109.67
2	A	401	CBI	C1-O5-C5	2.58	118.80	113.74
2	A	401	CBI	O4-C4-C5	2.65	116.20	109.23
2	A	401	CBI	C1-C2-C3	2.78	115.50	109.98
2	A	401	CBI	O5'-C5'-C4'	2.94	116.05	109.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	324/347 (93%)	-0.23	6 (1%) 70 70	29, 42, 66, 83	0
1	B	324/347 (93%)	-0.18	5 (1%) 76 76	30, 48, 68, 93	0
All	All	648/694 (93%)	-0.20	11 (1%) 73 74	29, 45, 67, 93	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	328	ASN	3.3
1	A	117	VAL	2.9
1	B	345	ALA	2.6
1	A	119	VAL	2.5
1	A	283	ALA	2.5
1	B	307	VAL	2.5
1	B	161	VAL	2.3
1	A	255	ASP	2.3
1	A	256	ASN	2.2
1	A	275	THR	2.0
1	B	283	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CBI	B	401	23/23	0.85	0.39	8.55	52,73,105,112	0
2	CBI	A	401	23/23	0.80	0.38	4.29	64,82,90,93	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.