wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 07:27 PM BST

PDB ID	:	2DAF
Title	:	Solution Structure of the Novel Identified Ubiquitin-like Domain in the Human Hypothetical Protein FLJ35834
Authors	:	Zhao, C.; Kigawa, T.; Saito, K.; Koshiba, S.; Inoue, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2005-12-14

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 92%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	118	53% 19% 28%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:15-A:99 (85)	0.22	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 17, 19
2	14, 20
Single-model clusters	16; 18

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1837 atoms, of which 934 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called FLJ35834 protein.

Mol	Chain	Residues	Atoms					Trace
			Total	C	H	N	O	
1	A	118	1837	576	934	152	175	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q6NUH6
A	2	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	3	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	4	GLY	-	CLONING ARTIFACT	UNP Q6NUH6
A	5	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	6	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	7	GLY	-	CLONING ARTIFACT	UNP Q6NUH6
A	113	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	114	GLY	-	CLONING ARTIFACT	UNP Q6NUH6
A	115	PRO	-	CLONING ARTIFACT	UNP Q6NUH6
A	116	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	117	SER	-	CLONING ARTIFACT	UNP Q6NUH6
A	118	GLY	-	CLONING ARTIFACT	UNP Q6NUH6

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: FLJ35834 protein

Chain A: 53% 19% 28%

G1 S2 S3 G4 S5 S6 G7 Q8 E9 S10 V11 E12 D13 S14 L22 I23 P24 V25 F33 I38 L39 K40 Y41 L42 H45 H48 L49 V56 L57 Y61 L66 T71 V77 E81 I87 F88 R99 I100 D101 G102 L103 T104 D105 V106 S107 Q108 I109

I110 T111 V112 S113 G114 P115 S116 S117 G118

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

- Molecule 1: FLJ35834 protein

Chain A: 41% 30% • 28%

G1 S2 S3 G4 S5 S6 G7 Q8 E9 S10 V11 E12 D13 S14 V20 V21 L22 I23 P24 V25 F33 K34 V35 D36 L39 K40 Y41 L42 K43 D44 H45 F46 S47 H48 L49 P53 V56 L57 Y61 L66 L72 V73 G76 V77 K78 P79 Q80 E81 I87 F88

N91 L94 Y95 R99 I100 D101 G102 L103 T104 D105 V106 S107 Q108 I109 I110 T111 V112 S113 G114 P115 S116 S117 G118

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics, restrainted molecular dynamics.*

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function,structures with the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.17
CYANA	refinement	2.0.17

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 11270
Number of chemical shift lists	1
Total number of shifts	1384
Number of shifts mapped to atoms	1384
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	92%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	685	734	730	26±3
All	All	13700	14680	14600	526

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

5 of 129 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:77:VAL:HG13	1:A:81:GLU:OE2	0.91	1.66	14	19
1:A:38:ILE:HD11	1:A:41:TYR:CD2	0.77	2.15	19	3
1:A:61:TYR:HB3	1:A:66:LEU:HD11	0.76	1.58	16	10
1:A:91:ASN:ND2	1:A:94:LEU:HD12	0.74	1.98	15	6
1:A:22:LEU:HD13	1:A:25:VAL:HG21	0.73	1.57	15	20

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	85/118 (72%)	77±2 (91±2%)	7±2 (8±2%)	1±1 (1±1%)	26 73
All	All	1700/2360 (72%)	1542 (91%)	142 (8%)	16 (1%)	26 73

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	38	ILE	11
1	A	76	GLY	2
1	A	99	ARG	1
1	A	44	ASP	1
1	A	54	HIS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	80/107 (75%)	77±2 (96±2%)	3±2 (4±2%)	40 83
All	All	1600/2140 (75%)	1531 (96%)	69 (4%)	40 83

5 of 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	49	LEU	9
1	A	71	THR	9
1	A	99	ARG	8
1	A	64	LYS	7
1	A	34	LYS	5

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 92% for the well-defined parts and 85% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 11270

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1384
Number of shifts mapped to atoms	1384
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	107	-0.24 ± 0.17	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	100	0.15 ± 0.12	None needed (< 0.5 ppm)
$^{13}C'$	103	-0.11 ± 0.14	None needed (< 0.5 ppm)
^{15}N	97	0.45 ± 0.47	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 92%, i.e. 1022 atoms were assigned a chemical shift out of a possible 1109. 0 out of 22 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	412/413 (100%)	163/164 (99%)	170/170 (100%)	79/79 (100%)
Sidechain	543/613 (89%)	329/357 (92%)	207/232 (89%)	7/24 (29%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	67/83 (81%)	35/43 (81%)	32/32 (100%)	0/8 (0%)
Overall	1022/1109 (92%)	527/564 (93%)	409/434 (94%)	86/111 (77%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	58	GLN	HB2	0.30	3.30 – 0.80	-7.0

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: