



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:05 AM GMT

PDB ID : 3DCB
Title : Crystal structure of the Drosophila kinesin family member NOD in complex with AMPPNP
Authors : Cochran, J.C.; Mulko, N.K.; Kull, F.J.
Deposited on : 2008-06-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	<div> <div>9%</div> <div>58%</div> <div>26%</div> <div>•</div> <div>13%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin-like protein Nod.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	299	2320	1463	416	427	14	0	0	0


There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	EXPRESSION TAG	UNP P18105
A	-13	ALA	-	EXPRESSION TAG	UNP P18105
A	-12	SER	-	EXPRESSION TAG	UNP P18105
A	-11	ASP	-	EXPRESSION TAG	UNP P18105
A	-10	TYR	-	EXPRESSION TAG	UNP P18105
A	-9	LYS	-	EXPRESSION TAG	UNP P18105
A	-8	ASP	-	EXPRESSION TAG	UNP P18105
A	-7	ASP	-	EXPRESSION TAG	UNP P18105
A	-6	ASP	-	EXPRESSION TAG	UNP P18105
A	-5	ASP	-	EXPRESSION TAG	UNP P18105
A	-4	LYS	-	EXPRESSION TAG	UNP P18105
A	-3	ARG	-	EXPRESSION TAG	UNP P18105
A	-2	ARG	-	EXPRESSION TAG	UNP P18105
A	-1	ARG	-	EXPRESSION TAG	UNP P18105
A	0	GLY	-	EXPRESSION TAG	UNP P18105
A	319	ALA	-	EXPRESSION TAG	UNP P18105
A	320	ALA	-	EXPRESSION TAG	UNP P18105
A	321	ALA	-	EXPRESSION TAG	UNP P18105
A	322	LEU	-	EXPRESSION TAG	UNP P18105
A	323	GLU	-	EXPRESSION TAG	UNP P18105
A	324	HIS	-	EXPRESSION TAG	UNP P18105
A	325	HIS	-	EXPRESSION TAG	UNP P18105
A	326	HIS	-	EXPRESSION TAG	UNP P18105
A	327	HIS	-	EXPRESSION TAG	UNP P18105
A	328	HIS	-	EXPRESSION TAG	UNP P18105
A	329	HIS	-	EXPRESSION TAG	UNP P18105

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kinesin-like protein Nod



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.86Å 74.78Å 79.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.00 – 2.50 19.59 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.1 (19.00-2.50) 97.2 (19.59-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	10.90	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 2.50Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.209 , 0.275 0.218 , 0.268	Depositor DCC
R_{free} test set	570 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 62.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 10899 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2501	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/2367	0.57	0/3209

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	265	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2320	0	2333	70	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	31	0	13	0	0
4	A	149	0	0	1	0
All	All	2501	0	2346	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:ARG:NH2	1:A:228:ALA:HB1	1.70	1.07
1:A:234:ARG:NH2	1:A:234:ARG:HB2	1.85	0.91
1:A:204:ARG:NH2	1:A:228:ALA:CB	2.36	0.88
1:A:267:HIS:O	1:A:268:THR:HB	1.81	0.79
1:A:264:ALA:O	1:A:265:ALA:HB2	1.86	0.75
1:A:10:ARG:HD2	4:A:472:HOH:O	1.88	0.73
1:A:204:ARG:HH21	1:A:228:ALA:HB1	1.53	0.72
1:A:138:TYR:HB2	1:A:212:HIS:HB2	1.72	0.72
1:A:234:ARG:HH21	1:A:234:ARG:HB2	1.55	0.70
1:A:127:ARG:NH1	1:A:132:LYS:HD2	2.07	0.70
1:A:14:ARG:HH12	1:A:60:GLN:NE2	1.90	0.69
1:A:250:ASN:HD22	1:A:253:LEU:H	1.41	0.69
1:A:264:ALA:O	1:A:265:ALA:CB	2.41	0.68
1:A:105:ILE:HD11	1:A:110:LEU:HD23	1.76	0.68
1:A:268:THR:O	1:A:269:VAL:HG23	1.95	0.67
1:A:266:GLY:O	1:A:267:HIS:HB2	1.95	0.66
1:A:88:GLN:HE22	1:A:303:ASP:HB3	1.59	0.66
1:A:142:ILE:HD11	1:A:169:CYS:SG	2.37	0.65
1:A:103:GLY:H	1:A:181:HIS:HD2	1.43	0.65
1:A:14:ARG:HH12	1:A:60:GLN:HE22	1.45	0.64
1:A:250:ASN:HB3	1:A:253:LEU:HD23	1.81	0.62
1:A:106:LEU:HB3	1:A:108:GLU:OE1	2.00	0.62
1:A:270:ILE:HB	1:A:272:TYR:CE1	2.35	0.61
1:A:162:VAL:HG12	1:A:163:ALA:N	2.16	0.60
1:A:267:HIS:O	1:A:268:THR:CB	2.49	0.60
1:A:59:SER:OG	1:A:62:GLU:HG3	2.01	0.59
1:A:250:ASN:ND2	1:A:253:LEU:H	2.01	0.59
1:A:234:ARG:HB2	1:A:234:ARG:CZ	2.34	0.57
1:A:82:THR:HG23	1:A:223:ASN:HB2	1.86	0.57
1:A:139:ALA:HB2	1:A:174:LEU:HD11	1.85	0.57
1:A:151:ASP:OD2	1:A:170:THR:HG23	2.05	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:ASP:OD1	1:A:123:ARG:NH1	2.38	0.57
1:A:162:VAL:HG12	1:A:163:ALA:H	1.69	0.56
1:A:97:MET:CE	1:A:294:LEU:HD23	2.36	0.56
1:A:108:GLU:CD	1:A:108:GLU:H	2.10	0.55
1:A:304:LEU:O	1:A:308:LEU:HD13	2.07	0.55
1:A:203:SER:O	1:A:204:ARG:HG2	2.08	0.53
1:A:97:MET:HE1	1:A:294:LEU:HD23	1.91	0.53
1:A:230:SER:HB2	1:A:234:ARG:NH1	2.25	0.51
1:A:41:LEU:HD23	1:A:41:LEU:N	2.24	0.51
1:A:260:VAL:O	1:A:264:ALA:HB2	2.11	0.51
1:A:144:ILE:HD13	1:A:276:VAL:HG23	1.93	0.50
1:A:69:LEU:HB3	1:A:70:PRO:HD3	1.93	0.49
1:A:80:GLN:OE1	1:A:221:ARG:HG2	2.12	0.49
1:A:250:ASN:HD22	1:A:253:LEU:HD22	1.78	0.49
1:A:102:PRO:HG2	1:A:180:LEU:HD23	1.95	0.49
1:A:255:SER:O	1:A:259:VAL:HG23	2.12	0.48
1:A:257:ASN:OD1	1:A:317:ALA:HA	2.13	0.48
1:A:204:ARG:HH22	1:A:228:ALA:HA	1.79	0.47
1:A:21:PHE:O	1:A:24:ARG:HG2	2.14	0.47
1:A:20:GLN:HB3	1:A:25:ARG:HE	1.80	0.47
1:A:308:LEU:O	1:A:312:ARG:HG3	2.16	0.46
1:A:269:VAL:HG12	1:A:269:VAL:O	2.16	0.46
1:A:188:THR:O	1:A:191:ARG:HG2	2.16	0.45
1:A:5:LYS:O	1:A:287:ALA:HA	2.17	0.45
1:A:132:LYS:HB3	1:A:132:LYS:NZ	2.32	0.44
1:A:29:VAL:HG21	1:A:304:LEU:HD13	1.98	0.44
1:A:182:HIS:CD2	1:A:183:ILE:HD12	2.53	0.44
1:A:126:ALA:O	1:A:129:GLU:HG2	2.18	0.44
1:A:33:PRO:HB2	1:A:35:TRP:CE2	2.53	0.43
1:A:225:VAL:HG12	1:A:227:LEU:HD13	2.00	0.42
1:A:120:ILE:O	1:A:124:VAL:HG23	2.19	0.42
1:A:143:GLU:HB3	1:A:152:LEU:HD11	2.02	0.42
1:A:204:ARG:NH2	1:A:228:ALA:CA	2.83	0.41
1:A:123:ARG:O	1:A:126:ALA:HB3	2.20	0.41
1:A:250:ASN:HD22	1:A:253:LEU:N	2.14	0.41
1:A:13:VAL:HA	1:A:295:ALA:O	2.20	0.41
1:A:64:TYR:CD1	1:A:68:ILE:HB	2.56	0.41
1:A:256:ILE:O	1:A:260:VAL:HG23	2.21	0.40
1:A:41:LEU:H	1:A:41:LEU:HD23	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/344 (85%)	270 (92%)	18 (6%)	5 (2%)	11	19

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	ALA
1	A	267	HIS
1	A	266	GLY
1	A	268	THR
1	A	131	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	256/293 (87%)	241 (94%)	15 (6%)	24	44

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	24	ARG
1	A	94	SER
1	A	130	ASN
1	A	132	LYS
1	A	141	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	153	LEU
1	A	165	ARG
1	A	168	ARG
1	A	184	LEU
1	A	204	ARG
1	A	227	LEU
1	A	234	ARG
1	A	253	LEU
1	A	298	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	46	ASN
1	A	60	GLN
1	A	65	GLN
1	A	88	GLN
1	A	146	ASN
1	A	158	HIS
1	A	167	GLN
1	A	181	HIS
1	A	182	HIS
1	A	206	HIS
1	A	250	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ANP	A	401	2	27,33,33	1.43	5 (18%)	30,52,52	2.51	9 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	401	2	-	0/12/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	ANP	C8-N7	-2.00	1.30	1.34
3	A	401	ANP	PB-O3A	2.45	1.62	1.59
3	A	401	ANP	PG-N3B	2.71	1.70	1.63
3	A	401	ANP	O4'-C1'	2.82	1.44	1.41
3	A	401	ANP	PB-N3B	3.04	1.71	1.63

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	ANP	N3-C2-N1	-9.40	121.69	128.89
3	A	401	ANP	PA-O3A-PB	-6.10	112.20	132.67
3	A	401	ANP	C2'-C1'-N9	-3.20	109.40	114.29
3	A	401	ANP	O1B-PB-N3B	-3.08	107.17	111.90
3	A	401	ANP	O3'-C3'-C2'	-2.83	102.64	111.83
3	A	401	ANP	C4-C5-N7	-2.76	106.94	109.48
3	A	401	ANP	C2'-C3'-C4'	-2.14	98.21	102.61
3	A	401	ANP	O2B-PB-O3A	2.12	114.70	105.09
3	A	401	ANP	O3A-PA-O5'	2.12	108.57	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/344 (86%)	0.56	32 (10%) 8 8	10, 29, 69, 98	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	ASN	8.8
1	A	247	VAL	7.5
1	A	248	ASN	6.5
1	A	265	ALA	6.1
1	A	267	HIS	5.8
1	A	268	THR	5.1
1	A	128	GLN	4.7
1	A	204	ARG	4.6
1	A	202	SER	4.5
1	A	130	ASN	4.4
1	A	249	ILE	4.4
1	A	203	SER	4.4
1	A	266	GLY	4.0
1	A	132	LYS	3.7
1	A	236	THR	3.5
1	A	104	GLU	2.9
1	A	106	LEU	2.9
1	A	129	GLU	2.8
1	A	133	ASP	2.8
1	A	233	VAL	2.8
1	A	168	ARG	2.8
1	A	302	CYS	2.7
1	A	231	GLU	2.5
1	A	105	ILE	2.5
1	A	234	ARG	2.4
1	A	269	VAL	2.3
1	A	163	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	SER	2.2
1	A	164	ALA	2.1
1	A	232	GLY	2.1
1	A	22	LEU	2.1
1	A	102	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ANP	A	401	31/31	0.94	0.17	-0.10	21,37,44,46	0
2	MG	A	402	1/1	0.95	0.06	-	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.