



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:31 AM GMT

PDB ID : 2DDU
Title : Crystal structure of the third repeat domain of reelin
Authors : Nogi, T.; Yasui, N.; Takagi, J.
Deposited on : 2006-02-03
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1192 (2.04-2.04)
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2465 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called reelin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2362	1504	407	437	14			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1220	GLY	-	CLONING ARTIFACT	UNP Q64FW1
A	1221	ARG	-	CLONING ARTIFACT	UNP Q64FW1
A	1598	SER	-	CLONING ARTIFACT	UNP Q64FW1
A	1599	ARG	-	CLONING ARTIFACT	UNP Q64FW1
A	1600	LEU	-	CLONING ARTIFACT	UNP Q64FW1
A	1601	GLU	-	CLONING ARTIFACT	UNP Q64FW1
A	1602	ASN	-	CLONING ARTIFACT	UNP Q64FW1
A	1603	LEU	-	CLONING ARTIFACT	UNP Q64FW1
A	1604	TYR	-	CLONING ARTIFACT	UNP Q64FW1
A	1605	PHE	-	CLONING ARTIFACT	UNP Q64FW1
A	1606	GLN	-	CLONING ARTIFACT	UNP Q64FW1

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	100	Total	O	0	0
			100	100		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: reelin



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	129.93Å 129.93Å 122.48Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.05 41.43 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.05) 100.0 (41.43-2.05)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.234 , 0.265 0.233 , 0.258	Depositor DCC
R_{free} test set	1254 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 64.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 25058 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2465	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2425	0.62	0/3291

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2362	0	2292	26	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	100	0	0	0	0
All	All	2465	0	2292	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1424:ILE:HG13	1:A:1429:PHE:HD1	1.50	0.75
1:A:1332:TRP:CZ3	1:A:1388:ARG:HB2	2.24	0.73
1:A:1424:ILE:HG13	1:A:1429:PHE:CD1	2.25	0.71
1:A:1318:ALA:HB1	1:A:1391:GLN:O	2.03	0.57
1:A:1297:LYS:HB2	1:A:1300:TYR:CE1	2.39	0.57
1:A:1404:LEU:O	1:A:1405:ASP:HB2	2.05	0.55
1:A:1487:ASN:HD22	1:A:1487:ASN:H	1.57	0.52
1:A:1378:ARG:HD3	1:A:1378:ARG:H	1.75	0.51
1:A:1350:GLY:O	1:A:1353:ARG:HD2	2.14	0.47
1:A:1454:ARG:NH2	1:A:1480:ASP:OD2	2.48	0.47
1:A:1474:GLY:HA3	1:A:1485:TYR:CZ	2.51	0.46
1:A:1391:GLN:HE22	1:A:1401:PRO:HA	1.81	0.46
1:A:1536:TYR:HD2	1:A:1547:LEU:HG	1.80	0.46
1:A:1361:TYR:HB3	1:A:1366:PHE:CE2	2.50	0.46
1:A:1528:ARG:HG2	1:A:1552:PHE:CE2	2.53	0.44
1:A:1452:PHE:HA	1:A:1595:VAL:O	2.18	0.44
1:A:1435:THR:HG23	1:A:1444:ASN:HD21	1.82	0.44
1:A:1533:VAL:HG12	1:A:1535:GLN:HG3	2.02	0.42
1:A:1322:LEU:HB3	1:A:1359:THR:HB	2.00	0.42
1:A:1303:GLN:HE22	1:A:1423:CYS:HB3	1.84	0.42
1:A:1384:LYS:HG2	1:A:1384:LYS:O	2.20	0.42
1:A:1391:GLN:O	1:A:1392:GLU:HB2	2.19	0.42
1:A:1459:LEU:HD21	1:A:1484:LEU:HD13	2.02	0.42
1:A:1311:THR:HG22	1:A:1313:GLN:H	1.83	0.41
1:A:1487:ASN:HD22	1:A:1487:ASN:N	2.16	0.41
1:A:1358:PRO:HB2	1:A:1360:VAL:HG23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	297/387 (77%)	276 (93%)	16 (5%)	5 (2%)	11 3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1379	SER
1	A	1298	PRO
1	A	1382	SER
1	A	1405	ASP
1	A	1309	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	257/333 (77%)	243 (95%)	14 (5%)	27 17

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1306	LEU
1	A	1311	THR
1	A	1357	GLU
1	A	1363	THR
1	A	1368	GLU
1	A	1378	ARG
1	A	1379	SER
1	A	1384	LYS
1	A	1424	ILE
1	A	1427	VAL
1	A	1470	GLN
1	A	1487	ASN
1	A	1599	ARG
1	A	1600	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1391	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1444	ASN
1	A	1448	HIS
1	A	1470	GLN
1	A	1487	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/387 (77%)	1.36	73 (24%) 1 1	24, 46, 84, 87	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1429	PHE	8.2
1	A	1344	ALA	7.5
1	A	1432	LEU	6.7
1	A	1333	PHE	6.4
1	A	1317	THR	5.8
1	A	1406	GLY	5.6
1	A	1330	MET	5.5
1	A	1314	PHE	5.3
1	A	1387	PHE	5.3
1	A	1405	ASP	5.2
1	A	1350	GLY	5.0
1	A	1364	GLY	5.0
1	A	1346	LYS	5.0
1	A	1438	GLN	4.9
1	A	1332	TRP	4.8
1	A	1402	PHE	4.8
1	A	1399	VAL	4.8
1	A	1294	LEU	4.8
1	A	1308	ILE	4.7
1	A	1400	PRO	4.6
1	A	1401	PRO	4.5
1	A	1318	ALA	4.4
1	A	1378	ARG	4.4
1	A	1363	THR	4.3
1	A	1296	LEU	4.1
1	A	1388	ARG	3.9
1	A	1309	GLY	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1403	GLY	3.8
1	A	1324	TYR	3.8
1	A	1349	GLU	3.7
1	A	1567	ARG	3.6
1	A	1424	ILE	3.4
1	A	1380	LEU	3.4
1	A	1389	TRP	3.4
1	A	1427	VAL	3.4
1	A	1352	SER	3.4
1	A	1391	GLN	3.3
1	A	1369	TRP	3.2
1	A	1311	THR	3.2
1	A	1331	SER	3.1
1	A	1390	ILE	3.0
1	A	1320	VAL	3.0
1	A	1351	ASN	2.9
1	A	1306	LEU	2.8
1	A	1382	SER	2.8
1	A	1301	VAL	2.8
1	A	1404	LEU	2.8
1	A	1326	HIS	2.8
1	A	1316	SER	2.7
1	A	1425	SER	2.7
1	A	1329	GLY	2.7
1	A	1381	ALA	2.6
1	A	1343	SER	2.5
1	A	1415	SER	2.5
1	A	1600	LEU	2.5
1	A	1436	ALA	2.4
1	A	1505	ARG	2.4
1	A	1313	GLN	2.4
1	A	1348	CYS	2.4
1	A	1297	LYS	2.4
1	A	1354	GLU	2.4
1	A	1325	SER	2.4
1	A	1431	ASP	2.3
1	A	1328	ALA	2.3
1	A	1408	TYR	2.3
1	A	1370	THR	2.2
1	A	1414	PRO	2.2
1	A	1599	ARG	2.2
1	A	1327	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1486	PHE	2.1
1	A	1449	SER	2.1
1	A	1590	TRP	2.1
1	A	1503	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	CA	A	1	1/1	0.98	0.09	-1.49	31,31,31,31	0
3	CL	A	2	1/1	0.99	0.16	-	25,25,25,25	1
4	MG	A	3	1/1	0.92	0.03	-	45,45,45,45	1

6.5 Other polymers [i](#)

There are no such residues in this entry.