



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:08 AM GMT

PDB ID : 3DH3
Title : Crystal Structure of RluF in complex with a 22 nucleotide RNA substrate
Authors : Alian, A.; DeGiovanni, A.; Stroud, R.M.; Finer-Moore, J.S.
Deposited on : 2008-06-16
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)
RNA backbone	2183	1036 (3.40-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	 2% 53% 25% 17%
1	B	290	 2% 54% 23% 5% 17%
1	C	290	 54% 24% 18%
1	D	290	 2% 51% 27% 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	22	
2	F	22	
2	G	22	
2	H	22	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal large subunit pseudouridine synthase F.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	240	Total 1902	C 1194	N 340	O 362	S 6	0	0	0
1	B	241	Total 1909	C 1199	N 341	O 363	S 6	0	0	0
1	C	239	Total 1895	C 1189	N 339	O 361	S 6	0	0	0
1	D	239	Total 1895	C 1189	N 339	O 361	S 6	0	0	0

- Molecule 2 is a RNA chain called stem loop fragment of E. Coli 23S RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	F	N	O				P
2	E	22	Total 475	C 212	F 1	N 90	O 151	P 21	0	0	0
2	F	22	Total 475	C 212	F 1	N 90	O 151	P 21	0	0	0
2	G	22	Total 475	C 212	F 1	N 90	O 151	P 21	0	0	0
2	H	22	Total 475	C 212	F 1	N 90	O 151	P 21	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	12	Total 12	O 12	0	0
3	E	5	Total 5	O 5	0	0
3	B	9	Total 9	O 9	0	0
3	F	1	Total 1	O 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	15	Total 15	O 15	0	0
3	G	3	Total 3	O 3	0	0
3	D	3	Total 3	O 3	0	0
3	H	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribosomal large subunit pseudouridine synthase F




- Molecule 1: Ribosomal large subunit pseudouridine synthase F




- Molecule 1: Ribosomal large subunit pseudouridine synthase F





- Molecule 1: Ribosomal large subunit pseudouridine synthase F



- Molecule 2: stem loop fragment of E. Coli 23S RNA



- Molecule 2: stem loop fragment of E. Coli 23S RNA



- Molecule 2: stem loop fragment of E. Coli 23S RNA



- Molecule 2: stem loop fragment of E. Coli 23S RNA



A2587	G2588	A2589	A2590	G2591	G2592	U2593	C2594	G2595	U2596	G2597	A2598	G2599	A2600	C2601	A2602	G2603	U2604	U2605	C2606	G2607	G2608
-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.55Å 84.13Å 91.35Å 90.00° 94.87° 90.00°	Depositor
Resolution (Å)	49.51 – 3.00 49.48 – 3.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (49.51-3.00) 98.7 (49.48-3.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, R_{free}	0.231 , 0.279 0.237 , 0.286	Depositor DCC
R_{free} test set	1355 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.0	EDS
Estimated twinning fraction	0.026 for l,-k,h	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 27006 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	9550	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FHU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.65	0/1928	0.76	1/2596 (0.0%)
1	B	0.66	0/1936	0.76	0/2607
1	C	0.66	0/1921	0.73	0/2586
1	D	0.63	0/1921	0.76	1/2586 (0.0%)
2	E	1.33	0/508	2.43	34/792 (4.3%)
2	F	1.38	0/508	2.41	34/792 (4.3%)
2	G	1.58	3/508 (0.6%)	2.64	45/792 (5.7%)
2	H	1.28	0/508	2.37	37/792 (4.7%)
All	All	0.86	3/9738 (0.0%)	1.36	152/13543 (1.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	2597	G	N9-C4	-10.44	1.29	1.38
2	G	2597	G	N3-C4	-7.42	1.30	1.35
2	G	2597	G	C5-C4	-5.26	1.34	1.38

The worst 5 of 152 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2597	G	C4'-C3'-C2'	-17.21	85.39	102.60
2	G	2597	G	N3-C4-N9	-16.07	116.36	126.00
2	F	2597	G	C4'-C3'-C2'	-14.02	88.58	102.60
2	G	2597	G	N3-C4-C5	13.58	135.39	128.60
2	H	2597	G	N3-C4-N9	-13.21	118.07	126.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1902	0	1942	62	0
1	B	1909	0	1950	67	0
1	C	1895	0	1933	67	0
1	D	1895	0	1933	75	0
2	E	475	0	241	27	0
2	F	475	0	241	27	0
2	G	475	0	241	26	0
2	H	475	0	241	33	0
3	A	12	0	0	1	0
3	B	9	0	0	2	0
3	C	15	0	0	1	0
3	D	3	0	0	0	0
3	E	5	0	0	0	0
3	F	1	0	0	0	0
3	G	3	0	0	0	0
3	H	1	0	0	1	0
All	All	9550	0	8722	334	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 334 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:51:VAL:HG11	1:C:218:LEU:O	1.45	1.15
2:G:2596:U:O2'	2:G:2597:G:O5'	1.70	1.10
1:D:236:ILE:HG23	1:D:238:ASN:ND2	1.70	1.07
1:C:8:ARG:HH21	2:G:2597:G:N2	1.52	1.06
1:C:236:ILE:HG23	1:C:238:ASN:ND2	1.71	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/290 (82%)	214 (90%)	21 (9%)	3 (1%)	15	53
1	B	239/290 (82%)	210 (88%)	24 (10%)	5 (2%)	9	40
1	C	237/290 (82%)	213 (90%)	19 (8%)	5 (2%)	9	40
1	D	237/290 (82%)	209 (88%)	23 (10%)	5 (2%)	9	40
All	All	951/1160 (82%)	846 (89%)	87 (9%)	18 (2%)	10	43

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	GLU
1	B	5	SER
1	B	237	GLU
1	C	5	SER
1	D	237	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/253 (85%)	190 (89%)	24 (11%)	7	29
1	B	215/253 (85%)	191 (89%)	24 (11%)	7	29
1	C	213/253 (84%)	189 (89%)	24 (11%)	7	28
1	D	213/253 (84%)	190 (89%)	23 (11%)	8	30
All	All	855/1012 (84%)	760 (89%)	95 (11%)	8	29

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	222	ARG
1	C	47	LYS
1	D	163	VAL
1	B	229	LEU
1	C	13	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	238	ASN
1	C	55	ASN
1	D	119	HIS
1	C	32	ASN
1	C	95	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	22/22 (100%)	11 (50%)	4 (18%)
2	F	22/22 (100%)	10 (45%)	3 (13%)
2	G	22/22 (100%)	9 (40%)	5 (22%)
2	H	22/22 (100%)	9 (40%)	4 (18%)
All	All	88/88 (100%)	39 (44%)	16 (18%)

5 of 39 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	2588	G
2	E	2589	A
2	E	2595	G
2	E	2596	U
2	E	2597	G

5 of 16 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	G	2587	A
2	G	2588	G
2	H	2587	A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	2597	G
2	H	2588	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FHU	E	2604	2	15,23,24	1.02	0	17,35,38	1.51	3 (17%)
2	FHU	F	2604	2	15,23,24	1.12	2 (13%)	17,35,38	1.75	4 (23%)
2	FHU	G	2604	2	15,23,24	1.18	1 (6%)	17,35,38	1.85	7 (41%)
2	FHU	H	2604	2	15,23,24	1.20	3 (20%)	17,35,38	2.11	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FHU	E	2604	2	-	0/3/47/48	0/2/2/2
2	FHU	F	2604	2	-	0/3/47/48	0/2/2/2
2	FHU	G	2604	2	-	0/3/47/48	0/2/2/2
2	FHU	H	2604	2	-	0/3/47/48	0/2/2/2

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	2604	FHU	C2'-C1'	-2.55	1.49	1.53
2	F	2604	FHU	C4-N3	-2.51	1.33	1.37
2	H	2604	FHU	C2-N3	-2.21	1.33	1.37
2	F	2604	FHU	C2-N3	-2.19	1.33	1.37
2	H	2604	FHU	C4-N3	-2.16	1.33	1.37

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	2604	FHU	O2'-C2'-C1'	-3.30	104.35	112.90
2	F	2604	FHU	O2'-C2'-C1'	-3.10	104.86	112.90
2	G	2604	FHU	C4-N3-C2	-2.96	121.56	126.00
2	G	2604	FHU	O2'-C2'-C1'	-2.91	105.37	112.90
2	E	2604	FHU	O2'-C2'-C1'	-2.62	106.12	112.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2604	FHU	1	0
2	G	2604	FHU	1	0
2	H	2604	FHU	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/290 (82%)	0.01	0 100 100	37, 52, 65, 78	0
1	B	241/290 (83%)	0.22	5 (2%) 67 36	37, 52, 67, 78	0
1	C	239/290 (82%)	-0.03	1 (0%) 93 80	36, 52, 65, 78	0
1	D	239/290 (82%)	0.26	3 (1%) 79 53	37, 52, 65, 78	0
2	E	21/22 (95%)	-0.18	1 (4%) 34 14	30, 41, 67, 90	0
2	F	21/22 (95%)	-0.13	1 (4%) 34 14	29, 41, 67, 90	0
2	G	21/22 (95%)	-0.11	0 100 100	28, 41, 66, 90	0
2	H	21/22 (95%)	-0.05	1 (4%) 34 14	31, 43, 68, 90	0
All	All	1043/1248 (83%)	0.10	12 (1%) 81 55	28, 51, 67, 90	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	2608	G	4.7
1	D	56	GLY	3.8
1	B	49	GLY	3.1
1	B	173	ALA	3.0
1	B	51	VAL	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FHU	H	2604	22/23	0.96	0.15	-	30,34,38,39	0
2	FHU	F	2604	22/23	0.97	0.13	-	31,34,39,41	0
2	FHU	G	2604	22/23	0.98	0.15	-	28,33,38,39	0
2	FHU	E	2604	22/23	0.97	0.14	-	27,33,40,40	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.