



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2016 – 06:19 PM EDT

PDB ID : 5DHP
Title : Crystal structure of NAD kinase 1 from *Listeria monocytogenes* in complex with a novel inhibitor
Authors : Gelin, M.; Paoletti, J.; Assairi, L.; Huteau, V.; Pochet, S.; Labesse, G.
Deposited on : 2015-08-31
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20028320
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20028320


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5193 (2.30-2.26)
Clashscore	102246	5929 (2.30-2.26)
Ramachandran outliers	100387	5851 (2.30-2.26)
Sidechain outliers	100360	5850 (2.30-2.26)
RSRZ outliers	91569	5204 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>3%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	B	272	<div> <div>6%</div> <div>84%</div> <div>12%</div> <div>..</div> </div>
1	C	272	<div> <div>10%</div> <div>88%</div> <div>8%</div> <div>.</div> </div>
1	D	272	<div> <div>15%</div> <div>78%</div> <div>14%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	302	-	-	-	X
4	5AQ	D	302	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2064	1322	345	388	9			
1	B	263	Total	C	N	O	S	0	2	0
			2113	1356	354	394	9			
1	C	261	Total	C	N	O	S	0	0	0
			2035	1303	342	381	9			
1	D	250	Total	C	N	O	S	0	1	0
			1953	1249	337	360	7			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	265	LEU	-	expression tag	UNP Q8Y8D7
A	266	GLU	-	expression tag	UNP Q8Y8D7
A	267	HIS	-	expression tag	UNP Q8Y8D7
A	268	HIS	-	expression tag	UNP Q8Y8D7
A	269	HIS	-	expression tag	UNP Q8Y8D7
A	270	HIS	-	expression tag	UNP Q8Y8D7
A	271	HIS	-	expression tag	UNP Q8Y8D7
A	272	HIS	-	expression tag	UNP Q8Y8D7
B	265	LEU	-	expression tag	UNP Q8Y8D7
B	266	GLU	-	expression tag	UNP Q8Y8D7
B	267	HIS	-	expression tag	UNP Q8Y8D7
B	268	HIS	-	expression tag	UNP Q8Y8D7
B	269	HIS	-	expression tag	UNP Q8Y8D7
B	270	HIS	-	expression tag	UNP Q8Y8D7
B	271	HIS	-	expression tag	UNP Q8Y8D7
B	272	HIS	-	expression tag	UNP Q8Y8D7
C	265	LEU	-	expression tag	UNP Q8Y8D7
C	266	GLU	-	expression tag	UNP Q8Y8D7
C	267	HIS	-	expression tag	UNP Q8Y8D7
C	268	HIS	-	expression tag	UNP Q8Y8D7
C	269	HIS	-	expression tag	UNP Q8Y8D7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	270	HIS	-	expression tag	UNP Q8Y8D7
C	271	HIS	-	expression tag	UNP Q8Y8D7
C	272	HIS	-	expression tag	UNP Q8Y8D7
D	265	LEU	-	expression tag	UNP Q8Y8D7
D	266	GLU	-	expression tag	UNP Q8Y8D7
D	267	HIS	-	expression tag	UNP Q8Y8D7
D	268	HIS	-	expression tag	UNP Q8Y8D7
D	269	HIS	-	expression tag	UNP Q8Y8D7
D	270	HIS	-	expression tag	UNP Q8Y8D7
D	271	HIS	-	expression tag	UNP Q8Y8D7
D	272	HIS	-	expression tag	UNP Q8Y8D7

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		
2	D	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 8-({2-oxo-2-[(2-phenylethyl)amino]ethyl}sulfanyl)adenosine (three-letter code: 5AQ) (formula: C₂₀H₂₄N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			32	20	6	5	1		
4	B	1	Total	C	N	O	S	0	0
			32	20	6	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			32	20	6	5	1		
4	D	1	Total	C	N	O	S	0	0
			32	20	6	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	86	Total	O	0	0
			86	86		
5	B	60	Total	O	0	1
			61	61		
5	C	72	Total	O	0	0
			72	72		
5	D	67	Total	O	0	0
			67	67		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: NAD kinase 1




• Molecule 1: NAD kinase 1




• Molecule 1: NAD kinase 1



• Molecule 1: NAD kinase 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.93Å 119.53Å 67.31Å 90.00° 100.28° 90.00°	Depositor
Resolution (Å)	44.26 – 2.27 44.26 – 2.27	Depositor EDS
% Data completeness (in resolution range)	96.1 (44.26-2.27) 96.1 (44.26-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 2.27Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.217 , 0.259 0.216 , 0.260	Depositor DCC
R_{free} test set	1408 reflections (3.05%)	DCC
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8624	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5AQ, GOL, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/2114	0.38	0/2862
1	B	0.23	0/2168	0.37	0/2929
1	C	0.22	0/2085	0.36	0/2824
1	D	0.22	0/2004	0.36	0/2712
All	All	0.22	0/8371	0.37	0/11327

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2064	0	1982	10	0
1	B	2113	0	2063	16	0
1	C	2035	0	1937	13	0
1	D	1953	0	1884	20	0
2	A	13	0	5	2	0
2	B	13	0	5	3	0
2	D	13	0	5	0	0
3	A	6	0	8	0	0
4	A	32	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	32	0	0	1	0
4	C	32	0	0	0	0
4	D	32	0	0	3	0
5	A	86	0	0	2	0
5	B	61	0	0	1	0
5	C	72	0	0	0	0
5	D	67	0	0	2	0
All	All	8624	0	7889	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:252:PRO:HA	2:B:301:CIT:H42	1.76	0.67
1:A:38:ILE:HD13	1:A:90:LEU:HD22	1.79	0.64
1:B:125:THR:HG1	1:B:220:SER:HG	1.51	0.59
1:D:160:THR:HG21	1:D:172:MET:HG2	1.84	0.58
1:C:255:ARG:HD2	1:D:142:ILE:HD13	1.86	0.57
4:D:302:5AQ:O5'	4:D:302:5AQ:CBC	2.53	0.57
1:A:100:TYR:OH	2:A:301:CIT:O5	2.18	0.55
1:C:38:ILE:HG21	1:C:90:LEU:HD21	1.89	0.55
1:C:38:ILE:HD13	1:C:90:LEU:HD11	1.87	0.55
1:D:2:LYS:N	5:D:403:HOH:O	2.39	0.55
1:C:106:THR:HG21	1:C:235:ARG:HH11	1.72	0.54
1:B:222:ASP:O	1:B:223[A]:HIS:HB2	2.07	0.54
2:B:301:CIT:O3	5:B:401:HOH:O	2.18	0.54
1:B:182:THR:HG22	1:B:198:PRO:HB3	1.92	0.52
1:B:148:ARG:HG2	1:B:187:ILE:HD13	1.92	0.51
1:C:157:PRO:HB2	1:C:174:PRO:HA	1.92	0.50
1:A:49:LEU:HD23	1:A:121:LEU:HD23	1.94	0.50
1:D:138:VAL:HB	1:D:208:SER:HB3	1.93	0.50
1:C:90:LEU:HD13	1:C:248:PHE:HZ	1.77	0.50
1:B:1:MET:N	1:B:1:MET:SD	2.71	0.49
1:C:255:ARG:HH11	1:D:142:ILE:HD13	1.78	0.49
4:D:302:5AQ:NAW	4:D:302:5AQ:C8	2.76	0.49
1:C:252:PRO:HD2	1:C:255:ARG:HD3	1.94	0.49
1:D:146:ARG:HG2	1:D:192:TYR:HD1	1.78	0.48
1:B:252:PRO:HD2	1:B:255:ARG:HD3	1.94	0.48
1:B:95:TYR:HB3	1:B:248:PHE:CE1	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:226:ILE:HG13	1:C:228:HIS:CE1	2.49	0.47
1:B:128:SER:HB2	1:B:133:PHE:HB2	1.97	0.47
2:B:301:CIT:O1	2:B:301:CIT:O7	2.28	0.47
1:B:89:LEU:HB3	1:B:248:PHE:CE1	2.50	0.46
1:D:211:PRO:HG3	1:D:217:PHE:HE2	1.80	0.46
1:D:256:ARG:NH2	5:D:407:HOH:O	2.48	0.46
1:B:148:ARG:HD3	1:B:192:TYR:CE2	2.51	0.46
1:B:160:THR:HA	1:B:164:ASN:HB3	1.98	0.46
1:D:68:HIS:ND1	1:D:68:HIS:O	2.49	0.45
2:A:301:CIT:O3	5:A:401:HOH:O	2.21	0.45
1:D:135:VAL:HG21	1:D:209:LEU:HB3	1.97	0.45
1:D:134:VAL:HG22	1:D:148:ARG:HG3	1.98	0.44
1:B:210:GLN:HA	1:B:211:PRO:HD3	1.82	0.44
1:D:252:PRO:HD2	1:D:255:ARG:HE	1.83	0.44
1:A:133:PHE:HB3	1:A:149:GLY:O	2.18	0.43
1:D:99:SER:HB3	1:D:242[B]:LYS:HD3	2.00	0.43
1:A:144:PHE:O	1:B:258:HIS:HD2	2.00	0.43
1:A:146:ARG:HG2	1:A:192:TYR:HD1	1.83	0.43
1:D:123:GLU:OE2	4:D:302:5AQ:O2'	2.37	0.43
1:D:127:LYS:O	1:D:218:GLN:N	2.48	0.43
1:A:173:HIS:HA	1:A:174:PRO:HD3	1.80	0.42
1:A:256:ARG:NH2	5:A:402:HOH:O	2.31	0.42
1:A:65:ILE:HG21	1:A:86:LEU:HD21	2.01	0.42
1:D:21:ILE:HD11	1:D:31:TYR:CD1	2.54	0.42
1:B:49:LEU:HD23	1:B:121:LEU:HD23	2.02	0.42
1:C:133:PHE:HB3	1:C:149:GLY:O	2.20	0.42
1:C:40:ILE:HG12	1:C:65:ILE:HB	2.01	0.42
1:D:220:SER:HA	1:D:225:SER:HA	2.02	0.42
1:D:65:ILE:HG21	1:D:86:LEU:HD21	2.03	0.41
1:D:66:GLY:HA3	1:D:76:ALA:HA	2.03	0.41
1:D:133:PHE:HB3	1:D:149:GLY:O	2.21	0.40
1:C:5:ILE:HG12	1:C:40:ILE:HB	2.03	0.40
1:A:119:LEU:HD11	1:A:241:LYS:HD2	2.02	0.40
4:B:302:5AQ:NAW	4:B:302:5AQ:CBA	2.82	0.40
1:C:119:LEU:HD11	1:C:241:LYS:HD2	2.02	0.40
1:B:108:LYS:HE3	1:B:233:GLU:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/272 (95%)	248 (96%)	11 (4%)	0	100	100
1	B	261/272 (96%)	248 (95%)	11 (4%)	2 (1%)	24	26
1	C	257/272 (94%)	244 (95%)	13 (5%)	0	100	100
1	D	245/272 (90%)	233 (95%)	12 (5%)	0	100	100
All	All	1022/1088 (94%)	973 (95%)	47 (5%)	2 (0%)	56	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	223[A]	HIS
1	B	223[B]	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/237 (92%)	214 (98%)	5 (2%)	58	73
1	B	229/237 (97%)	220 (96%)	9 (4%)	39	52
1	C	213/237 (90%)	210 (99%)	3 (1%)	74	86
1	D	206/237 (87%)	202 (98%)	4 (2%)	65	79
All	All	867/948 (92%)	846 (98%)	21 (2%)	57	72

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	LEU
1	A	123	GLU
1	A	163	TYR
1	A	251	PHE
1	A	263	GLU
1	B	1	MET
1	B	11	GLU
1	B	15	LEU
1	B	94	GLU
1	B	123	GLU
1	B	163	TYR
1	B	224	LEU
1	B	249	ARG
1	B	251	PHE
1	C	123	GLU
1	C	163	TYR
1	C	251	PHE
1	D	3	TYR
1	D	163	TYR
1	D	223	HIS
1	D	251	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	228	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CIT	A	301	-	3,12,12	1.21	0	3,17,17	1.41	1 (33%)
3	GOL	A	302	-	5,5,5	0.36	0	5,5,5	0.18	0
4	5AQ	A	303	-	31,35,35	1.45	6 (19%)	29,49,49	2.59	5 (17%)
2	CIT	B	301	-	3,12,12	1.39	0	3,17,17	2.37	2 (66%)
4	5AQ	B	302	-	31,35,35	1.61	6 (19%)	29,49,49	2.71	5 (17%)
4	5AQ	C	301	-	31,35,35	1.23	3 (9%)	29,49,49	2.61	3 (10%)
2	CIT	D	301	-	3,12,12	1.33	0	3,17,17	2.69	2 (66%)
4	5AQ	D	302	-	31,35,35	1.50	4 (12%)	29,49,49	2.71	7 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	301	-	-	0/6/16/16	0/0/0/0
3	GOL	A	302	-	-	0/4/4/4	0/0/0/0
4	5AQ	A	303	-	-	0/11/33/33	0/4/4/4
2	CIT	B	301	-	-	0/6/16/16	0/0/0/0
4	5AQ	B	302	-	-	0/11/33/33	0/4/4/4
4	5AQ	C	301	-	-	0/11/33/33	0/4/4/4
2	CIT	D	301	-	-	0/6/16/16	0/0/0/0
4	5AQ	D	302	-	-	0/11/33/33	0/4/4/4

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	302	5AQ	C8-N9	-5.27	1.30	1.36
4	A	303	5AQ	C8-N9	-4.48	1.31	1.36
4	D	302	5AQ	C8-N9	-4.27	1.32	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	301	5AQ	C8-N9	-3.70	1.32	1.36
4	D	302	5AQ	O4'-C4'	-3.45	1.37	1.45
4	B	302	5AQ	O4'-C4'	-3.44	1.37	1.45
4	B	302	5AQ	C4-N3	-3.06	1.31	1.35
4	D	302	5AQ	C8-SAT	-3.00	1.67	1.75
4	A	303	5AQ	C8-SAT	-2.76	1.68	1.75
4	A	303	5AQ	O4'-C4'	-2.69	1.38	1.45
4	D	302	5AQ	O4'-C1'	-2.61	1.37	1.41
4	C	301	5AQ	O4'-C4'	-2.55	1.39	1.45
4	C	301	5AQ	C4-N3	-2.52	1.31	1.35
4	A	303	5AQ	C4-N3	-2.48	1.31	1.35
4	B	302	5AQ	C8-SAT	-2.32	1.69	1.75
4	A	303	5AQ	O2'-C2'	-2.28	1.37	1.43
4	B	302	5AQ	CAU-CAV	-2.23	1.49	1.51
4	B	302	5AQ	C2'-C1'	-2.12	1.50	1.53
4	A	303	5AQ	C2-N1	2.08	1.37	1.33

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	301	5AQ	N3-C2-N1	-11.21	120.07	128.87
4	A	303	5AQ	N3-C2-N1	-11.08	120.17	128.87
4	B	302	5AQ	N3-C2-N1	-11.01	120.22	128.87
4	D	302	5AQ	N3-C2-N1	-10.12	120.92	128.87
4	D	302	5AQ	C4'-O4'-C1'	-7.39	101.81	109.64
4	B	302	5AQ	C4'-O4'-C1'	-7.28	101.93	109.64
4	C	301	5AQ	C1'-N9-C4	-5.37	121.19	126.92
4	C	301	5AQ	C4'-O4'-C1'	-5.13	104.20	109.64
4	A	303	5AQ	C1'-N9-C4	-4.29	122.34	126.92
4	A	303	5AQ	C5'-C4'-C3'	-4.22	105.02	115.08
4	D	302	5AQ	C1'-N9-C4	-4.13	122.50	126.92
4	A	303	5AQ	C4'-O4'-C1'	-3.83	105.58	109.64
2	D	301	CIT	C3-C4-C5	-3.57	109.37	114.95
4	B	302	5AQ	C1'-N9-C4	-3.45	123.23	126.92
2	B	301	CIT	C3-C2-C1	-3.28	109.82	114.95
2	D	301	CIT	C3-C2-C1	-2.96	110.33	114.95
4	B	302	5AQ	O4'-C4'-C3'	-2.83	99.43	105.16
2	A	301	CIT	C3-C4-C5	-2.40	111.20	114.95
2	B	301	CIT	C3-C4-C5	-2.39	111.22	114.95
4	B	302	5AQ	O4'-C4'-C5'	-2.34	104.22	109.18
4	A	303	5AQ	O4'-C4'-C5'	-2.31	104.28	109.18
4	D	302	5AQ	O4'-C4'-C3'	-2.25	100.59	105.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	302	5AQ	N6-C6-N1	2.04	121.94	118.52
4	D	302	5AQ	OBF-CAV-NAW	2.63	128.18	122.96
4	D	302	5AQ	CAX-NAW-CAV	3.02	128.78	122.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	CIT	2	0
2	B	301	CIT	3	0
4	B	302	5AQ	1	0
4	D	302	5AQ	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/272 (96%)	0.35	7 (2%) 58 66	25, 45, 77, 113	40 (15%)
1	B	263/272 (96%)	0.59	16 (6%) 25 31	28, 54, 87, 104	46 (17%)
1	C	261/272 (95%)	0.78	28 (10%) 8 11	27, 58, 103, 129	35 (13%)
1	D	250/272 (91%)	0.97	41 (16%) 2 3	30, 64, 117, 135	34 (13%)
All	All	1037/1088 (95%)	0.67	92 (8%) 12 16	25, 54, 103, 135	155 (14%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	86	LEU	5.4
1	D	89	LEU	5.4
1	D	15	LEU	5.4
1	D	87	VAL	4.9
1	C	3	TYR	4.8
1	C	40	ILE	4.6
1	D	83	ALA	4.6
1	C	31	TYR	4.6
1	B	91	ALA	4.3
1	D	95	TYR	4.2
1	C	90	LEU	4.1
1	D	248	PHE	4.1
1	D	34	VAL	4.0
1	D	88	LYS	4.0
1	D	227	LEU	3.9
1	D	18	LEU	3.9
1	D	91	ALA	3.9
1	B	267	HIS	3.9
1	D	247	ARG	3.8
1	D	119	LEU	3.8
1	D	231	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	223[A]	HIS	3.6
1	D	117	THR	3.6
1	D	84	ASP	3.5
1	C	18	LEU	3.5
1	D	250	SER	3.5
1	C	212	VAL	3.4
1	D	16	LEU	3.4
1	C	60	ASP	3.3
1	C	30	GLU	3.2
1	B	226	ILE	3.1
1	C	191	VAL	3.1
1	B	188	ASN	3.0
1	C	187	ILE	3.0
1	D	217	PHE	3.0
1	B	109	TYR	2.9
1	D	7	SER	2.9
1	D	81	ALA	2.9
1	B	92	LYS	2.8
1	D	10	ASP	2.8
1	D	42	ILE	2.8
1	D	214	ASP	2.8
1	D	85	LYS	2.8
1	D	129	SER	2.8
1	D	80	PRO	2.7
1	D	71	HIS	2.7
1	C	61	GLU	2.5
1	D	32	ASP	2.5
1	C	91	ALA	2.4
1	C	248	PHE	2.4
1	D	61	GLU	2.4
1	B	217	PHE	2.4
1	A	250	SER	2.4
1	A	196	GLY	2.4
1	C	94	GLU	2.4
1	B	33	ASP	2.4
1	B	95	TYR	2.4
1	A	72	LEU	2.4
1	D	216	ASP	2.3
1	B	3	TYR	2.3
1	C	198	PRO	2.3
1	D	79	ARG	2.3
1	D	230	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	44	GLY	2.3
1	A	69	THR	2.3
1	C	1	MET	2.3
1	C	32	ASP	2.3
1	C	78	TRP	2.3
1	D	210	GLN	2.3
1	B	19	ASN	2.2
1	D	70	GLY	2.2
1	B	232	GLN	2.2
1	C	39	VAL	2.2
1	A	248	PHE	2.2
1	C	38	ILE	2.2
1	A	232	GLN	2.2
1	A	94	GLU	2.2
1	C	167	LEU	2.2
1	C	11	GLU	2.1
1	C	21	ILE	2.1
1	C	2	LYS	2.1
1	C	19	ASN	2.1
1	D	105	THR	2.1
1	C	24	PHE	2.1
1	B	212	VAL	2.1
1	D	138	VAL	2.1
1	C	29	MET	2.1
1	D	31	TYR	2.1
1	D	226	ILE	2.0
1	D	234	ILE	2.0
1	B	229	ARG	2.0
1	B	167	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	302	6/6	0.79	0.31	3.50	72,73,73,73	0
4	5AQ	D	302	32/32	0.68	0.28	2.99	67,71,110,111	0
2	CIT	B	301	13/13	0.85	0.22	1.79	77,78,79,80	0
2	CIT	D	301	13/13	0.83	0.36	1.65	51,52,54,55	13
2	CIT	A	301	13/13	0.85	0.21	0.93	60,64,67,68	13
4	5AQ	B	302	32/32	0.93	0.12	-0.62	37,43,63,63	0
4	5AQ	C	301	32/32	0.91	0.13	-0.77	37,41,66,66	0
4	5AQ	A	303	32/32	0.93	0.13	-1.06	29,43,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.