



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:48 PM GMT

PDB ID : 4DIF
Title : Structure of A1-type ketoreductase
Authors : Zheng, J.; Keatinge-Clay, A.T.
Deposited on : 2012-01-30
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2658 (1.54-1.50)
Clashscore	102246	2887 (1.54-1.50)
Ramachandran outliers	100387	2818 (1.54-1.50)
Sidechain outliers	100360	2816 (1.54-1.50)
RSRZ outliers	91569	2660 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	<div> <div>4%</div> <div>88%</div> <div>8%</div> <div>.</div> </div>
1	B	496	<div> <div>3%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AmphB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	476	Total	C	N	O	S	0	0	0
			3469	2166	636	658	9			
1	B	476	Total	C	N	O	S	0	0	0
			3469	2166	636	658	9			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	EXPRESSION TAG	UNP Q93NW7
A	-19	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-18	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-17	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-16	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-15	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-14	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-13	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-12	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-11	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	-10	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-9	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-8	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-7	LEU	-	EXPRESSION TAG	UNP Q93NW7
A	-6	VAL	-	EXPRESSION TAG	UNP Q93NW7
A	-5	PRO	-	EXPRESSION TAG	UNP Q93NW7
A	-4	ARG	-	EXPRESSION TAG	UNP Q93NW7
A	-3	GLY	-	EXPRESSION TAG	UNP Q93NW7
A	-2	SER	-	EXPRESSION TAG	UNP Q93NW7
A	-1	HIS	-	EXPRESSION TAG	UNP Q93NW7
A	0	MET	-	EXPRESSION TAG	UNP Q93NW7
A	355	THR	GLY	ENGINEERED MUTATION	UNP Q93NW7
A	364	HIS	GLN	ENGINEERED MUTATION	UNP Q93NW7
B	-20	MET	-	EXPRESSION TAG	UNP Q93NW7
B	-19	GLY	-	EXPRESSION TAG	UNP Q93NW7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-17	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-16	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-15	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-14	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-13	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-12	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-11	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	-10	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-9	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-8	GLY	-	EXPRESSION TAG	UNP Q93NW7
B	-7	LEU	-	EXPRESSION TAG	UNP Q93NW7
B	-6	VAL	-	EXPRESSION TAG	UNP Q93NW7
B	-5	PRO	-	EXPRESSION TAG	UNP Q93NW7
B	-4	ARG	-	EXPRESSION TAG	UNP Q93NW7
B	-3	GLY	-	EXPRESSION TAG	UNP Q93NW7
B	-2	SER	-	EXPRESSION TAG	UNP Q93NW7
B	-1	HIS	-	EXPRESSION TAG	UNP Q93NW7
B	0	MET	-	EXPRESSION TAG	UNP Q93NW7
B	355	THR	GLY	ENGINEERED MUTATION	UNP Q93NW7
B	364	HIS	GLN	ENGINEERED MUTATION	UNP Q93NW7

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	
			48	21	7	17	3	
2	B	1	Total	C	N	O	P	
			48	21	7	17	3	

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O		
			6	3	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	309	Total	O		
			309	309	0	0
4	B	304	Total	O		
			304	304	0	0

- Molecule 1: AmphB



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.55Å 63.72Å 71.98Å 72.92° 67.22° 89.74°	Depositor
Resolution (Å)	50.00 – 1.52 26.61 – 1.52	Depositor EDS
% Data completeness (in resolution range)	86.9 (50.00-1.52) 76.9 (26.61-1.52)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.98 (at 1.52Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.198 , 0.224 0.196 , 0.221	Depositor DCC
R_{free} test set	7355 reflections (6.06%)	DCC
Wilson B-factor (Å ²)	12.7	Xtriage
Anisotropy	0.043	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 52.8	EDS
Estimated twinning fraction	0.064 for h,-k,h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 127760 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7653	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/3540 (0.0%)	0.50	0/4837
1	B	0.53	0/3540	0.51	0/4837
All	All	0.53	1/7080 (0.0%)	0.51	0/9674

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	396	TRP	NE1-CE2	-5.98	1.29	1.37

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	ARG	Sidechain
1	A	401	MET	Mainchain
1	B	124	ARG	Sidechain
1	B	405	PRO	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3469	0	3429	39	0
1	B	3469	0	3429	49	0
2	A	48	0	26	1	0
2	B	48	0	26	1	0
3	B	6	0	8	1	0
4	A	309	0	0	5	0
4	B	304	0	0	6	0
All	All	7653	0	6918	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:407:VAL:O	1:B:411:LEU:HD13	1.55	1.07
1:A:124:ARG:HE	1:A:166:GLU:HA	1.15	1.04
1:A:90:HIS:HD2	1:A:92:ASP:H	1.08	0.98
1:A:124:ARG:NE	1:A:166:GLU:HA	1.79	0.97
1:B:19:ARG:NH2	1:B:45:LEU:O	1.99	0.95
1:A:341:THR:HA	1:A:344:LEU:HD13	1.48	0.95
1:B:208:VAL:HG23	1:B:211:SER:OG	1.69	0.92
1:B:41:VAL:CG2	1:B:164:LEU:HD11	2.01	0.88
1:B:411:LEU:HD12	1:B:411:LEU:N	1.96	0.79
1:B:407:VAL:O	1:B:411:LEU:CD1	2.29	0.78
1:B:213:LYS:HE3	1:B:435:ASP:CB	2.15	0.77
1:B:59:ARG:HB2	1:B:105:LEU:CD1	2.15	0.76
1:B:59:ARG:HB2	1:B:105:LEU:HD12	1.68	0.76
1:B:404:ASP:C	1:B:406:GLU:H	1.87	0.75
1:A:139:VAL:HG22	4:A:859:HOH:O	1.87	0.74
1:A:90:HIS:CD2	1:A:92:ASP:H	2.01	0.70
1:A:203:HIS:HD2	4:A:740:HOH:O	1.75	0.69
1:B:213:LYS:HE3	1:B:435:ASP:HB2	1.72	0.69
1:B:139:VAL:HG22	4:B:685:HOH:O	1.91	0.69
1:A:203:HIS:HE1	1:A:467:GLU:OE2	1.76	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:411:LEU:CD1	1:B:411:LEU:N	2.57	0.67
1:A:364:HIS:HD2	4:A:685:HOH:O	1.78	0.67
1:B:245:HIS:HE1	1:B:274:THR:OG1	1.80	0.64
1:A:166:GLU:H	1:A:166:GLU:CD	1.98	0.64
1:A:402:ALA:O	1:A:408:HIS:HB2	1.98	0.63
1:A:245:HIS:HE1	1:A:274:THR:OG1	1.81	0.62
1:A:341:THR:CA	1:A:344:LEU:HD13	2.26	0.61
1:B:23:ARG:NH1	1:B:73:GLU:OE1	2.34	0.61
1:B:213:LYS:HE3	1:B:435:ASP:HB3	1.83	0.60
1:B:41:VAL:HG21	1:B:164:LEU:HD11	1.81	0.60
1:B:87:GLU:OE1	1:B:139:VAL:HG21	2.02	0.60
1:A:124:ARG:CZ	1:A:166:GLU:HA	2.32	0.59
1:B:402:ALA:O	1:B:408:HIS:HB2	2.03	0.59
1:B:41:VAL:CG2	1:B:164:LEU:CD1	2.79	0.58
1:B:124:ARG:HE	1:B:166:GLU:HA	1.66	0.58
1:B:208:VAL:HG23	1:B:211:SER:HG	1.68	0.58
1:B:411:LEU:CD1	1:B:411:LEU:H	2.17	0.57
1:A:41:VAL:CG2	1:A:164:LEU:HD11	2.35	0.57
1:B:364:HIS:HD2	4:B:661:HOH:O	1.88	0.57
1:A:77:HIS:HE1	1:A:180:LEU:O	1.88	0.57
1:A:391:VAL:HG21	1:A:428:LEU:HD13	1.87	0.57
1:B:391:VAL:HG21	1:B:428:LEU:HD13	1.87	0.56
1:B:380:HIS:HE1	4:B:789:HOH:O	1.87	0.56
1:A:41:VAL:HG21	1:A:164:LEU:HD11	1.87	0.56
1:B:404:ASP:C	1:B:406:GLU:N	2.59	0.55
1:B:87:GLU:OE1	1:B:139:VAL:CG2	2.55	0.54
1:B:404:ASP:O	1:B:406:GLU:N	2.39	0.53
1:B:37:TRP:HE3	1:B:164:LEU:HD12	1.74	0.53
1:B:237:ARG:HD2	1:B:425:LEU:HD13	1.91	0.53
1:B:77:HIS:HE1	1:B:180:LEU:O	1.92	0.53
1:B:278:CYS:SG	4:B:866:HOH:O	2.53	0.52
1:A:344:LEU:HD12	1:A:344:LEU:N	2.25	0.52
1:B:237:ARG:CD	1:B:425:LEU:HD13	2.40	0.51
1:A:450:ALA:HB3	1:A:451:PRO:HD3	1.93	0.51
1:A:335:ARG:HD2	4:A:778:HOH:O	2.09	0.51
1:A:224:THR:OG1	1:A:304:HIS:CD2	2.64	0.50
1:B:135:LEU:O	1:B:380:HIS:HD2	1.93	0.50
1:A:124:ARG:NH1	4:A:735:HOH:O	2.45	0.50
1:B:70:ARG:HH11	1:B:70:ARG:HG3	1.77	0.50
1:B:37:TRP:CE3	1:B:164:LEU:HD12	2.47	0.49
1:A:135:LEU:O	1:A:380:HIS:HD2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:THR:OG1	1:A:90:HIS:HE1	1.96	0.48
1:A:224:THR:OG1	1:A:304:HIS:HD2	1.97	0.47
1:B:125:GLY:HA2	4:B:733:HOH:O	2.15	0.47
1:B:208:VAL:HG23	1:B:211:SER:CB	2.44	0.47
1:B:41:VAL:HG22	1:B:164:LEU:HD11	1.91	0.46
1:B:164:LEU:HD13	1:B:168:VAL:HG21	1.98	0.45
1:A:41:VAL:CG2	1:A:164:LEU:CD1	2.95	0.45
1:B:429:ASP:OD1	3:B:502:GOL:H12	2.17	0.44
1:A:330:LYS:HD3	1:A:374:LEU:HD11	1.98	0.44
1:A:341:THR:HA	1:A:344:LEU:CD1	2.33	0.44
1:A:220:SER:OG	1:A:245:HIS:HD2	2.02	0.42
1:B:404:ASP:OD1	1:B:406:GLU:HB3	2.20	0.42
1:B:19:ARG:HG2	1:B:177:THR:HB	2.01	0.42
1:A:393:TRP:HB2	2:A:501:NDP:C5N	2.49	0.42
1:A:124:ARG:NE	1:A:166:GLU:CA	2.68	0.42
1:A:37:TRP:CE3	1:A:164:LEU:HD12	2.54	0.42
1:A:364:HIS:N	1:A:365:PRO:CD	2.83	0.42
1:B:292:GLU:HG2	4:B:811:HOH:O	2.19	0.42
1:A:304:HIS:CE1	1:A:329:ALA:O	2.73	0.41
1:A:344:LEU:CD1	1:A:344:LEU:N	2.84	0.41
1:A:411:LEU:O	1:A:414:GLN:HG2	2.20	0.41
1:A:89:LEU:HD11	1:A:335:ARG:HH22	1.86	0.41
1:B:450:ALA:HB3	1:B:451:PRO:HD3	2.03	0.41
1:B:393:TRP:HB2	2:B:501:NDP:C5N	2.50	0.41
1:B:213:LYS:CE	1:B:435:ASP:HB3	2.50	0.40
1:A:304:HIS:HE1	1:A:329:ALA:O	2.04	0.40
1:B:41:VAL:HG11	1:B:82:LEU:HD21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/496 (96%)	464 (98%)	10 (2%)	0	100	100
1	B	474/496 (96%)	464 (98%)	9 (2%)	1 (0%)	52	25
All	All	948/992 (96%)	928 (98%)	19 (2%)	1 (0%)	56	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	405	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/352 (95%)	334 (100%)	1 (0%)	94	86
1	B	335/352 (95%)	335 (100%)	0	100	100
All	All	670/704 (95%)	669 (100%)	1 (0%)	95	87

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	311	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	10	ASN
1	A	77	HIS
1	A	90	HIS
1	A	102	GLN
1	A	203	HIS
1	A	245	HIS
1	A	304	HIS
1	A	336	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	364	HIS
1	A	380	HIS
1	B	32	HIS
1	B	77	HIS
1	B	102	GLN
1	B	241	GLN
1	B	245	HIS
1	B	336	HIS
1	B	364	HIS
1	B	380	HIS
1	B	423	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	501	-	42,52,52	1.15	3 (7%)	55,80,80	1.91	5 (9%)
2	NDP	B	501	-	42,52,52	1.07	2 (4%)	55,80,80	1.94	7 (12%)
3	GOL	B	502	-	5,5,5	0.20	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	501	-	-	0/30/77/77	0/5/5/5
2	NDP	B	501	-	-	0/30/77/77	0/5/5/5
3	GOL	B	502	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NDP	C4N-C5N	-4.89	1.38	1.49
2	B	501	NDP	C4N-C5N	-4.65	1.39	1.49
2	A	501	NDP	C2N-C3N	2.00	1.39	1.34
2	B	501	NDP	C6N-C5N	2.57	1.38	1.33
2	A	501	NDP	C6N-C5N	2.98	1.39	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NDP	N3A-C2A-N1A	-11.43	120.14	128.89
2	B	501	NDP	N3A-C2A-N1A	-11.18	120.34	128.89
2	B	501	NDP	C4A-C5A-N7A	-2.81	106.90	109.48
2	B	501	NDP	O2B-P2B-O1X	-2.47	100.95	107.11
2	B	501	NDP	PN-O3-PA	-2.33	126.18	132.73
2	A	501	NDP	C3N-C2N-N1N	-2.27	119.89	123.14
2	A	501	NDP	C4A-C5A-N7A	-2.13	107.52	109.48
2	A	501	NDP	O2B-P2B-O1X	-2.11	101.84	107.11
2	B	501	NDP	C1B-N9A-C4A	-2.10	123.77	126.94
2	B	501	NDP	O3X-P2B-O2X	2.16	115.61	107.38
2	A	501	NDP	C5N-C4N-C3N	2.20	118.58	112.52
2	B	501	NDP	C5N-C4N-C3N	2.27	118.78	112.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NDP	1	0
2	B	501	NDP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/496 (95%)	0.30	20 (4%) 40 41	8, 14, 32, 79	0
1	B	476/496 (95%)	0.31	17 (3%) 46 49	8, 14, 32, 66	0
All	All	952/992 (95%)	0.30	37 (3%) 43 45	8, 14, 32, 79	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	405	PRO	7.5
1	A	407	VAL	7.3
1	A	406	GLU	5.5
1	A	412	VAL	5.0
1	A	411	LEU	4.9
1	A	410	ARG	4.6
1	A	405	PRO	4.5
1	A	208	VAL	4.4
1	A	413	ARG	4.1
1	B	411	LEU	3.7
1	A	131	ARG	3.6
1	A	21	ALA	3.5
1	B	413	ARG	3.3
1	B	131	ARG	3.1
1	A	213	LYS	3.0
1	B	410	ARG	2.9
1	A	212	GLY	2.9
1	B	406	GLU	2.8
1	A	295	GLU	2.8
1	B	407	VAL	2.7
1	A	209	PRO	2.7
1	B	208	VAL	2.7
1	A	404	ASP	2.6
1	A	-1	HIS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	302	VAL	2.4
1	B	351	LEU	2.4
1	A	409	ASP	2.4
1	A	408	HIS	2.3
1	A	342	ALA	2.3
1	A	352	PHE	2.3
1	B	132	GLY	2.3
1	B	260	ALA	2.2
1	B	71	GLU	2.2
1	B	-1	HIS	2.1
1	B	473	SER	2.1
1	B	412	VAL	2.1
1	B	21	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	502	6/6	0.92	0.10	0.54	21,21,23,23	0
2	NDP	A	501	48/48	0.98	0.07	-1.07	11,12,15,18	0
2	NDP	B	501	48/48	0.98	0.07	-1.15	10,11,13,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.