



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:48 PM GMT

PDB ID : 4DJH
Title : Structure of the human kappa opioid receptor in complex with JDTic
Authors : Wu, H.; Wacker, D.; Katritch, V.; Mileni, M.; Han, G.W.; Vardy, E.; Liu, W.; Thompson, A.A.; Huang, X.P.; Carroll, F.I.; Mascarella, S.W.; Westkaemper, R.B.; Mosier, P.D.; Roth, B.L.; Cherezov, V.; Stevens, R.C.; GPCR Network (GPCR)
Deposited on : 2012-02-01
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CIT	A	1301	-	-	-	X
4	OLC	B	1302	-	-	-	X
5	PEG	B	1305	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7089 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Kappa-type opioid receptor, Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3480	2277	564	616	23			
1	B	448	Total	C	N	O	S	0	0	0
			3452	2263	555	611	23			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	GLY	-	EXPRESSION TAG	UNP P41145
A	39	GLY	-	EXPRESSION TAG	UNP P41145
A	40	THR	-	EXPRESSION TAG	UNP P41145
A	41	THR	-	EXPRESSION TAG	UNP P41145
A	42	MET	-	EXPRESSION TAG	UNP P41145
A	135	LEU	ILE	engineered mutation	UNP P41145
A	1054	THR	CYS	engineered mutation	UNP P00720
A	1097	ALA	CYS	engineered mutation	UNP P00720
B	38	GLY	-	EXPRESSION TAG	UNP P41145
B	39	GLY	-	EXPRESSION TAG	UNP P41145
B	40	THR	-	EXPRESSION TAG	UNP P41145
B	41	THR	-	EXPRESSION TAG	UNP P41145
B	42	MET	-	EXPRESSION TAG	UNP P41145
B	135	LEU	ILE	engineered mutation	UNP P41145
B	1054	THR	CYS	engineered mutation	UNP P00720
B	1097	ALA	CYS	engineered mutation	UNP P00720

- Molecule 2 is (3R)-7-HYDROXY-N-[(2S)-1-[(3R,4R)-4-(3-HYDROXYPHENYL)-3,4-DIMETHYLPIPERIDIN-1-YL]-3-METHYLBUTAN-2-YL]-1,2,3,4-TETRAHYDROISOQUINOLINE-3-CARBOXAMIDE (three-letter code: JDC) (formula: C₂₈H₃₉N₃O₃).




-
- Chemical structure of Citric acid (CIT) is shown. The structure features a central carbon atom (C3) bonded to three hydroxyl groups (OH) and a carboxylate group. The carboxylate group consists of a carbon atom (C6) double-bonded to an oxygen atom (O5) and single-bonded to a hydroxyl group (OH, O6). The central carbon (C3) is also bonded to a carboxylate group (C1, O2, O1) and a carboxylate group (C5, O3, O4). The structure is labeled with atom names (C1, C2, C3, C4, C5, C6) and atom numbers (O1, O2, O3, O4, O5, O6, O7).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		


- 

code: OLC) (formula: C₂₁H₄₀O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			16	12	4		
4	B	1	Total	C	O	0	0
			21	17	4		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	11	Total	O	0	0
			11	11		
6	B	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Kappa-type opioid receptor, Lysozyme



- Molecule 1: Kappa-type opioid receptor, Lysozyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.90 Å 147.30 Å 205.29 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.86 – 2.90 35.86 – 2.86	Depositor EDS
% Data completeness (in resolution range)	97.8 (35.86-2.90) 96.5 (35.86-2.86)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.85 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, R_{free}	0.227 , 0.265 0.227 , 0.264	Depositor DCC
R_{free} test set	1854 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	75.2	Xtriage
Anisotropy	0.213	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 57.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 38260 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7089	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, OLC, JDC, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3555	0.42	0/4843
1	B	0.30	0/3528	0.45	0/4818
All	All	0.29	0/7083	0.44	0/9661

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3480	0	3491	49	0
1	B	3452	0	3430	45	0
2	A	34	0	38	0	0
2	B	34	0	38	2	0
3	A	13	0	5	0	0
4	B	37	0	50	2	0
5	B	14	0	20	0	0
6	A	11	0	0	0	0
6	B	14	0	0	0	0
All	All	7089	0	7072	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:TRP:CH2	1:B:134:VAL:HG11	2.23	0.73
1:A:1011:GLU:HG2	1:A:1030:GLY:HA3	1.76	0.67
1:B:138:ASP:OD1	2:B:1300:JDC:N2	2.31	0.64
1:A:122:ASN:ND2	1:A:122:ASN:O	2.32	0.63
1:A:161:CYS:SG	1:A:252:ARG:NH2	2.74	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/480 (92%)	427 (96%)	16 (4%)	0	100	100
1	B	444/480 (92%)	434 (98%)	10 (2%)	0	100	100
All	All	887/960 (92%)	861 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/422 (88%)	353 (95%)	17 (5%)	33	69
1	B	363/422 (86%)	353 (97%)	10 (3%)	51	84
All	All	733/844 (87%)	706 (96%)	27 (4%)	41	77

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1115	THR
1	A	340	CYS
1	B	1125	ARG
1	A	1127	ASP
1	A	168	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	JDC	A	1300	-	36,37,37	2.61	13 (36%)	41,54,54	1.55	7 (17%)
3	CIT	A	1301	-	3,12,12	1.31	0	3,17,17	2.50	2 (66%)
2	JDC	B	1300	-	36,37,37	2.59	13 (36%)	41,54,54	1.63	9 (21%)
4	OLC	B	1302	-	15,15,24	1.10	1 (6%)	16,16,25	1.14	1 (6%)
4	OLC	B	1303	-	20,20,24	1.03	1 (5%)	20,21,25	0.95	1 (5%)
5	PEG	B	1304	-	6,6,6	0.60	0	5,5,5	0.74	0
5	PEG	B	1305	-	6,6,6	0.72	0	5,5,5	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JDC	A	1300	-	-	0/22/47/47	0/4/4/4
3	CIT	A	1301	-	-	0/6/16/16	0/0/0/0
2	JDC	B	1300	-	-	0/22/47/47	0/4/4/4
4	OLC	B	1302	-	-	0/15/15/24	0/0/0/0
4	OLC	B	1303	-	-	0/20/20/24	0/0/0/0
5	PEG	B	1304	-	-	0/4/4/4	0/0/0/0
5	PEG	B	1305	-	-	0/4/4/4	0/0/0/0

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1300	JDC	C41-C31	-7.19	1.46	1.55
2	A	1300	JDC	C41-C31	-7.00	1.46	1.55
2	A	1300	JDC	C51-C41	-6.23	1.44	1.54
2	B	1300	JDC	C51-C41	-6.11	1.44	1.54
2	A	1300	JDC	C1-N2	-4.03	1.41	1.46

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1300	JDC	C4E-C41-C51	-4.20	104.31	109.08
3	A	1301	CIT	C3-C4-C5	-3.80	108.89	114.96
2	B	1300	JDC	C3A-C2A-N2A	-3.59	103.44	110.88
2	B	1300	JDC	C4E-C41-C31	-3.59	106.48	110.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	1300	JDC	C3A-C2A-N2A	-3.19	104.29	110.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1300	JDC	2	0
4	B	1302	OLC	1	0
4	B	1303	OLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/480 (93%)	-0.26	2 (0%) 93 92	48, 79, 139, 260	0
1	B	448/480 (93%)	-0.15	14 (3%) 52 45	39, 72, 141, 239	0
All	All	895/960 (93%)	-0.20	16 (1%) 71 68	39, 76, 142, 260	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	204	ASP	4.5
1	B	217	ASP	3.4
1	A	206	ASP	3.4
1	B	201	VAL	3.2
1	B	305	SER	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	PEG	B	1305	7/7	0.82	0.35	10.94	93,94,100,101	0
3	CIT	A	1301	13/13	0.76	0.35	5.04	129,148,161,166	0
4	OLC	B	1302	16/25	0.93	0.23	2.52	54,68,72,78	0
4	OLC	B	1303	21/25	0.87	0.22	1.21	52,75,134,145	0
2	JDC	B	1300	34/34	0.95	0.18	-0.03	48,65,92,99	0
5	PEG	B	1304	7/7	0.74	0.19	-0.23	75,93,104,105	0
2	JDC	A	1300	34/34	0.96	0.16	-0.25	43,62,79,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.