



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:10 AM GMT

PDB ID : 3DKQ
Title : Crystal structure of Putative Oxygenase (YP_001051978.1) from SHE-
WANELLA BALTICA OS155 at 2.26 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2008-06-25
Resolution : 2.26 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1640 (2.28-2.24)
Clashscore	102246	1095 (2.26-2.26)
Ramachandran outliers	100387	1063 (2.26-2.26)
Sidechain outliers	100360	1063 (2.26-2.26)
RSRZ outliers	91569	1647 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	<div> <div>13%</div> <div>88%</div> <div>7%</div> <div>5%</div> </div>
1	B	243	<div> <div>16%</div> <div>80%</div> <div>5%</div> <div>15%</div> </div>
1	C	243	<div> <div>16%</div> <div>83%</div> <div>7%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	B	502	-	-	-	X
4	GOL	A	503	-	-	-	X
4	GOL	A	507	-	-	-	X
4	GOL	C	503	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PKHD-type hydroxylase Sbal_3634.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	Se	0	7	0
			1867	1180	332	349	6			
1	B	207	Total	C	N	O	Se	0	2	0
			1640	1036	288	312	4			
1	C	221	Total	C	N	O	Se	0	4	0
			1778	1125	309	339	5			

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	leader sequence	UNP A3D8P6
A	-17	GLY	-	leader sequence	UNP A3D8P6
A	-16	SER	-	leader sequence	UNP A3D8P6
A	-15	ASP	-	leader sequence	UNP A3D8P6
A	-14	LYS	-	leader sequence	UNP A3D8P6
A	-13	ILE	-	leader sequence	UNP A3D8P6
A	-12	HIS	-	leader sequence	UNP A3D8P6
A	-11	HIS	-	leader sequence	UNP A3D8P6
A	-10	HIS	-	leader sequence	UNP A3D8P6
A	-9	HIS	-	leader sequence	UNP A3D8P6
A	-8	HIS	-	leader sequence	UNP A3D8P6
A	-7	HIS	-	leader sequence	UNP A3D8P6
A	-6	GLU	-	leader sequence	UNP A3D8P6
A	-5	ASN	-	leader sequence	UNP A3D8P6
A	-4	LEU	-	leader sequence	UNP A3D8P6
A	-3	TYR	-	leader sequence	UNP A3D8P6
A	-2	PHE	-	leader sequence	UNP A3D8P6
A	-1	GLN	-	leader sequence	UNP A3D8P6
A	0	GLY	-	leader sequence	UNP A3D8P6
B	-18	MSE	-	leader sequence	UNP A3D8P6
B	-17	GLY	-	leader sequence	UNP A3D8P6
B	-16	SER	-	leader sequence	UNP A3D8P6
B	-15	ASP	-	leader sequence	UNP A3D8P6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	LYS	-	leader sequence	UNP A3D8P6
B	-13	ILE	-	leader sequence	UNP A3D8P6
B	-12	HIS	-	leader sequence	UNP A3D8P6
B	-11	HIS	-	leader sequence	UNP A3D8P6
B	-10	HIS	-	leader sequence	UNP A3D8P6
B	-9	HIS	-	leader sequence	UNP A3D8P6
B	-8	HIS	-	leader sequence	UNP A3D8P6
B	-7	HIS	-	leader sequence	UNP A3D8P6
B	-6	GLU	-	leader sequence	UNP A3D8P6
B	-5	ASN	-	leader sequence	UNP A3D8P6
B	-4	LEU	-	leader sequence	UNP A3D8P6
B	-3	TYR	-	leader sequence	UNP A3D8P6
B	-2	PHE	-	leader sequence	UNP A3D8P6
B	-1	GLN	-	leader sequence	UNP A3D8P6
B	0	GLY	-	leader sequence	UNP A3D8P6
C	-18	MSE	-	leader sequence	UNP A3D8P6
C	-17	GLY	-	leader sequence	UNP A3D8P6
C	-16	SER	-	leader sequence	UNP A3D8P6
C	-15	ASP	-	leader sequence	UNP A3D8P6
C	-14	LYS	-	leader sequence	UNP A3D8P6
C	-13	ILE	-	leader sequence	UNP A3D8P6
C	-12	HIS	-	leader sequence	UNP A3D8P6
C	-11	HIS	-	leader sequence	UNP A3D8P6
C	-10	HIS	-	leader sequence	UNP A3D8P6
C	-9	HIS	-	leader sequence	UNP A3D8P6
C	-8	HIS	-	leader sequence	UNP A3D8P6
C	-7	HIS	-	leader sequence	UNP A3D8P6
C	-6	GLU	-	leader sequence	UNP A3D8P6
C	-5	ASN	-	leader sequence	UNP A3D8P6
C	-4	LEU	-	leader sequence	UNP A3D8P6
C	-3	TYR	-	leader sequence	UNP A3D8P6
C	-2	PHE	-	leader sequence	UNP A3D8P6
C	-1	GLN	-	leader sequence	UNP A3D8P6
C	0	GLY	-	leader sequence	UNP A3D8P6

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ni 1 1	0	0
2	A	1	Total Ni 1 1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Ni	0	0
			1	1		

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			5	3	2		
3	A	1	Total	C	N	0	0
			5	3	2		
3	B	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	C	1	Total	C	N	0	0
			5	3	2		
3	B	1	Total	C	N	0	0
			5	3	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	119	Total	O	0	0
			119	119		
5	B	52	Total	O	0	2
			54	54		
5	C	73	Total	O	0	0
			73	73		

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	153.45Å 153.45Å 84.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.86 – 2.26 28.86 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.86-2.26) 100.0 (28.86-2.26)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.26Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.232 0.204 , 0.237	Depositor DCC
R_{free} test set	2424 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 54.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 47883 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5600	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/1919	0.96	6/2586 (0.2%)
1	B	0.68	0/1670	0.84	4/2255 (0.2%)
1	C	0.70	0/1821	0.88	5/2458 (0.2%)
All	All	0.70	0/5410	0.89	15/7299 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	ARG	NE-CZ-NH2	-10.97	114.81	120.30
1	A	184	ARG	NE-CZ-NH1	10.66	125.63	120.30
1	C	184	ARG	NE-CZ-NH1	9.78	125.19	120.30
1	C	184	ARG	NE-CZ-NH2	-9.60	115.50	120.30
1	A	179	ARG	NE-CZ-NH1	9.45	125.03	120.30
1	A	184	ARG	NE-CZ-NH2	-8.50	116.05	120.30
1	B	184	ARG	NE-CZ-NH2	-7.83	116.39	120.30
1	B	184	ARG	NE-CZ-NH1	6.85	123.73	120.30
1	B	219	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	B	220	ARG	NE-CZ-NH2	-5.81	117.39	120.30
1	A	219[A]	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	219[B]	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	C	110	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	C	220	ARG	NE-CZ-NH2	-5.55	117.53	120.30
1	C	80	TYR	N-CA-C	-5.06	97.34	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1867	0	1849	10	0
1	B	1640	0	1590	5	0
1	C	1778	0	1740	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	10	0	10	0	0
3	B	10	0	10	0	0
3	C	10	0	10	0	0
4	A	30	0	40	0	0
4	C	6	0	8	0	0
5	A	119	0	0	1	0
5	B	54	0	0	0	0
5	C	73	0	0	0	0
All	All	5600	0	5257	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:MSE:HE2	1:A:97:ILE:HB	1.44	0.98
1:C:-4:LEU:HD21	1:C:3:ILE:HD11	1.55	0.86
1:A:36:MSE:HE2	1:A:97:ILE:CB	2.11	0.80
1:A:75:LEU:HB3	1:A:177:MSE:HE3	1.68	0.75
1:A:32:THR:HG21	1:A:35:ALA:HB3	1.77	0.66
1:A:36:MSE:HE2	1:A:97:ILE:CG2	2.34	0.57
1:A:171[B]:PHE:HE2	1:A:173:TRP:CZ2	2.24	0.55
1:A:186:LEU:HD21	1:B:213:VAL:HG13	1.89	0.54
1:C:2:LEU:HD11	1:C:139:GLN:HE22	1.74	0.53
1:A:32:THR:CG2	1:A:35:ALA:HB3	2.38	0.52
1:C:14:VAL:HG11	1:C:145:ALA:HB1	1.91	0.52
1:C:-4:LEU:HD12	1:C:69:GLN:CD	2.32	0.50
1:C:177:MSE:CE	1:C:218:LEU:HD11	2.41	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:-4:LEU:HD21	1:C:3:ILE:CD1	2.37	0.49
1:B:75:LEU:HB3	1:B:177:MSE:HE3	1.94	0.48
1:A:100:ALA:C	1:A:101:ILE:HD12	2.36	0.47
1:C:-4:LEU:HD22	1:C:-2:PHE:CE1	2.51	0.45
1:B:14:VAL:HG11	1:B:145:ALA:HB1	1.97	0.45
1:C:3:ILE:HG23	1:C:150:LEU:HB3	1.98	0.45
1:C:132:ILE:HG23	1:C:155:SER:HB3	2.02	0.42
1:B:186:LEU:HD23	1:B:186:LEU:C	2.40	0.41
1:B:203:ALA:HB3	1:B:206:GLU:HG2	2.03	0.41
1:C:177:MSE:HE2	1:C:218:LEU:HD11	2.03	0.41
1:A:177:MSE:HE2	5:A:588:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/243 (97%)	234 (100%)	1 (0%)	0	100	100
1	B	203/243 (84%)	200 (98%)	3 (2%)	0	100	100
1	C	221/243 (91%)	216 (98%)	5 (2%)	0	100	100
All	All	659/729 (90%)	650 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/204 (99%)	199 (98%)	3 (2%)	72	82
1	B	174/204 (85%)	172 (99%)	2 (1%)	80	88
1	C	193/204 (95%)	187 (97%)	6 (3%)	47	58
All	All	569/612 (93%)	558 (98%)	11 (2%)	66	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	93	PHE
1	A	148	LEU
1	B	83	LEU
1	B	179	ARG
1	C	-4	LEU
1	C	-2	PHE
1	C	19[A]	GLU
1	C	19[B]	GLU
1	C	28	ASP
1	C	93	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	139	GLN
1	B	69	GLN
1	B	124	ASN
1	B	189	GLN
1	C	124	ASN
1	C	139	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 3 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	IMD	A	501	2	3,5,5	0.38	0	4,5,5	0.97	0
3	IMD	A	502	-	3,5,5	0.42	0	4,5,5	0.90	0
4	GOL	A	503	-	5,5,5	0.45	0	5,5,5	0.78	0
4	GOL	A	504	-	5,5,5	0.30	0	5,5,5	0.30	0
4	GOL	A	505	-	5,5,5	0.38	0	5,5,5	0.68	0
4	GOL	A	506	-	5,5,5	0.47	0	5,5,5	0.75	0
4	GOL	A	507	-	5,5,5	0.39	0	5,5,5	0.62	0
3	IMD	B	501	-	3,5,5	0.54	0	4,5,5	0.73	0
3	IMD	B	502	-	3,5,5	0.46	0	4,5,5	0.80	0
3	IMD	C	501	-	3,5,5	0.32	0	4,5,5	0.74	0
3	IMD	C	502	-	3,5,5	0.26	0	4,5,5	0.88	0
4	GOL	C	503	-	5,5,5	0.45	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	A	501	2	-	0/0/0/0	0/1/1/1
3	IMD	A	502	-	-	0/0/0/0	0/1/1/1
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
4	GOL	A	504	-	-	0/4/4/4	0/0/0/0
4	GOL	A	505	-	-	0/4/4/4	0/0/0/0
4	GOL	A	506	-	-	0/4/4/4	0/0/0/0
4	GOL	A	507	-	-	0/4/4/4	0/0/0/0
3	IMD	B	501	-	-	0/0/0/0	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	B	502	-	-	0/0/0/0	0/1/1/1
3	IMD	C	501	-	-	0/0/0/0	0/1/1/1
3	IMD	C	502	-	-	0/0/0/0	0/1/1/1
4	GOL	C	503	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/243 (92%)	0.78	32 (14%) 4 4	45, 52, 83, 99	0
1	B	203/243 (83%)	1.21	39 (19%) 2 1	42, 52, 74, 109	0
1	C	216/243 (88%)	1.04	39 (18%) 2 1	44, 52, 78, 104	0
All	All	643/729 (88%)	1.00	110 (17%) 2 2	42, 52, 79, 109	0

All (110) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-4	LEU	11.3
1	B	95	TYR	8.6
1	C	135	THR	7.9
1	C	30	ASN	7.9
1	B	136	TYR	7.8
1	C	134	ASP	7.4
1	B	97	ILE	7.2
1	B	-5	ASN	6.9
1	A	38	THR	6.8
1	A	171[A]	PHE	6.7
1	C	-1	GLN	6.3
1	C	29	GLY	6.3
1	C	-2	PHE	6.2
1	B	135	THR	5.9
1	B	96	HIS	5.6
1	A	30	ASN	5.5
1	B	90	GLY	5.3
1	A	32	THR	4.8
1	C	31	GLN	4.7
1	B	201	THR	4.5
1	A	224	LEU	4.5
1	B	29	GLY	4.2
1	B	42	ARG	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	115	ALA	4.2
1	B	47	ASP	4.2
1	C	-3	TYR	4.2
1	C	12	GLN	4.1
1	A	31	GLN	4.1
1	B	156	LEU	4.1
1	C	165	GLY	4.0
1	C	137	GLY	3.9
1	C	123	GLU	3.9
1	B	198	THR	3.9
1	B	27	ILE	3.9
1	C	136	TYR	3.8
1	A	33	SER	3.7
1	B	89	GLY	3.7
1	A	35	ALA	3.7
1	B	94	GLY	3.6
1	A	-3	TYR	3.6
1	B	49	ASP	3.5
1	C	-4	LEU	3.5
1	A	29	GLY	3.5
1	A	37	ALA	3.5
1	A	28	ASP	3.4
1	C	28	ASP	3.4
1	C	90	GLY	3.3
1	C	169	ALA	3.2
1	A	-1	GLN	3.2
1	C	170	ALA	3.2
1	C	41	LYS	3.2
1	B	28	ASP	3.2
1	B	116	THR	3.2
1	A	105	PRO	3.1
1	B	150	LEU	3.0
1	C	115	ALA	3.0
1	C	40	ARG	3.0
1	A	116	THR	2.9
1	A	136	TYR	2.9
1	A	130	LEU	2.9
1	A	205	GLN	2.8
1	C	42	ARG	2.8
1	A	15	SER	2.8
1	B	149	VAL	2.8
1	B	117	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	24	ARG	2.8
1	B	170	ALA	2.8
1	B	148	LEU	2.7
1	C	105	PRO	2.7
1	A	115	ALA	2.6
1	A	150	LEU	2.6
1	B	98	ASP	2.6
1	A	-4	LEU	2.5
1	C	140	SER	2.5
1	B	169	ALA	2.5
1	A	-5	ASN	2.5
1	A	169	ALA	2.5
1	C	116	THR	2.5
1	B	8	VAL	2.4
1	A	-2	PHE	2.4
1	B	155	SER	2.4
1	B	199	ALA	2.4
1	C	27	ILE	2.4
1	C	171	PHE	2.4
1	C	151	TYR	2.4
1	B	203	ALA	2.4
1	C	149	VAL	2.3
1	A	16	HIS	2.3
1	C	106	ASP	2.3
1	C	163	LEU	2.3
1	A	190	LEU	2.3
1	A	151	TYR	2.3
1	C	162	VAL	2.3
1	A	201	THR	2.2
1	C	114	SER	2.2
1	B	-2	PHE	2.2
1	B	23	ALA	2.1
1	C	150	LEU	2.1
1	C	138	GLN	2.1
1	C	139	GLN	2.1
1	C	122	PRO	2.1
1	C	117	LEU	2.1
1	B	110	ARG	2.1
1	B	21	LEU	2.1
1	B	46	LEU	2.1
1	A	12[A]	GLN	2.0
1	A	114	SER	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	-1	GLN	2.0
1	A	117	LEU	2.0
1	C	119	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	C	503	6/6	0.86	0.26	8.73	92,96,97,99	0
3	IMD	B	502	5/5	0.73	0.44	6.36	72,76,77,78	0
4	GOL	A	507	6/6	0.88	0.21	5.54	76,84,85,87	0
4	GOL	A	503	6/6	0.76	0.22	2.53	79,81,84,84	0
4	GOL	A	505	6/6	0.87	0.22	1.32	74,80,82,83	0
4	GOL	A	506	6/6	0.73	0.24	0.61	79,83,85,88	0
3	IMD	B	501	5/5	0.92	0.23	0.07	87,88,90,90	0
3	IMD	C	502	5/5	0.92	0.19	0.02	62,63,65,68	0
3	IMD	A	502	5/5	0.98	0.16	-0.88	46,48,51,53	0
3	IMD	C	501	5/5	0.96	0.12	-1.18	48,49,53,53	0
3	IMD	A	501	5/5	0.98	0.10	-1.36	36,40,45,46	0
2	NI	A	500	1/1	0.99	0.06	-1.53	42,42,42,42	0
2	NI	C	500	1/1	0.98	0.04	-2.71	53,53,53,53	0
2	NI	B	500	1/1	0.86	0.09	-	59,59,59,59	1
4	GOL	A	504	6/6	0.90	0.22	-	70,75,78,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.