



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2016 – 10:49 PM EDT

PDB ID : 5DMZ  
Title : Structure of human Bub1 kinase domain phosphorylated at Ser969  
Authors : Breit, C.; Weir, J.R.; Musacchio, A.  
Deposited on : 2015-09-09  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>10%</div> <div>•</div> <div>7%</div> </div> </div>
1	B	365	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>7%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11114 atoms, of which 5508 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Mitotic checkpoint serine/threonine-protein kinase BUB1.

Mol	Chain	Residues	Atoms								ZeroOcc	AltConf	Trace
1	A	340	Total	C	H	N	O	P	S	0	0	0	
			5495	1796	2732	456	488	1	22				
1	B	340	Total	C	H	N	O	P	S	0	0	0	
			5515	1796	2752	456	488	1	22				

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	721	GLY	-	expression tag	UNP O43683
A	722	PRO	-	expression tag	UNP O43683
A	723	MET	-	expression tag	UNP O43683
A	724	ASP	-	expression tag	UNP O43683
A	725	PRO	-	expression tag	UNP O43683
B	721	GLY	-	expression tag	UNP O43683
B	722	PRO	-	expression tag	UNP O43683
B	723	MET	-	expression tag	UNP O43683
B	724	ASP	-	expression tag	UNP O43683
B	725	PRO	-	expression tag	UNP O43683

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		
2	B	1	Total	C	H	N	O	P	0	0
			39	10	12	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	O	0	0
			11	11		
4	B	13	Total	O	0	0
			13	13		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1



- Molecule 1: Mitotic checkpoint serine/threonine-protein kinase BUB1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.38Å 66.34Å 106.54Å 90.00° 98.34° 90.00°	Depositor
Resolution (Å)	43.30 – 2.40 56.15 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.30-2.40) 99.8 (56.15-2.40)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.35 (at 2.40Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.210 , 0.245 0.211 , 0.245	Depositor DCC
$R_{free}$ test set	1691 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	56.4	Xtriage
Anisotropy	0.532	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11114	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/2824	0.43	0/3811
1	B	0.26	0/2824	0.43	0/3811
All	All	0.26	0/5648	0.43	0/7622

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2763	2732	2763	25	0
1	B	2763	2752	2763	26	0
2	A	27	12	12	1	0
2	B	27	12	12	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	11	0	0	0	0
4	B	13	0	0	0	0
All	All	5606	5508	5550	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:906:GLU:OE1	1:B:1051:ARG:NH1	2.12	0.82
1:B:889:MET:HE3	1:B:897:PHE:HD2	1.56	0.71
1:A:821:LYS:NZ	2:A:1101:ADP:O2A	2.23	0.71
1:A:900:ARG:NH1	1:A:937:ASP:O	2.26	0.68
1:B:900:ARG:NH1	1:B:937:ASP:O	2.29	0.65
1:B:821:LYS:NZ	2:B:1101:ADP:O1A	2.29	0.65
1:A:961:ILE:HG21	1:A:981:PRO:HB2	1.85	0.59
1:A:932:GLN:NE2	1:A:934:ASP:O	2.36	0.59
1:B:1024:HIS:NE2	1:B:1062:TYR:OH	2.25	0.57
1:B:932:GLN:NE2	1:B:934:ASP:O	2.33	0.55
1:B:938:LEU:CD2	1:B:1051:ARG:HD3	2.37	0.55
1:A:956:PHE:HB3	1:A:960:THR:HG21	1.90	0.53
1:A:889:MET:HE3	1:A:897:PHE:HD2	1.72	0.53
1:A:831:PHE:CD1	1:A:855:ALA:HB2	2.47	0.49
1:B:840:ARG:CZ	1:B:954:LYS:HD2	2.41	0.49
1:A:782:LEU:HD11	1:A:856:HIS:CG	2.48	0.48
1:B:1034:VAL:CG1	1:B:1045:PRO:HB3	2.44	0.48
1:A:736:PHE:HB3	1:A:960:THR:HG22	1.96	0.47
1:B:772:PRO:HD3	1:B:858:PHE:CE2	2.50	0.47
1:B:736:PHE:CD2	1:B:960:THR:HG22	2.51	0.46
1:A:983:ASN:OD1	1:A:984:TYR:N	2.44	0.46
1:B:850:MET:HB2	1:B:945:ILE:HG22	1.99	0.44
1:A:1078:LEU:CD2	1:B:930:LEU:HD11	2.47	0.44
1:A:1059:GLN:HA	1:A:1063:THR:HG23	2.00	0.44
1:B:889:MET:HE3	1:B:897:PHE:CD2	2.44	0.44
1:B:1027:MET:SD	1:B:1057:VAL:HG21	2.57	0.44
1:A:961:ILE:HG21	1:A:981:PRO:CB	2.48	0.44
1:B:821:LYS:NZ	2:B:1101:ADP:PA	2.91	0.43
1:A:975:GLU:OE1	1:A:975:GLU:N	2.50	0.43
1:A:889:MET:HE1	1:A:894:VAL:HA	2.00	0.43
1:B:938:LEU:HD23	1:B:1051:ARG:HD3	2.00	0.43
1:A:741:PRO:HG3	1:A:962:PHE:CD1	2.54	0.43
1:A:771:LEU:HD12	1:A:861:GLY:HA3	2.00	0.43
1:A:929:PHE:HA	1:A:941:GLY:HA2	1.99	0.43
1:A:782:LEU:CD1	1:A:856:HIS:CD2	3.02	0.43
1:A:929:PHE:CZ	1:A:1073:LEU:HD23	2.54	0.43
1:B:838:MET:HE2	1:B:851:LYS:HE2	2.01	0.43
1:B:759:SER:OG	1:B:766:GLU:OE2	2.13	0.42
1:B:929:PHE:HA	1:B:941:GLY:HA2	2.01	0.42
1:B:741:PRO:HG3	1:B:962:PHE:CD1	2.55	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:884:THR:HB	1:B:885:PRO:HD2	2.00	0.41
1:A:961:ILE:CG2	1:A:981:PRO:HB2	2.50	0.41
1:B:929:PHE:CZ	1:B:1073:LEU:HD23	2.56	0.41
1:B:1082:ARG:O	1:B:1083:SER:OG	2.32	0.41
1:A:1027:MET:SD	1:A:1057:VAL:HG21	2.61	0.41
1:A:870:SER:HB2	1:A:1081:LYS:HE2	2.02	0.41
1:B:1027:MET:SD	1:B:1054:LEU:HD23	2.61	0.40
1:A:758:VAL:O	1:A:764:THR:HG21	2.21	0.40
1:B:737:ILE:HD12	1:B:737:ILE:N	2.37	0.40
1:A:767:TRP:HD1	1:A:858:PHE:CZ	2.39	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/365 (92%)	326 (97%)	9 (3%)	0	100	100
1	B	335/365 (92%)	328 (98%)	7 (2%)	0	100	100
All	All	670/730 (92%)	654 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/323 (94%)	300 (99%)	3 (1%)	82	93
1	B	303/323 (94%)	301 (99%)	2 (1%)	88	95
All	All	606/646 (94%)	601 (99%)	5 (1%)	86	94

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	782	LEU
1	A	930	LEU
1	A	1078	LEU
1	B	930	LEU
1	B	939	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	SEP	A	969	1	7,9,10	1.64	1 (14%)	8,12,14	1.68	1 (12%)
1	SEP	B	969	1	7,9,10	1.67	1 (14%)	8,12,14	1.92	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	969	1	-	0/5/8/10	0/0/0/0
1	SEP	B	969	1	-	0/5/8/10	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	969	SEP	P-O1P	3.23	1.61	1.50
1	B	969	SEP	P-O1P	3.36	1.61	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	969	SEP	O-C-CA	-2.12	120.04	125.72
1	A	969	SEP	OG-CB-CA	3.97	111.72	108.26
1	B	969	SEP	OG-CB-CA	4.63	112.30	108.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ADP	A	1101	3	24,29,29	0.98	1 (4%)	23,45,45	1.83	1 (4%)
2	ADP	B	1101	3	24,29,29	0.97	1 (4%)	23,45,45	1.92	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1101	3	-	0/12/32/32	0/3/3/3
2	ADP	B	1101	3	-	0/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	ADP	C5-C4	2.90	1.47	1.40
2	A	1101	ADP	C5-C4	3.00	1.47	1.40

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1101	ADP	N3-C2-N1	-7.61	122.89	128.87
2	A	1101	ADP	N3-C2-N1	-7.15	123.25	128.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	ADP	1	0
2	B	1101	ADP	2	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/365 (92%)	0.40	11 (3%) 51 51	48, 74, 112, 126	2 (0%)
1	B	339/365 (92%)	0.43	11 (3%) 51 51	51, 77, 116, 133	1 (0%)
All	All	678/730 (92%)	0.41	22 (3%) 51 51	48, 76, 115, 133	3 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1020	ARG	4.6
1	B	1018	LEU	3.5
1	A	1021	ARG	2.9
1	B	1019	PHE	2.9
1	A	783	GLY	2.8
1	A	1020	ARG	2.8
1	B	783	GLY	2.7
1	A	748	PHE	2.7
1	A	767	TRP	2.7
1	B	1030	GLU	2.6
1	A	784	SER	2.5
1	A	756	LYS	2.5
1	B	935	GLU	2.3
1	A	782	LEU	2.3
1	B	941	GLY	2.2
1	A	938	LEU	2.2
1	A	1025	LEU	2.2
1	B	765	PHE	2.2
1	B	1025	LEU	2.2
1	A	765	PHE	2.1
1	B	1083	SER	2.1
1	B	1022	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	SEP	A	969	10/11	0.92	0.21	-	68,71,83,93	0
1	SEP	B	969	10/11	0.93	0.18	-	73,78,95,95	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ADP	A	1101	27/27	0.96	0.18	0.57	48,61,83,122	0
2	ADP	B	1101	27/27	0.96	0.19	0.39	51,65,79,131	0
3	MG	A	1102	1/1	0.86	0.16	-0.49	57,57,57,57	0
3	MG	B	1102	1/1	0.95	0.12	-1.03	53,53,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.