wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 07:49 PM BST

PDB ID	:	2DNW
Title	:	Solution structure of RSGI RUH-059, an ACP domain of acyl carrier protein, mitochondrial [Precursor] from human cDNA
Authors	:	Abe, T.; Hirota, H.; Kurosaki, C.; Yoshida, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2006-04-26

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	99	27% 46% • 23%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 9 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy, target function.*

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:13-A:88 (76)	0.24	9

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 5 single-model clusters were found.

Cluster number	Models
1	3, 4, 8, 10, 16, 17
2	6, 7, 9, 12, 13, 20
3	1, 2, 5
Single-model clusters	11; 14; 15; 18; 19

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1511 atoms, of which 746 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Acyl carrier protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	99	1511	484	746	115	160	6	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP O14561
A	2	SER	-	CLONING ARTIFACT	UNP O14561
A	3	SER	-	CLONING ARTIFACT	UNP O14561
A	4	GLY	-	CLONING ARTIFACT	UNP O14561
A	5	SER	-	CLONING ARTIFACT	UNP O14561
A	6	SER	-	CLONING ARTIFACT	UNP O14561
A	7	GLY	-	CLONING ARTIFACT	UNP O14561
A	94	SER	-	CLONING ARTIFACT	UNP O14561
A	95	GLY	-	CLONING ARTIFACT	UNP O14561
A	96	PRO	-	CLONING ARTIFACT	UNP O14561
A	97	SER	-	CLONING ARTIFACT	UNP O14561
A	98	SER	-	CLONING ARTIFACT	UNP O14561
A	99	GLY	-	CLONING ARTIFACT	UNP O14561

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Acyl carrier protein

Chain A: 27% 46% • 23%

G1 S2 S3 G4 S5 S6 G7 M8 P9 P10 L11 T12 L13 E14 G15 I16 V20 L21 Y22 V23 L24 K25 L26 Y27 D28 K29 I30 D31 P32 E33 K34 L35 S36 V37 N38 S39 H40 F41 M42 K43 D44 L45 G46 L47 D48 S49 L50 D51 Q52 V53 E54 I55 A58 M59 E60 D61 E62 F63

G64 F65 E66 I67 P68 D69 I70 D71 A72 E73 K74 L75 M76 C77 E80 I81 V82 D83 Y84 I85 A86 D87 K88 K89 D90 V91 Y92 E93 S94 G95 P96 S97 S98 G99

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 9. Colouring as in section 4.1 above.

- Molecule 1: Acyl carrier protein

Chain A: 36% 35% • • 23%

G1 S2 S3 G4 S5 S6 G7 M8 P9 P10 L11 T12 L13 E14 G15 I16 Q17 V20 V23 L24 K25 L26 Y27 D28 K29 I30 D31 P32 E33 K34 L35 S36 S39 H40 F41 M42 K43 D44 L45 G46 L47 D48 S49 Q52 V53 E54 I55 E62 F63 I67 P68 D69 I70 D71

K74 C77 P78 Q79 E80 I81 V82 D83 Y84 I85 A86 D87 K88 K89 D90 V91 Y92 E93 S94 G95 P96 S97 S98 G99

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy, structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	refinement	1.0.8

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	614	607	607	34±7
All	All	12280	12140	12140	689

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

5 of 211 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:30:ILE:HD11	1:A:55:ILE:HD11	0.97	1.35	12	9
1:A:23:VAL:CG1	1:A:58:ALA:HB1	0.88	1.97	13	16
1:A:35:LEU:HD23	1:A:36:SER:N	0.82	1.89	5	4
1:A:51:ASP:OD1	1:A:55:ILE:HD12	0.81	1.74	19	1
1:A:50:LEU:O	1:A:53:VAL:HG12	0.79	1.78	3	4

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	76/99 (77%)	62±2 (82±3%)	13±3 (17±3%)	1±1 (2±1%)	18 63
All	All	1520/1980 (77%)	1242 (82%)	255 (17%)	23 (2%)	18 63

5 of 7 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	45	LEU	8
1	A	44	ASP	5
1	A	35	LEU	5
1	A	14	GLU	2
1	A	40	HIS	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	70/88 (80%)	53±2 (75±3%)	17±2 (25±3%)	3 27
All	All	1400/1760 (80%)	1056 (75%)	344 (25%)	3 27

5 of 42 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	74	LYS	20
1	A	77	CYS	20
1	A	33	GLU	19
1	A	13	LEU	19
1	A	70	ILE	18

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided