



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:38 AM GMT

PDB ID : 2DQW  
Title : Crystal Structure of Dihydropteroate Synthase (FolP) from *Thermus thermophilus* HB8  
Authors : Bagautdinov, B.; Kunishima, N.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2006-05-31  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	<div> <div>9%</div> <div>70%</div> <div>13%</div> <div>•</div> <div>15%</div> </div>
1	B	294	<div> <div>15%</div> <div>66%</div> <div>17%</div> <div>•</div> <div>16%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydropteroate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1926	1215	366	338	7			
1	B	247	Total	C	N	O	S	0	0	0
			1908	1205	363	335	5			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	339	Total	O	0	0
			339	339		
2	B	272	Total	O	0	0
			272	272		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Dihydropteroate synthase



#### • Molecule 1: Dihydropteroate synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	109.99 Å 109.99 Å 88.32 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.88 – 1.65 29.88 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.88-1.65) 99.8 (29.88-1.65)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.38 (at 1.65 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.213 , 0.237 0.213 , 0.237	Depositor DCC
$R_{free}$ test set	3683 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 68.5	EDS
Estimated twinning fraction	0.036 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 72756 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4445	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/1954	0.59	0/2644
1	B	0.25	0/1936	0.54	0/2618
All	All	0.27	0/3890	0.56	0/5262

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1926	0	2005	42	0
1	B	1908	0	1987	42	0
2	A	339	0	0	11	0
2	B	272	0	0	12	0
All	All	4445	0	3992	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:VAL:HG13	1:B:157:PRO:HD3	1.48	0.94

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:ARG:HD3	2:B:375:HOH:O	1.73	0.87
1:A:232:ARG:HG2	2:A:350:HOH:O	1.79	0.81
1:B:56:GLU:HA	1:B:59:ARG:HD3	1.65	0.75
1:A:55:LEU:O	1:A:59:ARG:HG3	1.89	0.73
1:A:53:ARG:HA	1:A:53:ARG:NE	2.05	0.71
1:B:15:ARG:NH2	2:B:442:HOH:O	2.23	0.71
1:B:268:ARG:NH1	2:B:522:HOH:O	2.24	0.70
1:A:184:ARG:NH1	2:A:417:HOH:O	2.26	0.67
1:B:154:MET:HE2	2:B:542:HOH:O	1.96	0.66
1:A:53:ARG:HA	1:A:53:ARG:HE	1.62	0.65
1:B:120:LEU:HD23	1:B:124:ALA:HB3	1.79	0.65
1:A:271:ASP:HB3	2:A:478:HOH:O	1.97	0.64
1:B:60:GLU:O	1:B:64:GLU:HG3	1.97	0.64
1:A:93:LEU:HB2	1:A:94:PRO:HD3	1.79	0.63
1:B:154:MET:HG2	2:B:509:HOH:O	1.99	0.63
1:B:128:ASN:OD1	1:B:268:ARG:NH2	2.32	0.62
1:B:157:PRO:HG2	2:B:381:HOH:O	1.99	0.62
1:A:271:ASP:HB3	2:A:568:HOH:O	1.99	0.62
1:B:181:ALA:O	1:B:185:ARG:HG3	2.00	0.62
2:A:394:HOH:O	1:B:250:HIS:HD2	1.83	0.61
1:B:157:PRO:HG3	2:B:543:HOH:O	2.03	0.59
1:A:53:ARG:NH1	2:A:533:HOH:O	2.36	0.59
1:B:271:ASP:HB2	2:B:346:HOH:O	2.03	0.58
1:B:21:ASP:OD1	1:B:22:ARG:HD2	2.05	0.56
1:A:86:GLU:CD	1:A:112:LYS:HD3	2.26	0.56
1:A:87:GLU:HG2	1:A:90:ARG:HD3	1.89	0.54
1:A:37:LEU:HD11	1:A:95:VAL:HG11	1.88	0.54
1:A:127:LEU:HD11	1:A:138:MET:HE1	1.90	0.54
1:B:157:PRO:HA	2:B:543:HOH:O	2.07	0.54
1:B:58:ALA:O	1:B:62:VAL:HG23	2.07	0.54
1:A:159:PRO:HA	1:A:162:MET:CE	2.38	0.53
1:A:39:PRO:HD3	1:A:91:ARG:NH1	2.24	0.53
1:B:213:LEU:HD21	1:B:230:LEU:HD11	1.90	0.53
1:A:158:ASP:OD2	1:A:161:THR:HG23	2.08	0.53
1:B:200:PHE:CE2	1:B:213:LEU:HG	2.44	0.53
1:A:55:LEU:HD21	1:A:59:ARG:HH21	1.74	0.53
1:A:114:GLU:H	1:A:114:GLU:CD	2.13	0.51
1:B:87:GLU:HG2	1:B:90:ARG:NH1	2.26	0.51
1:A:261:MET:SD	1:B:261:MET:SD	3.09	0.50
1:B:185:ARG:NH2	2:B:413:HOH:O	2.44	0.50
1:B:93:LEU:HB2	1:B:94:PRO:HD3	1.94	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:GLY:HA3	1:A:276:ARG:HG3	1.95	0.48
1:A:200:PHE:CE2	1:A:213:LEU:HG	2.49	0.48
1:B:133:LEU:HD12	1:B:151:VAL:HG22	1.95	0.48
1:B:227:LEU:HD21	1:B:268:ARG:NH2	2.28	0.48
1:A:53:ARG:NH2	2:A:533:HOH:O	2.47	0.47
1:A:86:GLU:HG3	1:A:112:LYS:HD3	1.95	0.47
1:A:129:ASP:OD2	1:A:132:GLY:N	2.47	0.47
1:A:74:GLU:HG2	2:A:493:HOH:O	2.15	0.47
1:A:213:LEU:HD21	1:A:230:LEU:HD11	1.97	0.47
1:B:261:MET:HG3	1:B:286:TYR:CE2	2.51	0.46
1:B:96:LEU:O	1:B:100:LEU:HG	2.16	0.46
1:A:159:PRO:HA	1:A:162:MET:HE2	1.98	0.45
1:A:86:GLU:HG3	1:A:112:LYS:CD	2.47	0.45
1:B:20:ARG:HD3	1:B:124:ALA:O	2.17	0.45
1:B:130:VAL:O	1:B:154:MET:SD	2.76	0.44
1:A:86:GLU:CG	1:A:112:LYS:HD3	2.48	0.43
1:B:19:LEU:HB2	1:B:22:ARG:HG2	2.00	0.43
1:A:130:VAL:HG12	1:A:130:VAL:O	2.18	0.43
1:B:86:GLU:C	1:B:88:GLU:H	2.22	0.43
1:B:147:VAL:HG12	1:B:148:ALA:O	2.18	0.43
1:B:155:PRO:C	1:B:157:PRO:HD2	2.39	0.43
1:A:271:ASP:OD2	2:A:612:HOH:O	2.21	0.43
1:B:135:ASP:OD1	1:B:137:ARG:HB2	2.18	0.43
1:B:51:PRO:HB3	2:B:558:HOH:O	2.18	0.42
1:A:127:LEU:HD11	1:A:138:MET:CE	2.49	0.42
1:A:109:ASP:OD1	1:A:128:ASN:HB3	2.19	0.42
1:A:184:ARG:NH1	1:A:184:ARG:HB2	2.35	0.42
1:B:38:THR:HG22	1:B:73:ALA:O	2.20	0.42
1:A:184:ARG:NH1	2:A:513:HOH:O	2.53	0.42
1:A:25:ASP:OD2	1:A:27:ASP:HB3	2.20	0.42
1:A:127:LEU:HD21	1:A:138:MET:HE3	2.00	0.41
1:A:88:GLU:OE1	1:A:115:VAL:HG21	2.20	0.41
1:B:87:GLU:HG2	1:B:90:ARG:HH12	1.84	0.41
1:A:261:MET:HG3	1:A:286:TYR:CE1	2.55	0.41
1:B:95:VAL:O	1:B:99:VAL:HG23	2.21	0.41
1:A:96:LEU:O	1:A:100:LEU:HG	2.21	0.41
1:A:152:MET:HE1	2:A:481:HOH:O	2.20	0.40
1:B:180:GLU:OE2	2:B:470:HOH:O	2.21	0.40
1:B:89:LYS:HE3	1:B:118:GLU:HG3	2.04	0.40
1:B:87:GLU:OE2	1:B:89:LYS:HB2	2.21	0.40
1:A:184:ARG:CB	1:A:184:ARG:HH11	2.34	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/294 (83%)	235 (96%)	9 (4%)	0	100	100
1	B	239/294 (81%)	227 (95%)	12 (5%)	0	100	100
All	All	483/588 (82%)	462 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/232 (85%)	193 (98%)	5 (2%)	55	26
1	B	197/232 (85%)	193 (98%)	4 (2%)	63	38
All	All	395/464 (85%)	386 (98%)	9 (2%)	58	30

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	GLU
1	A	90	ARG
1	A	128	ASN
1	A	129	ASP
1	A	286	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	22	ARG
1	B	91	ARG
1	B	128	ASN
1	B	156	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	HIS
1	A	270	HIS
1	B	36	ASN
1	B	250	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/294 (85%)	0.69	25 (10%) 9 7	13, 26, 55, 65	0
1	B	247/294 (84%)	0.94	45 (18%) 2 1	13, 31, 64, 77	1 (0%)
All	All	497/588 (84%)	0.81	70 (14%) 4 3	13, 28, 59, 77	1 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	VAL	12.7
1	A	39	PRO	11.0
1	B	156	VAL	11.0
1	B	53	ARG	8.9
1	B	39	PRO	8.7
1	B	85	VAL	8.1
1	B	86	GLU	6.3
1	A	38	THR	5.8
1	B	157	PRO	5.7
1	B	84	PRO	5.6
1	B	75	SER	5.2
1	A	37	LEU	5.0
1	A	55	LEU	4.9
1	A	90	ARG	4.5
1	A	53	ARG	4.3
1	B	56	GLU	4.2
1	B	52	GLU	4.2
1	A	54	ALA	4.1
1	B	155	PRO	4.0
1	A	56	GLU	4.0
1	B	164	ALA	4.0
1	B	91	ARG	3.9
1	A	286	TYR	3.8
1	B	87	GLU	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	74	GLU	3.6
1	A	91	ARG	3.6
1	B	59	ARG	3.2
1	B	121	LYS	3.2
1	B	55	LEU	3.2
1	B	165	HIS	3.2
1	B	102	LEU	3.2
1	A	36	ASN	3.1
1	B	93	LEU	3.0
1	B	88	GLU	3.0
1	A	227	LEU	3.0
1	B	126	LEU	2.9
1	B	228	VAL	2.9
1	B	18	TRP	2.9
1	B	227	LEU	2.9
1	A	59	ARG	2.9
1	A	86	GLU	2.7
1	B	97	GLU	2.7
1	B	54	ALA	2.7
1	A	87	GLU	2.7
1	B	94	PRO	2.7
1	B	101	SER	2.6
1	B	51	PRO	2.6
1	A	35	LEU	2.6
1	B	267	LEU	2.6
1	B	115	VAL	2.5
1	A	267	LEU	2.5
1	B	90	ARG	2.5
1	B	123	GLY	2.5
1	A	73	ALA	2.5
1	B	15	ARG	2.4
1	B	286	TYR	2.4
1	B	95	VAL	2.4
1	A	157	PRO	2.3
1	B	269	VAL	2.3
1	B	230	LEU	2.2
1	A	130	VAL	2.2
1	B	169	ARG	2.2
1	B	32	LEU	2.2
1	B	103	GLY	2.2
1	A	150	VAL	2.2
1	A	195	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	184	ARG	2.1
1	B	36	ASN	2.1
1	B	150	VAL	2.1
1	B	38	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.