



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:38 AM GMT

PDB ID : 2DQZ  
Title : Crystal structure of human carboxylesterase in complex with homatropine, coenzyme A, and palmitate  
Authors : Bencharit, S.; Redinbo, M.R.  
Deposited on : 2006-06-02  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	542	
1	B	542	
1	C	542	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	C	379	X	-	-	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	COA	C	1	-	X	-	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Liver carboxylesterase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			
1	B	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			
1	C	532	Total	C	N	O	S	0	0	0
			4130	2662	685	763	20			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLN	DELETION	UNP P23141
B	?	-	GLN	DELETION	UNP P23141
C	?	-	GLN	DELETION	UNP P23141

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is SUGAR (O-SIALIC ACID) (three-letter code: SIA) (formula:  $C_{11}H_{19}NO_9$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			21	11	1	9		
3	B	1	Total	C	N	O	0	0
			21	11	1	9		
3	C	1	Total	C	N	O	0	0
			21	11	1	9		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is FLUORIDE ION (three-letter code: F) (formula: F).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	F	0	0
			1	1		
5	A	1	Total	F	0	0
			1	1		
5	C	1	Total	F	0	0
			1	1		

- Molecule 6 is COENZYME A (three-letter code: COA) (formula: C<sub>21</sub>H<sub>36</sub>N<sub>7</sub>O<sub>16</sub>P<sub>3</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	C	1	Total	C	N	O	S	0	0
			7	4	1	1	1		

- Molecule 7 is PALMITIC ACID (three-letter code: PLM) (formula: C<sub>16</sub>H<sub>32</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			11	9	2		
7	B	1	Total	C	O	0	0
			11	9	2		
7	C	1	Total	C	O	0	0
			11	9	2		

- Molecule 8 is HOMOTROPINE (three-letter code: HTQ) (formula:  $C_{16}H_{21}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			20	16	1	3		
8	B	1	Total	C	N	O	0	0
			20	16	1	3		
8	C	1	Total	C	N	O	0	1
			40	32	2	6		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	179	Total	O	0	0
			179	179		
9	B	135	Total	O	0	0
			135	135		
9	C	194	Total	O	0	0
			194	194		




### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Liver carboxylesterase 1




- Molecule 1: Liver carboxylesterase 1



- Molecule 1: Liver carboxylesterase 1

Chain C:  70% 27% ..



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.56 Å   181.02 Å   202.56 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	22.60 – 2.80	Depositor
% Data completeness (in resolution range)	97.9 (22.60-2.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.193 , 0.244	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13161	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, F, COA, SIA, SO4, HTQ, PLM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/4236	0.58	0/5754
1	B	0.38	0/4236	0.56	0/5754
1	C	0.39	0/4236	0.58	0/5754
All	All	0.39	0/12708	0.57	0/17262

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4130	0	4131	125	0
1	B	4130	0	4131	132	0
1	C	4130	0	4130	111	0
2	A	14	0	13	1	0
2	B	14	0	13	2	0
2	C	14	0	13	1	0
3	A	21	0	18	5	0
3	B	21	0	18	6	0
3	C	21	0	18	4	0
4	A	10	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	15	0	0	0	0
4	C	10	0	0	1	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
6	C	7	0	6	0	0
7	A	11	0	14	0	0
7	B	11	0	14	2	0
7	C	11	0	14	0	0
8	A	20	0	21	2	0
8	B	20	0	21	0	0
8	C	40	0	42	0	0
9	A	179	0	0	17	0
9	B	135	0	0	15	0
9	C	194	0	0	23	0
All	All	13161	0	12617	379	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 379 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:3292:GLU:HB2	9:C:7340:HOH:O	1.52	1.10
1:A:1199:ARG:HH11	1:A:1199:ARG:HB3	1.22	1.02
1:C:3134:PRO:HG2	1:C:3163:VAL:HG12	1.45	0.96
1:C:3199:ARG:HB3	1:C:3199:ARG:HH11	1.33	0.93
1:A:1237:LYS:HE2	1:A:1342:HIS:HB2	1.55	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/542 (98%)	485 (92%)	40 (8%)	5 (1%)	21	55
1	B	530/542 (98%)	477 (90%)	47 (9%)	6 (1%)	17	50
1	C	530/542 (98%)	490 (92%)	36 (7%)	4 (1%)	24	58
All	All	1590/1626 (98%)	1452 (91%)	123 (8%)	15 (1%)	21	55

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2105	LYS
1	A	1320	GLY
1	B	2320	GLY
1	C	3105	LYS
1	C	3406	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/457 (98%)	437 (98%)	11 (2%)	55	86
1	B	448/457 (98%)	436 (97%)	12 (3%)	52	85
1	C	448/457 (98%)	440 (98%)	8 (2%)	66	91
All	All	1344/1371 (98%)	1313 (98%)	31 (2%)	58	88

5 of 31 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2264	LEU
1	B	2366	TYR
1	C	3264	LEU
1	B	2318	LEU
1	B	2372	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	2131	ASN
1	B	2202	GLN
1	B	2528	GLN
1	B	2069	GLN
1	B	2436	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 3 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	HTQ	A	11	-	22,22,22	2.84	14 (63%)	31,31,31	1.85	5 (16%)
7	PLM	A	111	-	7,10,17	0.75	0	7,10,17	0.74	0
2	NAG	A	179	1	14,14,15	0.61	0	15,19,21	0.64	0
3	SIA	A	182	-	17,21,21	1.05	1 (5%)	19,31,31	0.90	1 (5%)
4	SO4	A	184	-	4,4,4	0.55	0	6,6,6	0.12	0
4	SO4	A	284	-	4,4,4	0.28	0	6,6,6	0.11	0
7	PLM	B	112	-	7,10,17	0.74	0	7,10,17	0.73	0
8	HTQ	B	21	-	22,22,22	2.72	13 (59%)	31,31,31	1.80	6 (19%)
2	NAG	B	279	1	14,14,15	0.49	0	15,19,21	0.94	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SIA	B	282	-	17,21,21	0.71	0	19,31,31	1.22	1 (5%)
4	SO4	B	285	-	4,4,4	0.41	0	6,6,6	0.07	0
4	SO4	B	286	-	4,4,4	0.63	0	6,6,6	0.07	0
4	SO4	B	385	-	4,4,4	0.35	0	6,6,6	0.09	0
6	COA	C	1	-	6,6,50	4.04	2 (33%)	5,6,75	5.71	4 (80%)
7	PLM	C	113	-	7,10,17	0.87	0	7,10,17	0.75	0
4	SO4	C	185	-	4,4,4	0.46	0	6,6,6	0.06	0
8	HTQ	C	31[A]	-	22,22,22	2.66	13 (59%)	31,31,31	1.93	5 (16%)
8	HTQ	C	31[B]	-	22,22,22	2.67	13 (59%)	31,31,31	1.78	5 (16%)
2	NAG	C	379	1	14,14,15	0.79	0	15,19,21	0.63	0
3	SIA	C	382	-	17,21,21	0.77	1 (5%)	19,31,31	1.13	1 (5%)
4	SO4	C	384	-	4,4,4	0.43	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	HTQ	A	11	-	-	0/12/33/33	0/1/3/3
7	PLM	A	111	-	-	0/6/8/15	0/0/0/0
2	NAG	A	179	1	-	0/6/23/26	0/1/1/1
3	SIA	A	182	-	-	0/14/38/38	0/1/1/1
4	SO4	A	184	-	-	0/0/0/0	0/0/0/0
4	SO4	A	284	-	-	0/0/0/0	0/0/0/0
7	PLM	B	112	-	-	0/6/8/15	0/0/0/0
8	HTQ	B	21	-	-	0/12/33/33	0/1/3/3
2	NAG	B	279	1	-	0/6/23/26	0/1/1/1
3	SIA	B	282	-	-	0/14/38/38	0/1/1/1
4	SO4	B	285	-	-	0/0/0/0	0/0/0/0
4	SO4	B	286	-	-	0/0/0/0	0/0/0/0
4	SO4	B	385	-	-	0/0/0/0	0/0/0/0
6	COA	C	1	-	-	1/4/4/64	0/0/0/3
7	PLM	C	113	-	-	0/6/8/15	0/0/0/0
4	SO4	C	185	-	-	0/0/0/0	0/0/0/0
8	HTQ	C	31[A]	-	-	0/12/33/33	0/1/3/3
8	HTQ	C	31[B]	-	-	0/12/33/33	0/1/3/3
2	NAG	C	379	1	1/1/5/7	0/6/23/26	0/1/1/1
3	SIA	C	382	-	-	0/14/38/38	0/1/1/1
4	SO4	C	384	-	-	0/0/0/0	0/0/0/0

The worst 5 of 57 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	382	SIA	C6-C5	2.04	1.56	1.53
8	C	31[A]	HTQ	C5-N8	2.31	1.51	1.48
8	C	31[B]	HTQ	C5-N8	2.31	1.51	1.48
8	B	21	HTQ	O10-C11	2.32	1.39	1.34
8	C	31[A]	HTQ	C7-C1	2.34	1.59	1.53

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1	COA	O5P-C5P-N4P	-9.70	104.53	121.79
6	C	1	COA	O5P-C5P-C6P	-6.49	110.15	122.06
6	C	1	COA	C6P-C5P-N4P	-3.71	110.53	116.19
3	B	282	SIA	C7-C6-C5	-3.62	108.84	114.32
8	C	31[B]	HTQ	C15-C14-C13	-3.57	114.63	120.48

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	C	379	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	1	COA	O5P-C5P-N4P-C3P

There are no ring outliers.

9 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	11	HTQ	2	0
2	A	179	NAG	1	0
3	A	182	SIA	5	0
7	B	112	PLM	2	0
2	B	279	NAG	2	0
3	B	282	SIA	6	0
4	C	185	SO4	1	0
2	C	379	NAG	1	0
3	C	382	SIA	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.