



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:14 AM GMT

PDB ID : 3DUQ
Title : E(L212)A, D(L213)A, N(M5)D triple mutant structure of photosynthetic reaction center from Rhodobacter sphaeroides
Authors : Pokkuluri, P.R.; Schiffer, M.
Deposited on : 2008-07-17
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	281	<div> <div>2%</div> <div>69%</div> <div>27%</div> <div>• •</div> </div>
2	M	314	<div> <div>%</div> <div>67%</div> <div>24%</div> <div>• 5%</div> </div>
3	H	260	<div> <div>2%</div> <div>61%</div> <div>27%</div> <div>• 8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
10	CDL	M	800	-	-	-	X
4	BCL	L	501	X	-	-	-
4	BCL	M	501	X	-	-	X
5	BPH	M	503	X	-	-	-
6	U10	L	504	-	-	-	X
7	LDA	H	703	-	-	-	X
7	LDA	L	709	-	-	-	X
7	LDA	M	701	-	-	-	X
9	SPN	M	600	-	-	-	X

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 7261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reaction center protein L chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	281	Total	C	N	O	S	0	0	0
			2225	1504	355	358	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	212	ALA	GLU	ENGINEERED	UNP P0C0Y8
L	213	ALA	ASP	ENGINEERED	UNP P0C0Y8

- Molecule 2 is a protein called Reaction center protein M chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	298	Total	C	N	O	S	0	0	0
			2373	1585	388	390	10			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	5	ASP	ASN	ENGINEERED	UNP P0C0Y9
M	308	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	309	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	310	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	311	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	312	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	313	HIS	-	EXPRESSION TAG	UNP P0C0Y9
M	314	HIS	-	EXPRESSION TAG	UNP P0C0Y9

- Molecule 3 is a protein called Reaction center protein H chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	H	240	Total	C	N	O	S	0	0	0
			1829	1169	314	337	9			

- Molecule 4 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula: $C_{55}H_{74}MgN_4O_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	L	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		
4	M	1	Total	C	Mg	N	O	0	0
			66	55	1	4	6		

- Molecule 5 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula: $C_{55}H_{76}N_4O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	N	O	0	0
			65	55	4	6		
5	M	1	Total	C	N	O	10	0
			65	55	4	6		

- Molecule 6 is UBIQUINONE-10 (three-letter code: U10) (formula: $C_{59}H_{90}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total	C	O	0	0
			48	44	4		
6	M	1	Total	C	O	0	0
			48	44	4		

- Molecule 7 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula: $C_{14}H_{31}NO$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	L	1	Total	C	N	O	0	0
			16	14	1	1		
7	L	1	Total	C	N	O	0	0
			16	14	1	1		
7	M	1	Total	C	N	O	0	0
			16	14	1	1		
7	H	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 8 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	M	1	Total	Fe	0	0
			1	1		

- Molecule 9 is SPEROIDENONE (three-letter code: SPN) (formula: C₄₁H₇₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	M	1	Total	C	O	0	0
			43	41	2		

- Molecule 10 is CARDIOLIPIN (three-letter code: CDL) (formula: $C_{81}H_{156}O_{17}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	M	1	Total	C	O	P	0	0
			81	62	17	2		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	H	71	Total 71	O 71	0	0
11	L	36	Total 36	O 36	0	0
11	M	48	Total 48	O 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Reaction center protein L chain



• Molecule 2: Reaction center protein M chain



• Molecule 3: Reaction center protein H chain





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	141.30Å 141.30Å 187.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.70 29.12 – 2.70	Depositor EDS
% Data completeness (in resolution range)	93.0 (30.00-2.70) 93.3 (29.12-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.68Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.209 , 0.217 0.202 , 0.207	Depositor DCC
R_{free} test set	5668 reflections (11.29%)	DCC
Wilson B-factor (Å ²)	46.4	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 64.0	EDS
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 55879 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7261	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BCL, LDA, CDL, BPH, FE, SPN, U10

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.96	0/2313	1.10	14/3166 (0.4%)
2	M	1.07	3/2464 (0.1%)	1.15	19/3364 (0.6%)
3	H	0.95	2/1877 (0.1%)	1.28	22/2553 (0.9%)
All	All	1.00	5/6654 (0.1%)	1.17	55/9083 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	1
2	M	0	2
All	All	0	3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	76	PRO	C-N	-6.35	1.21	1.33
3	H	77	GLY	N-CA	5.74	1.54	1.46
2	M	34	PRO	C-N	-5.64	1.21	1.34
2	M	37	THR	C-N	-5.30	1.21	1.34
2	M	301	HIS	CA-C	-5.19	1.39	1.52

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	248	ARG	CD-NE-CZ	13.03	141.85	123.60
1	L	103	ARG	NE-CZ-NH1	10.98	125.79	120.30
3	H	250	SER	CA-C-O	-8.27	102.73	120.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	231	ARG	NE-CZ-NH2	7.48	124.04	120.30
2	M	164	ARG	NE-CZ-NH2	7.37	123.98	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	280	ASN	Mainchain
2	M	34	PRO	Mainchain
2	M	35	PHE	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	2225	0	2187	73	0
2	M	2373	0	2287	69	0
3	H	1829	0	1836	66	0
4	L	132	0	148	15	0
4	M	132	0	148	12	0
5	L	65	0	76	2	0
5	M	65	0	76	0	0
6	L	48	0	62	3	0
6	M	48	0	63	6	0
7	H	16	0	31	4	0
7	L	32	0	62	8	0
7	M	16	0	31	4	0
8	M	1	0	0	0	0
9	M	43	0	69	2	0
10	M	81	0	106	2	0
11	H	71	0	0	5	0
11	L	36	0	0	1	0
11	M	48	0	0	0	0
All	All	7261	0	7182	209	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:H:118:ARG:HH11	3:H:118:ARG:CG	1.81	0.94
3:H:33:THR:O	3:H:59:PRO:HG3	1.69	0.91
3:H:118:ARG:HH11	3:H:118:ARG:HG3	1.35	0.89
1:L:34:PHE:O	1:L:38:THR:HG23	1.71	0.89
6:M:504:U10:H202	7:H:703:LDA:H112	1.54	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	279/281 (99%)	266 (95%)	13 (5%)	0	100	100
2	M	296/314 (94%)	284 (96%)	10 (3%)	2 (1%)	26	55
3	H	238/260 (92%)	229 (96%)	9 (4%)	0	100	100
All	All	813/855 (95%)	779 (96%)	32 (4%)	2 (0%)	52	80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	M	22	GLU
2	M	301	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	218/218 (100%)	208 (95%)	10 (5%)	33	64
2	M	233/247 (94%)	227 (97%)	6 (3%)	54	83
3	H	195/208 (94%)	187 (96%)	8 (4%)	37	69
All	All	646/673 (96%)	622 (96%)	24 (4%)	41	72

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	136	ARG
2	M	204	LEU
3	H	231	ASP
2	M	144	LYS
2	M	196	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	280	ASN
2	M	188	ASN
3	H	68	HIS
3	H	206	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	LDA	H	703	-	15,15,15	4.78	4 (26%)	16,17,17	0.85	1 (6%)
4	BCL	L	501	1	53,74,74	1.20	4 (7%)	57,115,115	2.29	11 (19%)
4	BCL	L	502	1	53,74,74	1.37	5 (9%)	57,115,115	2.17	10 (17%)
5	BPH	L	503	-	64,70,70	1.31	9 (14%)	73,101,101	2.00	16 (21%)
6	U10	L	504	-	48,48,63	1.91	14 (29%)	58,61,79	2.98	17 (29%)
7	LDA	L	704	-	15,15,15	3.74	1 (6%)	16,17,17	0.60	0
7	LDA	L	709	-	15,15,15	4.57	2 (13%)	16,17,17	0.59	0
4	BCL	M	501	2	53,74,74	1.30	7 (13%)	57,115,115	2.35	16 (28%)
4	BCL	M	502	2	53,74,74	0.95	3 (5%)	57,115,115	1.77	8 (14%)
5	BPH	M	503	-	64,70,70	1.32	6 (9%)	73,101,101	2.30	24 (32%)
6	U10	M	504	-	48,48,63	2.62	21 (43%)	58,61,79	1.39	10 (17%)
9	SPN	M	600	-	41,42,42	4.32	20 (48%)	41,52,52	3.03	18 (43%)
7	LDA	M	701	-	15,15,15	4.89	2 (13%)	16,17,17	0.52	0
10	CDL	M	800	-	80,80,99	0.50	0	82,92,111	0.94	4 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	LDA	H	703	-	-	0/13/13/13	0/0/0/0
4	BCL	L	501	1	1/1/21/25	0/37/137/137	0/0/9/9
4	BCL	L	502	1	-	0/37/137/137	0/0/9/9
5	BPH	L	503	-	-	0/54/105/105	0/1/6/6
6	U10	L	504	-	-	0/45/69/87	0/1/1/1
7	LDA	L	704	-	-	0/13/13/13	0/0/0/0
7	LDA	L	709	-	-	0/13/13/13	0/0/0/0
4	BCL	M	501	2	2/2/21/25	1/37/137/137	0/0/9/9
4	BCL	M	502	2	-	0/37/137/137	0/0/9/9
5	BPH	M	503	-	1/1/18/22	0/54/105/105	0/1/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	U10	M	504	-	-	0/45/69/87	0/1/1/1
9	SPN	M	600	-	-	0/50/51/51	0/0/0/0
7	LDA	M	701	-	-	0/13/13/13	0/0/0/0
10	CDL	M	800	-	-	0/91/91/110	0/0/0/0

The worst 5 of 98 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	M	701	LDA	O1-N1	-18.70	1.21	1.39
7	H	703	LDA	O1-N1	-17.69	1.22	1.39
7	L	709	LDA	O1-N1	-17.45	1.22	1.39
7	L	704	LDA	O1-N1	-14.41	1.25	1.39
6	M	504	U10	C27-C28	-8.48	1.26	1.50

The worst 5 of 135 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	501	BCL	O1D-CGD-CBD	-8.05	113.08	124.62
4	M	501	BCL	C4-C3-C5	-6.61	105.31	115.41
9	M	600	SPN	C6-C5-C4	-5.50	110.62	121.05
4	M	502	BCL	O1D-CGD-CBD	-5.44	116.82	124.62
9	M	600	SPN	C17-C18-C19	-5.17	111.25	121.05

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	M	501	BCL	C8
4	M	501	BCL	C13
4	L	501	BCL	C13
5	M	503	BPH	C8

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	M	501	BCL	C1-C2-C3-C4

There are no ring outliers.

13 monomers are involved in 53 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	H	703	LDA	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	501	BCL	6	0
4	L	502	BCL	11	0
5	L	503	BPH	2	0
6	L	504	U10	3	0
7	L	704	LDA	4	0
7	L	709	LDA	4	0
4	M	501	BCL	5	0
4	M	502	BCL	7	0
6	M	504	U10	6	0
9	M	600	SPN	2	0
7	M	701	LDA	4	0
10	M	800	CDL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	281/281 (100%)	-0.33	7 (2%) 61 61	26, 41, 79, 100	0
2	M	298/314 (94%)	-0.38	4 (1%) 79 79	22, 46, 83, 102	0
3	H	240/260 (92%)	-0.40	4 (1%) 73 74	26, 43, 68, 111	0
All	All	819/855 (95%)	-0.37	15 (1%) 71 72	22, 44, 79, 111	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	250	SER	5.2
1	L	59	TRP	4.2
3	H	249	LYS	3.7
1	L	51	TRP	3.1
2	M	302	GLY	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	U10	L	504	48/63	0.70	0.47	10.35	65,87,116,119	0
10	CDL	M	800	81/100	0.84	0.51	5.76	25,68,84,87	81
7	LDA	L	709	16/16	0.74	0.54	5.10	94,99,111,111	0
7	LDA	M	701	16/16	0.89	0.20	4.64	55,66,76,79	0
9	SPN	M	600	43/43	0.82	0.28	3.12	44,62,83,87	0
7	LDA	H	703	16/16	0.90	0.27	2.88	63,70,78,82	0
4	BCL	M	501	66/66	0.95	0.18	2.76	25,36,104,105	0
6	U10	M	504	48/63	0.93	0.18	1.93	22,39,73,75	0
4	BCL	L	502	66/66	0.97	0.17	0.77	23,34,52,62	0
5	BPH	L	503	65/65	0.96	0.15	0.69	19,31,48,55	0
4	BCL	M	502	66/66	0.97	0.15	0.61	21,33,69,78	0
4	BCL	L	501	66/66	0.97	0.15	0.17	17,29,57,63	0
5	BPH	M	503	65/65	0.96	0.14	-0.10	31,45,72,78	10
8	FE	M	500	1/1	0.99	0.05	-2.26	26,26,26,26	0
7	LDA	L	704	16/16	0.68	0.31	-	84,92,119,120	0

6.5 Other polymers [i](#)

There are no such residues in this entry.