Full wwPDB X-ray Structure Validation Report (i)

Sep 15, 2016 – 05:44 PM EDT

PDB ID	:	5DUI
Title	:	Identification of a new FoxO1 binding site that precludes CREB binding at the glucose-6-phosphatase catalytic subunit gene promoter
Authors	:	Singh, P.; Endrizzi, J.A.; Chi, Y.-I.
Deposited on	:	2015-09-18
Resolution	:	2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	111	15%; 66%; 12%; 23%
1	B	111	5%; 74%; •; 23%
2	C	21	90%; 10%
3	D	21	81%; 19%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2274 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Forkhead box protein O1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	86	691	439	121	129	2	0	0	0
1	B	85	677	428	119	128	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	149	GLY	-	expression tag	UNP Q12778
A	150	SER	-	expression tag	UNP Q12778
B	149	GLY	-	expression tag	UNP Q12778
B	150	SER	-	expression tag	UNP Q12778

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*TP*TP*TP*CP*TP*AP*TP*TP*TP*TP*AP*CP*GP*TP*AP*AP*AP*TP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	C	21	422	207	66	129	20	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(*AP*TP*GP*AP*TP*TP*TP*AP*CP*GP*TP*AP*AP*AP*AP*TP*AP*GP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	D	21	433	209	85	119	20	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	13	13	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	B	14	14	14	0	0
			Total	O		
4	C	12	12	12	0	0
			Total	O		
4	D	12	12	12	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Forkhead box protein O1

Chain A:



- Molecule 1: Forkhead box protein O1

Chain B:



- Molecule 2: DNA (5'-D(*TP*TP*TP*TP*CP*TP*AP*TP*TP*TP*TP*AP*CP*GP*TP*AP*AP*AP*TP*CP*A)-3')

Chain C:

- Molecule 3: DNA (5'-D(*AP*TP*GP*AP*TP*TP*TP*AP*CP*GP*TP*AP*AP*AP*AP*TP*AP*GP*AP*AP*A)-3')

Chain D:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	46.61Å 79.03Å 48.27Å 90.00° 90.79° 90.00°	Depositor
Resolution (Å)	40.14 – 2.31 40.14 – 2.31	Depositor EDS
% Data completeness (in resolution range)	78.7 (40.14-2.31) 73.8 (40.14-2.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.14 (at 2.31Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.208 , 0.245 0.205 , 0.244	Depositor DCC
R_{free} test set	605 reflections (4.97%)	DCC
Wilson B-factor (Å^2)	41.1	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.33 , 42.2	EDS
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Estimated twinning fraction	0.008 for l,k,-h 0.048 for h,-k,-l 0.196 for l,-k,h	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	2274	wwPDB-VP
Average B, all atoms (Å^2)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.23	0/709	0.40	0/955
1	B	0.22	0/693	0.38	0/932
2	C	0.45	0/470	1.18	0/723
3	D	0.43	0/488	1.01	0/752
All	All	0.33	0/2360	0.78	0/3362

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	691	0	676	8	0
1	B	677	0	666	1	0
2	C	422	0	244	2	0
3	D	433	0	239	2	0
4	A	13	0	0	0	0
4	B	14	0	0	0	0
4	C	12	0	0	0	0
4	D	12	0	0	0	0
All	All	2274	0	1825	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:SER:O	1:A:210:LYS:NZ	2.33	0.59
1:B:199:ASP:OD2	1:B:199:ASP:N	2.37	0.53
1:A:160:TRP:NE1	1:A:168:LEU:HD11	2.23	0.53
1:A:160:TRP:HE1	1:A:168:LEU:HD11	1.75	0.51
1:A:206:SER:HA	1:A:207:ALA:O	2.11	0.50
2:C:2:DT:H2"	2:C:3:DT:O5'	2.12	0.50
3:D:11:DT:H2"	3:D:12:DA:C8	2.49	0.47
3:D:9:DC:H2"	3:D:10:DG:C8	2.51	0.46
1:A:169:ILE:HG21	1:A:217:LEU:HD21	2.00	0.42
1:A:206:SER:HA	1:A:207:ALA:C	2.41	0.41
2:C:2:DT:H2'	2:C:3:DT:C6	2.55	0.41
1:A:196:TYR:O	1:A:200:LYS:NZ	2.54	0.40
1:A:191:VAL:O	1:A:198:LYS:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	84/111 (76%)	79 (94%)	5 (6%)	0	100 100
1	B	83/111 (75%)	79 (95%)	3 (4%)	1 (1%)	16 16
All	All	167/222 (75%)	158 (95%)	8 (5%)	1 (1%)	30 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	233	LYS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	75/95 (79%)	74 (99%)	1 (1%)	76	87
1	B	74/95 (78%)	73 (99%)	1 (1%)	74	86
All	All	149/190 (78%)	147 (99%)	2 (1%)	76	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	229	GLU
1	B	202	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	86/111 (77%)	1.25	17 (19%) 1 2	35, 60, 124, 155	0
1	B	85/111 (76%)	0.65	5 (5%) 26 34	31, 58, 113, 126	0
2	C	21/21 (100%)	-0.34	0 100 100	40, 63, 88, 92	0
3	D	21/21 (100%)	-0.17	0 100 100	42, 71, 105, 125	0
All	All	213/264 (80%)	0.71	22 (10%) 9 13	31, 61, 122, 155	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	201	GLY	12.3
1	A	244	GLY	5.2
1	A	232	GLY	5.1
1	A	199	ASP	4.9
1	A	231	THR	4.7
1	B	198	LYS	3.9
1	A	196	TYR	3.8
1	A	160	TRP	3.7
1	B	231	THR	3.3
1	A	197	PHE	3.3
1	A	198	LYS	3.2
1	A	206	SER	3.2
1	A	200	LYS	3.1
1	A	202	ASP	3.0
1	B	233	LYS	2.7
1	A	205	SER	2.6
1	B	244	GLY	2.5
1	A	243	GLY	2.4
1	B	203	SER	2.4
1	A	163	LEU	2.4
1	A	233	LYS	2.2
1	A	203	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.