



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 1, 2016 – 07:32 PM EDT

PDB ID : 3DZG
Title : Crystal structure of human CD38 extracellular domain, ara-F-ribose-5'-phosphate/nicotinamide complex
Authors : Liu, Q.; Kriksunov, I.A.; Jiang, H.; Graeff, R.; Lin, H.; Lee, H.C.; Hao, Q.
Deposited on : 2008-07-29
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1226 (1.66-1.66)
Clashscore	102246	1323 (1.66-1.66)
Ramachandran outliers	100387	1295 (1.66-1.66)
Sidechain outliers	100360	1295 (1.66-1.66)
RSRZ outliers	91569	1227 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	
1	B	262	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4612 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ADP-ribosyl cyclase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	252	Total	C	N	O	S	0	0	0
			2050	1290	358	386	16			
1	B	252	Total	C	N	O	S	0	0	0
			2050	1290	358	386	16			

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	39	LYS	-	EXPRESSION TAG	UNP P28907
A	40	ARG	-	EXPRESSION TAG	UNP P28907
A	41	GLU	-	EXPRESSION TAG	UNP P28907
A	42	ALA	-	EXPRESSION TAG	UNP P28907
A	43	GLU	-	EXPRESSION TAG	UNP P28907
A	44	ALA	-	EXPRESSION TAG	UNP P28907
A	49	THR	GLN	ENGINEERED	UNP P28907
A	100	ASP	ASN	ENGINEERED	UNP P28907
A	164	ASP	ASN	ENGINEERED	UNP P28907
A	209	ASP	ASN	ENGINEERED	UNP P28907
A	219	ASP	ASN	ENGINEERED	UNP P28907
B	39	LYS	-	EXPRESSION TAG	UNP P28907
B	40	ARG	-	EXPRESSION TAG	UNP P28907
B	41	GLU	-	EXPRESSION TAG	UNP P28907
B	42	ALA	-	EXPRESSION TAG	UNP P28907
B	43	GLU	-	EXPRESSION TAG	UNP P28907
B	44	ALA	-	EXPRESSION TAG	UNP P28907
B	49	THR	GLN	ENGINEERED	UNP P28907
B	100	ASP	ASN	ENGINEERED	UNP P28907
B	164	ASP	ASN	ENGINEERED	UNP P28907
B	209	ASP	ASN	ENGINEERED	UNP P28907
B	219	ASP	ASN	ENGINEERED	UNP P28907

- Molecule 2 is SUGAR (2-DEOXY-2-FLUORO-5-O-PHOSPHONO-ALPHA-D-ARABINOF URANOSE) (three-letter code: RF5) (formula: C₅H₁₀FO₇P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	O	P	0	0
			13	5	1	6	1		
2	B	1	Total	C	F	O	P	0	0
			13	5	1	6	1		

- Molecule 3 is NICOTINAMIDE (three-letter code: NCA) (formula: $C_6H_6N_2O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	6	2	1		
3	B	1	Total	C	N	O	0	0
			9	6	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	236	Total 236	O 236	0	0
4	B	232	Total 232	O 232	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADP-ribosyl cyclase 1



• Molecule 1: ADP-ribosyl cyclase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.69 Å 52.81 Å 65.38 Å 106.06° 91.95° 95.12°	Depositor
Resolution (Å)	20.00 – 1.65 30.12 – 1.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.65) 87.2 (30.12-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.12 (at 1.64 Å)	Xtriage
Refinement program	REFMAC 5.3.0021	Depositor
R, R_{free}	0.174 , 0.211 0.262 , 0.287	Depositor DCC
R_{free} test set	3125 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.373	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4612	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NCA, RF5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	3/2101 (0.1%)	0.77	2/2846 (0.1%)
1	B	0.59	0/2101	0.67	0/2846
All	All	0.66	3/4202 (0.1%)	0.72	2/5692 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	72	GLU	CD-OE2	13.79	1.40	1.25
1	A	68	VAL	CA-CB	5.68	1.66	1.54
1	A	72	GLU	CD-OE1	5.51	1.31	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	VAL	CA-CB-CG1	-7.06	100.31	110.90
1	A	131	LEU	CB-CG-CD2	5.37	120.13	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2050	0	1976	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2050	0	1976	34	0
2	A	13	0	6	0	0
2	B	13	0	6	0	0
3	A	9	0	6	0	0
3	B	9	0	6	0	0
4	A	236	0	0	6	0
4	B	232	0	0	10	0
All	All	4612	0	3976	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:194:ARG:HD2	4:B:541:HOH:O	1.67	0.94
1:B:115:GLN:HE22	1:B:149:LEU:H	1.18	0.92
1:A:115:GLN:HE22	1:A:149:LEU:H	1.18	0.90
1:A:165:THR:HG23	1:A:167:LYS:H	1.36	0.88
1:B:165:THR:HG23	1:B:167:LYS:H	1.35	0.87
1:B:266:ILE:HD11	1:B:273:PHE:HB2	1.60	0.82
1:B:202:ASP:HB3	4:B:503:HOH:O	1.80	0.80
1:A:75:PRO:HA	1:A:78:ARG:HG3	1.67	0.77
1:A:195:ARG:HD3	4:A:531:HOH:O	1.84	0.75
1:B:90:LYS:HG3	1:B:94:ILE:HG13	1.70	0.73
1:B:236:GLN:NE2	4:B:630:HOH:O	2.27	0.68
1:A:127:ARG:NH1	1:A:217:ASP:OD2	2.21	0.67
1:A:294:SER:O	1:A:296:CYS:N	2.28	0.67
1:B:266:ILE:HD11	1:B:273:PHE:CB	2.25	0.65
1:A:183:ASN:ND2	1:A:186:SER:H	1.95	0.65
1:A:268:LYS:HD3	1:B:163:PHE:CE1	2.32	0.65
1:A:140:ARG:NH2	4:A:431:HOH:O	2.29	0.64
1:B:162:GLU:HB2	1:B:165:THR:HG22	1.79	0.64
1:A:162:GLU:HB2	1:A:165:THR:HG22	1.80	0.63
1:A:182:ASN:HB2	4:A:413:HOH:O	1.99	0.62
1:A:86:TRP:CE2	1:A:90:LYS:HG3	2.35	0.61
1:A:110:MET:HE1	1:A:150:LEU:HD13	1.82	0.61
1:B:137:GLN:HG2	4:B:627:HOH:O	2.02	0.60
1:A:127:ARG:HB3	1:A:212:ARG:HE	1.66	0.60
1:A:268:LYS:HD3	1:B:163:PHE:HE1	1.66	0.58
1:B:115:GLN:NE2	1:B:149:LEU:H	1.96	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:GLU:HA	1:B:107:GLN:HG2	1.88	0.56
1:A:46:TRP:HE1	1:A:47:ARG:NH2	2.04	0.56
1:B:244:HIS:HD2	1:B:250:SER:HB3	1.70	0.55
1:A:48:GLN:NE2	1:A:171:GLN:HB3	2.22	0.55
1:B:195:ARG:HB2	4:B:528:HOH:O	2.07	0.54
1:B:45:ARG:HG3	1:B:46:TRP:N	2.23	0.54
1:A:78:ARG:NH1	4:A:395:HOH:O	2.41	0.53
1:B:194:ARG:NH1	4:B:634:HOH:O	2.41	0.53
1:B:195:ARG:HG3	4:B:527:HOH:O	2.09	0.52
1:B:293:ASP:CG	1:B:294:SER:H	2.13	0.52
1:B:244:HIS:CD2	1:B:250:SER:HB3	2.44	0.52
1:B:47:ARG:NH2	4:B:639:HOH:O	2.41	0.52
1:B:266:ILE:HD11	1:B:273:PHE:CA	2.40	0.52
1:A:244:HIS:HE1	1:A:277:ASN:OD1	1.94	0.50
1:A:183:ASN:HD21	1:A:186:SER:H	1.58	0.49
1:A:243:ILE:HG23	1:A:281:PRO:HG3	1.95	0.49
1:B:90:LYS:CG	1:B:94:ILE:HG13	2.39	0.49
1:B:238:LEU:HB3	1:B:266:ILE:HD13	1.95	0.47
1:A:134:GLN:CD	4:A:444:HOH:O	2.53	0.47
1:A:76:GLU:HG2	1:A:77:MET:HG2	1.97	0.46
1:B:98:PRO:O	1:B:183:ASN:HA	2.15	0.46
1:B:45:ARG:HG3	1:B:47:ARG:H	1.81	0.46
1:B:195:ARG:HD3	4:B:528:HOH:O	2.14	0.46
1:B:248:GLU:HG3	1:B:249:ASP:H	1.81	0.45
1:A:115:GLN:NE2	1:A:149:LEU:H	1.99	0.45
1:A:98:PRO:O	1:A:183:ASN:HA	2.18	0.44
1:B:134:GLN:O	1:B:137:GLN:HB2	2.18	0.43
1:A:104:GLU:HA	1:A:107:GLN:HG2	2.00	0.43
1:A:291:PRO:O	1:A:292:GLU:HB3	2.18	0.43
1:B:199:ALA:HB3	4:B:591:HOH:O	2.19	0.42
1:B:162:GLU:HB2	1:B:165:THR:CG2	2.47	0.42
1:A:94:ILE:O	1:A:95:SER:HB2	2.20	0.42
1:A:128:ILE:HG13	4:A:418:HOH:O	2.19	0.42
1:A:243:ILE:CG2	1:A:281:PRO:HG3	2.49	0.41
1:B:73:ILE:O	1:B:75:PRO:HD3	2.21	0.41
1:B:47:ARG:HH11	1:B:47:ARG:HG2	1.85	0.41
1:A:110:MET:HE1	1:A:192:VAL:HG12	2.02	0.41
1:A:78:ARG:HB2	1:A:78:ARG:HE	1.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/262 (95%)	240 (96%)	8 (3%)	2 (1%)	24	5
1	B	250/262 (95%)	239 (96%)	8 (3%)	3 (1%)	16	2
All	All	500/524 (95%)	479 (96%)	16 (3%)	5 (1%)	19	3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	295	SER
1	B	291	PRO
1	B	295	SER
1	A	128	ILE
1	B	128	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/241 (97%)	229 (98%)	4 (2%)	68	45
1	B	233/241 (97%)	228 (98%)	5 (2%)	61	34
All	All	466/482 (97%)	457 (98%)	9 (2%)	65	40

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	89	PHE
1	A	131	LEU
1	A	292	GLU
1	B	89	PHE
1	B	131	LEU
1	B	236	GLN
1	B	251	ARG
1	B	286	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	GLN
1	A	115	GLN
1	A	134	GLN
1	A	139	GLN
1	A	183	ASN
1	A	244	HIS
1	B	115	GLN
1	B	134	GLN
1	B	229	ASN
1	B	244	HIS
1	B	270	ASN
1	B	286	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	RF5	A	301	-	13,13,14	0.75	0	16,19,21	1.21	1 (6%)
3	NCA	A	302	-	9,9,9	0.56	0	11,11,11	2.06	4 (36%)
2	RF5	B	301	-	13,13,14	0.75	0	16,19,21	1.17	0
3	NCA	B	302	-	9,9,9	0.43	0	11,11,11	1.66	4 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RF5	A	301	-	-	0/6/19/22	0/1/1/1
3	NCA	A	302	-	-	0/4/4/4	0/1/1/1
2	RF5	B	301	-	-	0/6/19/22	0/1/1/1
3	NCA	B	302	-	-	0/4/4/4	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	NCA	O7-C7-C3	-3.40	115.82	119.60
3	A	302	NCA	C3-C2-N1	-3.23	118.59	123.47
3	B	302	NCA	C3-C2-N1	-2.73	119.35	123.47
3	B	302	NCA	O7-C7-C3	-2.34	117.00	119.60
2	A	301	RF5	O3-C3-C4	2.05	117.12	111.01
3	B	302	NCA	C3-C7-N7	2.48	120.62	117.82
3	B	302	NCA	C6-N1-C2	2.66	121.71	116.82
3	A	302	NCA	C6-N1-C2	2.73	121.84	116.82
3	A	302	NCA	C4-C3-C2	2.87	120.94	117.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.







5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	252/262 (96%)	1.57	72 (28%)  	24, 30, 39, 48	0
1	B	252/262 (96%)	1.81	90 (35%)  	26, 31, 39, 43	0
All	All	504/524 (96%)	1.69	162 (32%)  	24, 30, 39, 48	0

All (162) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	ASP	9.9
1	A	249	ASP	9.5
1	B	247	ARG	9.0
1	B	246	GLY	8.8
1	B	291	PRO	8.5
1	B	213	SER	8.2
1	B	249	ASP	7.8
1	A	246	GLY	7.7
1	B	295	SER	7.1
1	B	248	GLU	6.9
1	B	290	ASN	6.7
1	B	294	SER	6.7
1	A	294	SER	6.6
1	B	245	GLY	6.5
1	A	292	GLU	6.4
1	A	248	GLU	6.4
1	A	124	LEU	6.2
1	B	292	GLU	6.1
1	B	287	CYS	6.1
1	B	296	CYS	6.1
1	B	293	ASP	6.0
1	B	282	ASP	5.7
1	A	47	ARG	5.6
1	B	286	GLN	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	LEU	5.5
1	A	213	SER	5.2
1	B	159	TRP	5.2
1	A	247	ARG	5.2
1	B	153	LEU	5.1
1	B	189	TRP	5.1
1	B	280	ARG	4.8
1	B	284	PHE	4.7
1	A	230	LEU	4.6
1	A	123	LEU	4.6
1	A	189	TRP	4.5
1	A	222	PHE	4.5
1	B	211	SER	4.5
1	B	150	LEU	4.4
1	A	204	VAL	4.4
1	A	238	LEU	4.4
1	B	185	VAL	4.4
1	A	46	TRP	4.3
1	B	123	LEU	4.3
1	A	192	VAL	4.3
1	B	188	PHE	4.2
1	B	285	LEU	4.2
1	B	250	SER	4.1
1	A	45	ARG	4.1
1	A	196	PHE	4.1
1	B	127	ARG	4.0
1	B	228	HIS	4.0
1	B	141	ASP	4.0
1	A	206	VAL	3.9
1	A	225	VAL	3.9
1	A	227	VAL	3.9
1	A	185	VAL	3.9
1	B	116	THR	3.8
1	B	244	HIS	3.7
1	A	164	ASP	3.7
1	A	266	ILE	3.7
1	B	145	LEU	3.7
1	A	84	SER	3.7
1	B	138	VAL	3.7
1	A	79	HIS	3.7
1	A	245	GLY	3.6
1	A	296	CYS	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	89	PHE	3.6
1	A	291	PRO	3.6
1	B	124	LEU	3.6
1	B	289	LYS	3.5
1	A	150	LEU	3.5
1	A	48	GLN	3.5
1	B	269	ARG	3.5
1	A	188	PHE	3.4
1	B	151	GLY	3.4
1	B	215	ILE	3.4
1	B	196	PHE	3.4
1	A	193	SER	3.4
1	A	125	TRP	3.4
1	A	73	ILE	3.3
1	A	295	SER	3.3
1	A	271	ILE	3.3
1	A	159	TRP	3.3
1	A	235	VAL	3.3
1	B	212	ARG	3.3
1	B	176	TRP	3.3
1	B	252	ASP	3.2
1	A	144	THR	3.2
1	B	251	ARG	3.1
1	A	148	THR	3.1
1	B	219	ASP	3.0
1	A	80	VAL	3.0
1	B	184	PRO	3.0
1	A	143	PHE	3.0
1	B	93	PHE	3.0
1	B	279	TYR	2.9
1	B	149	LEU	2.9
1	B	257	PRO	2.9
1	B	288	VAL	2.9
1	A	122	ILE	2.9
1	A	154	ALA	2.9
1	A	195	ARG	2.9
1	B	137	GLN	2.8
1	B	224	SER	2.8
1	A	58	ARG	2.8
1	A	262	LEU	2.8
1	B	148	THR	2.8
1	A	153	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	144	THR	2.7
1	A	187	VAL	2.7
1	A	197	ALA	2.7
1	B	214	LYS	2.7
1	B	255	GLN	2.7
1	A	78	ARG	2.7
1	B	281	PRO	2.7
1	B	142	MET	2.6
1	A	205	HIS	2.6
1	B	154	ALA	2.6
1	B	270	ASN	2.6
1	B	283	LYS	2.6
1	A	280	ARG	2.6
1	B	260	LYS	2.6
1	B	143	PHE	2.5
1	B	140	ARG	2.5
1	B	63	VAL	2.5
1	B	187	VAL	2.5
1	B	192	VAL	2.5
1	A	111	LYS	2.5
1	B	173	CYS	2.5
1	B	253	LEU	2.5
1	B	256	ASP	2.5
1	B	99	CYS	2.5
1	A	200	ALA	2.4
1	B	106	TYR	2.4
1	B	267	SER	2.3
1	B	258	THR	2.3
1	A	203	VAL	2.3
1	A	208	LEU	2.3
1	B	55	THR	2.3
1	A	69	LYS	2.3
1	B	160	CYS	2.3
1	B	50	TRP	2.3
1	A	265	ILE	2.3
1	A	68	VAL	2.2
1	A	136	THR	2.2
1	A	151	GLY	2.2
1	B	254	CYS	2.2
1	B	278	ILE	2.2
1	B	218	LYS	2.1
1	B	233	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	65	ALA	2.1
1	A	207	MET	2.1
1	B	54	GLY	2.1
1	B	259	ILE	2.1
1	A	49	THR	2.1
1	B	158	THR	2.1
1	A	221	THR	2.0
1	B	157	LEU	2.0
1	B	180	CYS	2.0
1	A	165	THR	2.0
1	A	191	THR	2.0
1	B	174	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NCA	B	302	9/9	0.94	0.12	-1.25	19,24,25,26	0
2	RF5	B	301	13/14	0.95	0.09	-1.30	22,26,30,33	0
3	NCA	A	302	9/9	0.92	0.10	-1.80	19,23,25,26	0
2	RF5	A	301	13/14	0.97	0.07	-2.64	14,18,19,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.