



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:07 PM GMT

PDB ID : 1E1Q
Title : BOVINE MITOCHONDRIAL F1-ATPASE AT 100K
Authors : Braig, K.; Menz, R.I.; Montgomery, M.G.; Leslie, A.G.W.; Walker, J.E.
Deposited on : 2000-05-10
Resolution : 2.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3065 (2.64-2.60)
Ramachandran outliers	100387	3015 (2.64-2.60)
Sidechain outliers	100360	3015 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	510	
1	B	510	
1	C	510	
2	D	482	
2	E	482	
2	F	482	
3	G	272	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 23366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	0	0	0
			3715	2341	656	706	12			
1	B	479	Total	C	N	O	S	0	0	0
			3656	2303	647	694	12			
1	C	492	Total	C	N	O	S	0	0	0
			3748	2360	661	715	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	481	GLY	SER	CONFLICT	UNP P19483
B	481	GLY	SER	CONFLICT	UNP P19483
C	481	GLY	SER	CONFLICT	UNP P19483


- Molecule 2 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	467	Total	C	N	O	S	0	0	0
			3539	2243	601	684	11			
2	E	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			
2	F	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			

- Molecule 3 is a protein called BOVINE MITOCHONDRIAL F1-ATPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	122	Total	C	N	O	S	0	0	0
			945	591	171	176	7			

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	B	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	C	1	Total 31	C 10	N 6	O 12	P 3	0	0
4	F	1	Total 31	C 10	N 6	O 12	P 3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0
5	D	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0
5	F	1	Total Mg 1 1	0	0

- Molecule 6 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $\text{C}_{10}\text{H}_{15}\text{N}_5\text{O}_{10}\text{P}_2$).



-
- PO4
- 03
- O⁻
- 02 O⁻ — P — O 01
- P
- O⁻
- 04

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	E	1	Total	O	P	0	0
			5	4	1		

- 


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	100	Total 100	O 100	0	0
8	B	83	Total 83	O 83	0	0
8	C	109	Total 109	O 109	0	0
8	D	92	Total 92	O 92	0	0
8	E	44	Total 44	O 44	0	0
8	F	107	Total 107	O 107	0	0
8	G	7	Total 7	O 7	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: BOVINE MITOCHONDRIAL F1-ATPASE




• Molecule 1: BOVINE MITOCHONDRIAL F1-ATPASE






• Molecule 1: BOVINE MITOCHONDRIAL F1-ATPASE




• Molecule 2: BOVINE MITOCHONDRIAL F1-ATPASE




• Molecule 2: BOVINE MITOCHONDRIAL F1-ATPASE






• Molecule 2: BOVINE MITOCHONDRIAL F1-ATPASE

Chain F: 66% 27%



• Molecule 3: BOVINE MITOCHONDRIAL F1-ATPASE

Chain G: 30% 13% 55%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	280.80 Å 107.40 Å 139.60 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.61	Depositor
% Data completeness (in resolution range)	95.0 (20.00-2.61)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.232 , 0.280	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	23366	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ANP, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/3766	1.24	28/5080 (0.6%)
1	B	0.60	0/3704	1.24	23/4995 (0.5%)
1	C	0.63	0/3799	1.33	28/5126 (0.5%)
2	D	0.59	0/3596	1.28	26/4879 (0.5%)
2	E	0.55	0/3587	1.18	20/4867 (0.4%)
2	F	0.63	0/3587	1.29	26/4867 (0.5%)
3	G	0.45	0/949	0.97	4/1266 (0.3%)
All	All	0.59	0/22988	1.25	155/31080 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 155 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	274	ARG	NE-CZ-NH1	14.94	127.77	120.30
1	C	304	ARG	NE-CZ-NH2	13.00	126.80	120.30
1	C	423	ARG	NE-CZ-NH2	-12.63	113.98	120.30
1	C	219	ARG	NE-CZ-NH2	-12.04	114.28	120.30
1	A	210	ARG	CD-NE-CZ	11.64	139.89	123.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	97	VAL	Mainchain
2	E	223	ASN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3814	136	0
1	B	3656	0	3765	142	0
1	C	3748	0	3845	125	0
2	D	3539	0	3592	119	0
2	E	3530	0	3587	145	0
2	F	3530	0	3586	94	0
3	G	945	0	1019	30	0
4	A	31	0	13	1	0
4	B	31	0	13	2	0
4	C	31	0	13	2	0
4	F	31	0	13	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	F	1	0	0	0	0
6	D	27	0	12	1	0
7	E	5	0	0	1	0
8	A	100	0	0	6	0
8	B	83	0	0	6	0
8	C	109	0	0	3	0
8	D	92	0	0	4	0
8	E	44	0	0	4	0
8	F	107	0	0	3	0
8	G	7	0	0	0	0
All	All	23366	0	23272	741	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 741 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:282:GLN:H	2:D:282:GLN:HE21	0.96	0.96
1:C:294:TYR:HB3	1:C:298:VAL:HG21	1.52	0.91
2:D:139:VAL:HB	8:D:2027:HOH:O	1.72	0.90
1:C:187:LYS:HE3	1:C:224:ASP:HB3	1.54	0.89
2:D:145:PRO:HB2	2:D:357:ILE:HD11	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/510 (95%)	452 (93%)	25 (5%)	8 (2%)	12	23
1	B	475/510 (93%)	434 (91%)	38 (8%)	3 (1%)	30	54
1	C	490/510 (96%)	457 (93%)	28 (6%)	5 (1%)	19	37
2	D	465/482 (96%)	429 (92%)	32 (7%)	4 (1%)	21	41
2	E	464/482 (96%)	419 (90%)	40 (9%)	5 (1%)	17	34
2	F	464/482 (96%)	431 (93%)	31 (7%)	2 (0%)	39	63
3	G	116/272 (43%)	107 (92%)	6 (5%)	3 (3%)	7	10
All	All	2959/3248 (91%)	2729 (92%)	200 (7%)	30 (1%)	19	37

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	409	ASP
2	E	211	ALA
1	B	452	TYR
1	C	337	TYR
1	C	388	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/412 (95%)	365 (93%)	28 (7%)	18	35
1	B	388/412 (94%)	371 (96%)	17 (4%)	35	62
1	C	397/412 (96%)	368 (93%)	29 (7%)	17	34
2	D	377/386 (98%)	358 (95%)	19 (5%)	30	55
2	E	376/386 (97%)	355 (94%)	21 (6%)	26	49
2	F	376/386 (97%)	359 (96%)	17 (4%)	34	61
3	G	102/230 (44%)	97 (95%)	5 (5%)	31	56
All	All	2409/2624 (92%)	2273 (94%)	136 (6%)	26	49

5 of 136 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	349	GLN
2	D	44	ARG
2	F	232	VAL
1	C	390	MET
1	C	470	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	282	GLN
2	E	263	GLN
2	F	282	GLN
2	E	223	ASN
2	E	246	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	ANP	A	600	5	27,33,33	1.53	6 (22%)	30,52,52	1.66	6 (20%)
4	ANP	B	600	5	27,33,33	1.72	7 (25%)	30,52,52	2.70	6 (20%)
4	ANP	C	600	5	27,33,33	1.63	6 (22%)	30,52,52	1.52	8 (26%)
6	ADP	D	600	5	22,29,29	0.97	2 (9%)	27,45,45	1.39	3 (11%)
7	PO4	E	602	-	4,4,4	0.29	0	6,6,6	0.27	0
4	ANP	F	600	5	27,33,33	1.63	6 (22%)	30,52,52	1.39	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ANP	A	600	5	-	1/12/38/38	0/3/3/3
4	ANP	B	600	5	-	2/12/38/38	0/3/3/3
4	ANP	C	600	5	-	0/12/38/38	0/3/3/3
6	ADP	D	600	5	-	0/12/32/32	0/3/3/3
7	PO4	E	602	-	-	0/0/0/0	0/0/0/0
4	ANP	F	600	5	-	2/12/38/38	0/3/3/3

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	600	ANP	PG-O3G	-4.15	1.45	1.56
4	C	600	ANP	PG-O3G	-3.83	1.46	1.56
4	C	600	ANP	PB-O3A	-3.66	1.54	1.59
4	F	600	ANP	PG-O3G	-3.53	1.46	1.56
4	A	600	ANP	PG-O3G	-3.30	1.47	1.56

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	600	ANP	C2'-C1'-N9	-3.96	108.24	114.29
4	B	600	ANP	O2G-PG-O1G	-3.96	102.97	113.49
6	D	600	ADP	C1'-N9-C4	-3.52	121.64	126.94
4	F	600	ANP	N6-C6-N1	-3.41	111.89	119.20
6	D	600	ADP	C2'-C1'-N9	-3.21	109.39	114.29

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	600	ANP	O1G-PG-N3B-PB
4	F	600	ANP	O1B-PB-N3B-PG
4	F	600	ANP	O1G-PG-N3B-PB
4	B	600	ANP	O1B-PB-N3B-PG
4	B	600	ANP	O1G-PG-N3B-PB

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	600	ANP	1	0
4	B	600	ANP	2	0
4	C	600	ANP	2	0
6	D	600	ADP	1	0
7	E	602	PO4	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.