



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 07:07 PM GMT

PDB ID : 1E3Z  
Title : ACARBOSE COMPLEX OF CHIMAERIC AMYLASE FROM B. AMY-  
LOLIQUEFACIENS AND B. LICHENIFORMIS AT 1.93Å  
Authors : Brzozowski, A.M.; Lawson, D.M.; Turkenburg, J.P.; Bisgaard-Frantzen, H.;  
Svendsen, A.; Borchert, T.V.; Dauter, Z.; Wilson, K.S.; Davies, G.J.  
Deposited on : 2000-06-27  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3095 (1.96-1.92)
Ramachandran outliers	100387	3062 (1.96-1.92)
Sidechain outliers	100360	3062 (1.96-1.92)
RSRZ outliers	91569	2915 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	<div> <div></div> <div>89%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GLC	A	810	X	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-AMYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	483	Total	C	N	O	S	9	16	0
			3948	2500	682	759	7			


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is SUGAR (6-AMINO-4-HYDROXYMETHYL-CYCLOHEX-4-ENE-1,2,3-TRIOL) (three-letter code: ACI) (formula: C<sub>7</sub>H<sub>13</sub>NO<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	7	1	4		

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	3	Total	C	N	O	0	0
			32	19	1	12		
5	A	3	Total	C	N	O	0	0
			32	19	1	12		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	3	Total	C	O	0	0
			32	18	14		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	680	Total	O	3	0
			680	680		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALPHA-AMYLASE

Chain A:  89% 11%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.72Å 78.27Å 238.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.93 29.44 – 1.92	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-1.93) 98.9 (29.44-1.92)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.23 (at 1.92Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.130 , 0.200 0.127 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	13.2	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 55.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 38068 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4741	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ACI, GLC, GLD, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	1/4144 (0.0%)	1.14	20/5619 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	A	1	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	214	LYS	CG-CD	7.79	1.78	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	437	ARG	CD-NE-CZ	10.35	138.09	123.60
1	A	167	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	A	442	ARG	CD-NE-CZ	9.25	136.56	123.60
1	A	167	ARG	NE-CZ-NH1	8.96	124.78	120.30
1	A	219	TYR	CB-CG-CD2	-8.16	116.10	121.00
1	A	483	ARG	NE-CZ-NH1	-7.65	116.47	120.30
1	A	144	ARG	NE-CZ-NH1	7.65	124.12	120.30
1	A	123	ARG	NE-CZ-NH2	7.09	123.84	120.30
1	A	413	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	354	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	A	480	TYR	CB-CG-CD1	-6.09	117.34	121.00
1	A	219	TYR	CB-CG-CD1	5.58	124.35	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	ARG	CD-NE-CZ	5.53	131.35	123.60
1	A	442	ARG	NE-CZ-NH2	5.36	122.98	120.30
1	A	249	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	91	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	32	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	214	LYS	CG-CD-CE	-5.13	96.50	111.90
1	A	108	ASP	CB-CG-OD2	5.13	122.92	118.30
1	A	404	ASP	CB-CG-OD1	5.01	122.81	118.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	810	GLC	C1

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3948	0	3698	26	0
2	A	4	0	0	0	0
3	A	1	0	0	0	0
4	A	12	0	11	0	0
5	A	64	0	58	1	0
6	A	32	0	29	2	0
7	A	680	0	0	7	1
All	All	4741	0	3796	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189[A]:GLU:OE2	1:A:234:LYS:CE	2.25	0.85

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:GLN:HG3	7:A:2159:HOH:O	1.98	0.63
6:A:810:GLC:H5	7:A:2405:HOH:O	1.96	0.63
1:A:167:ARG:HD2	7:A:2290:HOH:O	2.02	0.59
1:A:6[B]:MET:HE2	1:A:30:LEU:HD22	1.85	0.59
1:A:189[A]:GLU:OE2	1:A:234:LYS:HE2	2.03	0.58
1:A:8:TYR:HB2	1:A:38[B]:VAL:HG13	1.85	0.58
1:A:189[A]:GLU:OE2	1:A:234:LYS:HE3	2.05	0.55
1:A:7:GLN:HE22	1:A:326:ASN:HB2	1.72	0.55
1:A:169:ILE:HD11	7:A:2290:HOH:O	2.06	0.54
1:A:416[A]:ASP:OD2	1:A:418:SER:OG	2.26	0.54
1:A:184:TRP:CD2	1:A:185:GLU:HA	2.46	0.50
1:A:189[A]:GLU:OE2	1:A:234:LYS:NZ	2.46	0.48
1:A:184:TRP:CG	1:A:185:GLU:HA	2.48	0.47
1:A:6[B]:MET:HE3	1:A:8:TYR:HD1	1.78	0.47
1:A:465:GLU:HG3	7:A:2641:HOH:O	2.14	0.47
1:A:189[A]:GLU:CD	1:A:234:LYS:HZ1	2.18	0.47
1:A:6[B]:MET:CE	1:A:30:LEU:HD22	2.45	0.46
1:A:385:GLU:N	1:A:386:PRO:CD	2.80	0.44
5:A:803:GLC:H5	5:A:804:ACI:H62	1.99	0.44
1:A:370[B]:LYS:HG3	7:A:2009:HOH:O	2.17	0.44
1:A:139:PHE:HB2	1:A:153:TRP:HB2	2.00	0.44
1:A:286:VAL:N	1:A:287:PRO:CD	2.82	0.42
1:A:114:THR:OG1	1:A:132:GLN:NE2	2.52	0.41
6:A:810:GLC:H3	7:A:2679:HOH:O	2.20	0.41
1:A:162:ASP:HA	1:A:171:ARG:O	2.21	0.41
1:A:458:GLU:HA	1:A:459:PRO:HD3	1.83	0.41
1:A:189[A]:GLU:CD	1:A:264:GLN:HE22	2.24	0.40
1:A:234:LYS:HG2	1:A:262:TYR:CE1	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:2419:HOH:O	7:A:2662:HOH:O[5_445]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	497/483 (103%)	480 (97%)	16 (3%)	1 (0%)	52 42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	337	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	420/404 (104%)	412 (98%)	8 (2%)	65 56

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLU
1	A	39	TRP
1	A	44	TYR
1	A	193	TYR
1	A	293	GLN
1	A	372	ASP
1	A	381	LYS
1	A	383	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	25	ASN
1	A	66	GLN
1	A	132	GLN
1	A	246	GLN
1	A	293	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

9 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GLD	A	802	5,4	9,9,10	1.40	2 (22%)	10,12,14	1.91	3 (30%)
5	GLC	A	803	5	11,11,12	1.23	2 (18%)	14,15,17	1.65	4 (28%)
5	ACI	A	804	5	11,12,12	0.90	1 (9%)	9,17,17	1.06	1 (11%)
5	GLD	A	805	5	9,9,10	1.66	4 (44%)	10,12,14	1.66	4 (40%)
5	GLC	A	806	5	11,11,12	1.26	1 (9%)	14,15,17	1.40	1 (7%)
5	ACI	A	807	5,6	11,12,12	1.51	2 (18%)	9,17,17	1.05	0
6	GLD	A	808	5,6	9,9,10	1.68	3 (33%)	10,12,14	1.61	1 (10%)
6	GLC	A	809	6	11,11,12	0.96	0	14,15,17	2.04	4 (28%)
6	GLC	A	810	6	12,12,12	0.74	0	17,17,17	1.53	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLD	A	802	5,4	-	0/0/13/16	0/1/1/1
5	GLC	A	803	5	-	0/2/19/22	0/1/1/1
5	ACI	A	804	5	-	0/2/22/22	0/1/1/1
5	GLD	A	805	5	-	0/0/13/16	0/1/1/1
5	GLC	A	806	5	-	0/2/19/22	0/1/1/1
5	ACI	A	807	5,6	-	0/2/22/22	0/1/1/1
6	GLD	A	808	5,6	-	0/0/13/16	0/1/1/1
6	GLC	A	809	6	-	0/2/19/22	0/1/1/1
6	GLC	A	810	6	1/1/5/5	0/2/22/22	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	808	GLD	O5-C1	-2.79	1.39	1.43
5	A	805	GLD	O5-C1	-2.47	1.39	1.43
5	A	803	GLC	O5-C1	-2.05	1.40	1.43
5	A	805	GLD	C4-C3	2.23	1.56	1.52
5	A	806	GLC	C2-C3	2.32	1.55	1.52
5	A	802	GLD	C4-C5	2.38	1.56	1.51
5	A	805	GLD	C4-C5	2.40	1.56	1.51
5	A	804	ACI	C3-C4	2.40	1.56	1.53
5	A	805	GLD	C3-C2	2.49	1.55	1.52
6	A	808	GLD	C4-C3	2.57	1.56	1.52
5	A	807	ACI	C7-C5	2.72	1.36	1.32
5	A	803	GLC	C2-C3	2.85	1.56	1.52
6	A	808	GLD	C4-C5	2.93	1.57	1.51
5	A	802	GLD	C4-C3	3.00	1.57	1.52
5	A	807	ACI	C3-C4	3.46	1.57	1.53

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	808	GLD	O5-C5-C4	-3.92	104.35	109.41
5	A	805	GLD	O2-C2-C3	-3.31	102.84	110.27
6	A	810	GLC	C3-C4-C5	-3.11	104.77	110.20
6	A	809	GLC	O5-C1-C2	-3.00	105.99	110.86
5	A	802	GLD	O3-C3-C4	-2.69	103.47	110.06
5	A	805	GLD	C4-C3-C2	-2.33	107.14	110.23
6	A	809	GLC	O2-C2-C3	-2.26	105.57	110.12

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	803	GLC	O5-C5-C6	-2.11	102.77	107.35
5	A	802	GLD	C4-C3-C2	-2.11	107.44	110.23
6	A	810	GLC	C4-C3-C2	-2.10	106.88	110.79
6	A	810	GLC	O5-C1-C2	-2.04	106.55	109.80
5	A	803	GLC	C6-C5-C4	-2.03	108.01	113.02
5	A	805	GLD	O2-C2-C1	2.00	113.22	109.21
5	A	805	GLD	C1-O5-C5	2.16	117.65	113.19
6	A	810	GLC	O5-C5-C4	2.46	114.30	109.68
5	A	803	GLC	O2-C2-C1	2.54	114.29	109.21
6	A	810	GLC	O3-C3-C2	2.71	116.44	110.34
5	A	804	ACI	O2-C2-C1	2.75	114.41	108.99
5	A	803	GLC	C1-O5-C5	3.29	116.42	112.25
6	A	809	GLC	O2-C2-C1	3.33	115.88	109.21
5	A	806	GLC	C1-O5-C5	3.74	117.00	112.25
5	A	802	GLD	O5-C5-C6	4.19	111.55	105.99
6	A	809	GLC	C1-O5-C5	5.27	118.93	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	810	GLC	C1

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	803	GLC	1	0
5	A	804	ACI	1	0
6	A	810	GLC	2	0

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is

the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	ACI	A	801	5	11,12,12	0.84	0	9,17,17	2.00	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACI	A	801	5	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	A	801	ACI	O3-C3-C4	-2.63	104.73	109.49
4	A	801	ACI	O6-C6-C5	-2.03	107.22	112.12
4	A	801	ACI	C2-C1-N1	2.05	115.56	111.40
4	A	801	ACI	O2-C2-C1	3.86	116.61	108.99

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	483/483 (100%)	-0.55	2 (0%) 93 95	5, 11, 22, 35	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	372	ASP	2.7
1	A	1	VAL	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	GLC	A	810	12/12	0.77	0.38	10.45	24,34,37,38	0
5	GLD	A	802	9/10	0.96	0.09	0.23	13,15,17,18	0
6	GLC	A	809	11/12	0.96	0.09	-0.18	10,15,19,22	0
5	ACI	A	807	12/12	0.98	0.07	-0.35	4,5,9,14	0
5	GLC	A	806	11/12	0.99	0.05	-0.66	5,7,9,10	0
6	GLD	A	808	9/10	0.98	0.05	-0.91	6,7,8,9	0
5	GLC	A	803	11/12	0.98	0.07	-0.92	13,14,19,20	0
5	ACI	A	804	12/12	0.97	0.07	-	10,12,18,24	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	GLD	A	805	9/10	0.97	0.08	-	9,11,12,12	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	ACI	A	801	12/12	0.90	0.12	0.88	20,28,30,32	0
3	NA	A	505	1/1	0.99	0.07	-0.48	5,5,5,5	0
2	CA	A	504	1/1	1.00	0.03	-2.61	12,12,12,12	0
2	CA	A	501	1/1	0.99	0.05	-3.65	5,5,5,5	0
2	CA	A	503	1/1	1.00	0.02	-3.68	12,12,12,12	0
2	CA	A	502	1/1	1.00	0.03	-4.66	6,6,6,6	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.