



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:20 AM GMT

PDB ID : 3E9Y
Title : Arabidopsis thaliana acetohydroxyacid synthase in complex with monosulfuron
Authors : Guddat, L.W.; Duggleby, R.G.; Wang, J.-G.; Li, Z.-M.
Deposited on : 2008-08-24
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TDM	A	700	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4646 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetolactate synthase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	582	4462	2831	767	839	25	0	4	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	330	THR	SER	CONFLICT	UNP P17597

- Molecule 2 is N-[(4-METHYLPYRIMIDIN-2-YL)CARBAMOYL]-2-NITROBENZENESULFONAMIDE (three-letter code: 1MS) (formula: C₁₂H₁₁N₅O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	23	12	5	5	1	0	0

- Molecule 3 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula: C₈H₁₇NO₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 2-[(2E)-3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-2-(1-H YDROXYETHYLIDENE)-4-METHYL-2,3-DIHYDRO-1,3-THIAZOL-5-YL]ETHYL TRIHYDROGEN DIPHOSPHATE (three-letter code: TDM) (formula: C₁₄H₂₂N₄O₈P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	S	0	0
			29	14	4	8	2	1		

- Molecule 6 is FLAVIN-ADENINE DINUCLEOTIDE-N5-ISOBUTYL KETONE (three-letter code: FAB) (formula: $C_{31}H_{39}N_9O_{16}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	65	Total	O	0	0
			65	65		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	178.87Å 178.87Å 186.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	89.44 – 3.00 89.43 – 3.00	Depositor EDS
% Data completeness (in resolution range)	80.4 (89.44-3.00) 80.4 (89.43-3.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.06 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.192 , 0.223 0.185 , 0.215	Depositor DCC
R_{free} test set	1448 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	61.8	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 28694 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4646	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, MG, TDM, CSD, 1MS, FAB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/4556	0.51	0/6190

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4462	0	4426	42	0
2	A	23	0	11	1	0
3	A	13	0	17	0	0
4	A	1	0	0	0	0
5	A	29	0	19	5	0
6	A	53	0	31	0	0
7	A	65	0	0	2	1
All	All	4646	0	4504	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:622:ARG:HG3	1:A:622:ARG:HH11	1.33	0.92
1:A:644:GLN:HG3	1:A:667:ARG:HG3	1.63	0.81
1:A:143:HIS:HD2	1:A:145:GLN:H	1.30	0.77
1:A:145:GLN:HE21	1:A:544:ASN:HD21	1.33	0.77
1:A:622:ARG:CG	1:A:622:ARG:HH11	1.98	0.74
1:A:566:GLN:HE22	1:A:598:PHE:H	1.33	0.74
1:A:543:MET:O	7:A:766:HOH:O	2.12	0.68
1:A:513:MET:SD	5:A:700:TDM:HM43	2.39	0.62
1:A:117:ALA:H	1:A:138:ASN:HD21	1.46	0.61
1:A:350:GLY:HA2	1:A:378:VAL:HA	1.83	0.60
1:A:656:THR:HG22	1:A:657:PHE:N	2.21	0.56
2:A:695:1MS:N13	2:A:695:1MS:O9	2.39	0.55
1:A:332:LEU:HA	1:A:348:MET:HE2	1.89	0.55
1:A:394:ILE:HG12	1:A:411:VAL:HB	1.91	0.53
1:A:614:ARG:HD2	1:A:640:ILE:HD11	1.91	0.52
1:A:570:MET:HB3	5:A:700:TDM:S1	2.50	0.52
5:A:700:TDM:C27	5:A:700:TDM:H4'1	2.23	0.51
1:A:376:ASP:OD2	1:A:377:ARG:NH1	2.39	0.51
1:A:144:GLU:HG3	1:A:174:ASN:HB2	1.93	0.50
1:A:375:ASP:OD2	1:A:377:ARG:HB2	2.12	0.50
1:A:656:THR:HG22	1:A:657:PHE:H	1.76	0.49
1:A:622:ARG:CG	1:A:622:ARG:NH1	2.66	0.48
1:A:351[A]:MET:HG3	1:A:651:ILE:O	2.13	0.48
1:A:144:GLU:HG2	1:A:144:GLU:H	1.37	0.46
1:A:622:ARG:HG3	1:A:622:ARG:NH1	2.13	0.46
1:A:562:LEU:HD21	1:A:601:MET:HG3	1.98	0.45
1:A:346:LEU:HD21	1:A:444:LEU:HD13	1.97	0.45
1:A:305:TYR:CE2	1:A:372:VAL:HG21	2.52	0.44
1:A:460:GLU:OE2	1:A:460:GLU:HA	2.18	0.44
1:A:668:ILE:HD13	1:A:668:ILE:H	1.82	0.44
1:A:114:THR:HG21	1:A:525:VAL:HG11	2.01	0.43
1:A:552:ARG:O	1:A:552:ARG:HD3	2.19	0.43
1:A:319:ARG:HA	1:A:322:GLU:HG2	2.01	0.43
1:A:466:TYR:CE2	1:A:618:LYS:HE3	2.53	0.43
1:A:533:VAL:HG22	1:A:559:LYS:HB2	2.01	0.42
1:A:485:VAL:HG21	1:A:511:GLY:C	2.39	0.42
5:A:700:TDM:O28	5:A:700:TDM:H7'1	2.19	0.42
1:A:95:GLN:HA	1:A:96:PRO:HD3	1.87	0.42
1:A:370:PHE:HB3	1:A:415:VAL:HG21	2.02	0.42
1:A:488:HIS:HB2	1:A:563:LEU:HD12	2.02	0.42
5:A:700:TDM:H4'1	5:A:700:TDM:C2	2.33	0.41
1:A:617:LYS:HB2	1:A:620:ASP:OD2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:552:ARG:HG2	1:A:608:CYS:O	2.21	0.41
1:A:352:HIS:CE1	1:A:650:MET:HG3	2.56	0.41
1:A:137:ARG:HD2	7:A:717:HOH:O	2.21	0.40
1:A:219:THR:HA	1:A:249:PRO:HG2	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:766:HOH:O	7:A:766:HOH:O[10_665]	1.87	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	583/584 (100%)	575 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/480 (99%)	455 (96%)	20 (4%)	36	76

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ARG
1	A	130	LEU
1	A	137	ARG
1	A	144	GLU
1	A	148	VAL
1	A	204	ASP
1	A	277	MET
1	A	285	GLU
1	A	351[A]	MET
1	A	351[B]	MET
1	A	375	ASP
1	A	377	ARG
1	A	378	VAL
1	A	381	LYS
1	A	441	ARG
1	A	460	GLU
1	A	580	LYS
1	A	614	ARG
1	A	622	ARG
1	A	668	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
1	A	143	HIS
1	A	145	GLN
1	A	174	ASN
1	A	261	GLN
1	A	269	GLN
1	A	288	HIS
1	A	312	ASN
1	A	566	GLN
1	A	643	HIS
1	A	644	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSD	A	340	1	3,7,8	1.18	0	3,8,10	1.82	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	340	1	-	0/2/6/8	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	340	CSD	OD1-SG-CB	2.42	109.43	105.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1MS	A	695	-	22,24,24	4.66	9 (40%)	30,34,34	3.27	14 (46%)
3	NHE	A	696	-	12,13,13	0.98	0	15,17,17	1.50	1 (6%)
5	TDM	A	700	4	24,30,30	2.83	3 (12%)	36,45,45	1.81	11 (30%)
6	FAB	A	701	-	48,58,63	2.31	9 (18%)	54,89,97	3.28	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1MS	A	695	-	-	0/18/19/19	0/2/2/2
3	NHE	A	696	-	-	0/7/15/15	0/1/1/1
5	TDM	A	700	4	-	0/16/21/21	0/2/2/2
6	FAB	A	701	-	-	0/30/50/58	0/6/6/6

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	695	1MS	C21-C20	-4.31	1.39	1.50
2	A	695	1MS	C18-N16	-3.89	1.34	1.39
2	A	695	1MS	C14-N13	-3.04	1.34	1.39
6	A	701	FAB	C10-N10	2.01	1.41	1.39
2	A	695	1MS	C20-N19	2.71	1.40	1.34
6	A	701	FAB	C10-N1	2.93	1.40	1.35
6	A	701	FAB	C5X-N5	3.20	1.40	1.35
5	A	700	TDM	PB-O1B	3.23	1.61	1.51
6	A	701	FAB	O4B-C1B	3.27	1.45	1.41
6	A	701	FAB	C4A-N3A	3.45	1.40	1.35
6	A	701	FAB	C4-N3	4.00	1.40	1.33
6	A	701	FAB	C4X-N5	4.52	1.40	1.33
2	A	695	1MS	O11-S10	5.57	1.49	1.43
2	A	695	1MS	O12-S10	6.00	1.50	1.43
2	A	695	1MS	C4-C3	7.44	1.54	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	700	TDM	O28-C27	7.57	1.53	1.31
6	A	701	FAB	C2A-N1A	7.83	1.48	1.33
6	A	701	FAB	C2A-N3A	9.37	1.48	1.32
5	A	700	TDM	C2-N3	9.84	1.54	1.38
2	A	695	1MS	O8-N7	10.27	1.43	1.22
2	A	695	1MS	C3-C2	13.80	1.54	1.39

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	701	FAB	N3A-C2A-N1A	-22.07	112.00	128.89
2	A	695	1MS	O12-S10-O11	-8.49	108.28	119.54
2	A	695	1MS	N17-C18-N19	-7.00	119.22	126.67
5	A	700	TDM	CM4-C4-C5	-3.25	121.58	128.90
5	A	700	TDM	C5'-C7'-N3	-3.14	108.10	113.31
2	A	695	1MS	C3-C2-S10	-3.08	113.64	117.39
5	A	700	TDM	N1'-C2'-N3'	-3.03	120.00	125.60
5	A	700	TDM	C29-C27-C2	-2.88	120.58	124.73
6	A	701	FAB	P-O3P-PA	-2.73	125.06	132.73
6	A	701	FAB	C4X-C4-N3	-2.68	119.93	123.59
2	A	695	1MS	C18-N16-C14	-2.62	126.87	130.19
2	A	695	1MS	C5-C4-C3	-2.60	116.38	120.19
2	A	695	1MS	C22-C23-N17	-2.55	120.99	123.90
5	A	700	TDM	PA-O3A-PB	-2.52	124.23	132.67
6	A	701	FAB	C4A-C5A-N7A	-2.41	107.27	109.48
2	A	695	1MS	C4-C3-C2	-2.01	115.01	118.74
5	A	700	TDM	O3A-PA-O7	2.03	108.33	102.94
6	A	701	FAB	C2A-N1A-C6A	2.12	122.56	118.77
2	A	695	1MS	N16-C18-N17	2.30	122.63	116.56
2	A	695	1MS	C21-C20-N19	2.42	120.34	116.59
5	A	700	TDM	C2-N3-C4	2.48	112.61	109.14
6	A	701	FAB	C5X-C9A-N10	2.55	119.56	117.62
5	A	700	TDM	C6-C5-C4	2.66	129.95	127.56
2	A	695	1MS	C3-C2-C1	2.71	122.62	119.94
5	A	700	TDM	CM2-C2'-N1'	2.76	120.34	117.03
2	A	695	1MS	C5-C6-C1	2.79	123.36	118.63
2	A	695	1MS	C4-C5-C6	3.14	124.78	120.19
5	A	700	TDM	C6'-N1'-C2'	3.27	121.49	115.77
5	A	700	TDM	C5-C4-N3	3.29	115.49	107.83
3	A	696	NHE	O2-S-C2	4.67	110.89	106.91
6	A	701	FAB	C4-N3-C2	5.17	119.72	115.25
2	A	695	1MS	C2-S10-N13	7.34	114.72	106.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	695	1MS	C23-N17-C18	7.81	122.28	115.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	695	1MS	1	0
5	A	700	TDM	5	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	581/584 (99%)	-0.35	0 100 100	34, 49, 71, 79	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CSD	A	340	8/9	0.94	0.15	-	63,64,65,65	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	TDM	A	700	29/29	0.96	0.25	2.03	53,65,66,66	0
2	1MS	A	695	23/23	0.97	0.21	1.32	65,66,67,67	0
6	FAB	A	701	53/58	0.98	0.17	-0.65	32,36,43,43	0
3	NHE	A	696	13/13	0.98	0.16	-0.75	48,48,49,49	0
4	MG	A	699	1/1	0.96	0.06	-3.64	48,48,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.