



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:21 AM GMT

PDB ID : 3EA4  
Title : Arabidopsis thaliana acetohydroxyacid synthase in complex with monosulfuron-ester  
Authors : Guddat, L.W.; Wang, J.-G.; Li, Z.-M.  
Deposited on : 2008-08-24  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	<div> <div></div> <div>88%</div> <div>11%</div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4707 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetolactate synthase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	0	3	0
			4479	2843	769	842	25			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	330	THR	SER	CONFLICT	UNP P17597

- Molecule 2 is METHYL 2-([(4-METHYLPYRIMIDIN-2-YL)CARBAMOYL]SULFAMOYL}BENZOATE (three-letter code: 2SM) (formula: C<sub>14</sub>H<sub>14</sub>N<sub>4</sub>O<sub>5</sub>S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			24	14	4	5	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE-N5-ISOBUTYL KETONE (three-letter code: FAB) (formula: C<sub>31</sub>H<sub>39</sub>N<sub>9</sub>O<sub>16</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula:  $\text{C}_8\text{H}_{17}\text{NO}_3\text{S}$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			13	8	1	3	1		

- Molecule 5 is 2-[(2E)-3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-2-(1-HYDROXYETHYLIDENE)-4-METHYL-2,3-DIHYDRO-1,3-THIAZOL-5-YL]ETHYL

TRIHYDROGEN DIPHOSPHATE (three-letter code: TDM) (formula:  $C_{14}H_{22}N_4O_8P_2S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	S	
			29	14	4	8	2	1	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg		
			1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	108	Total	O		
			108	108	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetolactate synthase, chloroplastic



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.47Å 178.47Å 184.95Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	58.42 – 2.80 58.42 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.5 (58.42-2.80) 98.5 (58.42-2.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.51 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.199 , 0.213 0.192 , 0.206	Depositor DCC
$R_{free}$ test set	2149 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.1	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 33.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Outliers	0 of 42764 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4707	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, 2SM, MG, TDM, CSD, FAB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.37	0/4570	0.52	0/6204

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4479	0	4471	36	0
2	A	24	0	14	1	0
3	A	53	0	31	0	0
4	A	13	0	16	1	0
5	A	29	0	19	4	0
6	A	1	0	0	0	0
7	A	108	0	0	2	0
All	All	4707	0	4551	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:644:GLN:HG3	1:A:667:ARG:HG3	1.57	0.84
1:A:145:GLN:HE21	1:A:544:ASN:HD21	1.30	0.80
1:A:117:ALA:H	1:A:138:ASN:HD21	1.30	0.80
1:A:566:GLN:HE22	1:A:598:PHE:H	1.33	0.76
1:A:332:LEU:HA	1:A:348:MET:HE2	1.66	0.76
1:A:143:HIS:HD2	1:A:145:GLN:H	1.33	0.72
1:A:137:ARG:HD2	7:A:702:HOH:O	1.97	0.63
1:A:513:MET:SD	5:A:698:TDM:HM43	2.39	0.62
1:A:394:ILE:HG12	1:A:411:VAL:HB	1.81	0.62
1:A:312:ASN:HD21	1:A:502:ARG:HH21	1.47	0.61
2:A:695:2SM:O25	2:A:695:2SM:N8	2.34	0.60
1:A:460:GLU:HB3	1:A:617:LYS:HE2	1.84	0.60
1:A:143:HIS:CD2	1:A:145:GLN:H	2.17	0.59
1:A:614:ARG:HD2	1:A:640:ILE:HD11	1.86	0.57
1:A:266:ASN:HB3	1:A:269:GLN:HE21	1.69	0.57
1:A:117:ALA:N	1:A:138:ASN:HD21	2.01	0.55
1:A:656:THR:HG22	1:A:657:PHE:H	1.74	0.53
1:A:656:THR:HG22	1:A:657:PHE:N	2.25	0.51
1:A:90:ARG:NH2	7:A:779:HOH:O	2.43	0.51
5:A:698:TDM:C27	5:A:698:TDM:H4'1	2.24	0.50
1:A:562:LEU:HD21	1:A:601:MET:HG3	1.93	0.50
1:A:370:PHE:HB3	1:A:415:VAL:HG21	1.94	0.49
1:A:92:ALA:HB3	1:A:95:GLN:HB2	1.93	0.49
1:A:350:GLY:HA2	1:A:378:VAL:HA	1.95	0.48
1:A:139:VAL:HG13	1:A:554:GLU:HG3	1.95	0.48
5:A:698:TDM:H7'1	5:A:698:TDM:O28	2.14	0.48
1:A:144:GLU:HG2	1:A:174:ASN:HB2	1.96	0.48
1:A:277:MET:HA	1:A:280:MET:HG3	1.97	0.47
1:A:570:MET:HB3	5:A:698:TDM:S1	2.55	0.46
1:A:144:GLU:HG2	1:A:174:ASN:CB	2.47	0.45
1:A:552:ARG:O	1:A:552:ARG:HD3	2.16	0.45
1:A:114:THR:HG21	1:A:525:VAL:HG11	1.99	0.44
4:A:697:NHE:HC11	4:A:697:NHE:H6'2	1.63	0.43
1:A:482:SER:HA	1:A:505:LEU:O	2.19	0.42
1:A:369:ALA:HB1	1:A:372:VAL:CG2	2.50	0.42
1:A:485:VAL:HG21	1:A:511:GLY:C	2.40	0.41
1:A:463:PRO:HA	1:A:464:PRO:HD2	1.92	0.41
1:A:198:ARG:HD3	1:A:258:ILE:HD11	2.03	0.41
1:A:401:ILE:HG21	1:A:410:SER:HB3	2.02	0.40
1:A:501:PRO:O	1:A:502:ARG:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	582/584 (100%)	575 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	480/480 (100%)	467 (97%)	13 (3%)	52	85

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	ARG
1	A	130	LEU
1	A	137	ARG
1	A	148	VAL
1	A	344	LEU
1	A	375	ASP
1	A	377	ARG
1	A	378	VAL
1	A	390	LYS
1	A	456	LYS
1	A	473	GLU
1	A	614	ARG
1	A	644	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	ASN
1	A	143	HIS
1	A	145	GLN
1	A	174	ASN
1	A	261	GLN
1	A	269	GLN
1	A	312	ASN
1	A	527	ASN
1	A	566	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSD	A	340	1	3,7,8	1.09	0	3,8,10	1.75	2 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	340	1	-	0/2/6/8	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	CSD	O-C-CA	-2.10	120.01	125.49
1	A	340	CSD	OD1-SG-CB	2.16	109.00	105.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	2SM	A	695	-	25,25,25	2.41	7 (28%)	35,35,35	3.17	11 (31%)
3	FAB	A	696	-	48,58,63	2.31	9 (18%)	54,89,97	3.30	11 (20%)
4	NHE	A	697	-	12,13,13	1.08	0	15,17,17	1.41	2 (13%)
5	TDM	A	698	6	24,30,30	3.00	5 (20%)	36,45,45	1.66	11 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2SM	A	695	-	-	0/21/21/21	0/2/2/2
3	FAB	A	696	-	-	0/30/50/58	0/6/6/6
4	NHE	A	697	-	-	0/7/15/15	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TDM	A	698	6	-	0/16/21/21	0/2/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	695	2SM	C11-N10	-3.98	1.34	1.39
2	A	695	2SM	C9-N8	-2.82	1.34	1.39
5	A	698	TDM	PB-O3B	2.05	1.62	1.54
5	A	698	TDM	C7'-C5'	2.27	1.56	1.51
3	A	696	FAB	C9A-N10	2.36	1.42	1.38
2	A	695	2SM	C13-N12	2.49	1.40	1.34
3	A	696	FAB	C4A-N3A	2.50	1.39	1.35
3	A	696	FAB	C10-N1	2.61	1.40	1.35
2	A	695	2SM	C15-N16	2.85	1.40	1.34
5	A	698	TDM	PB-O1B	3.40	1.62	1.51
3	A	696	FAB	O4B-C1B	3.47	1.45	1.41
3	A	696	FAB	C5X-N5	3.92	1.41	1.35
3	A	696	FAB	C4-N3	4.31	1.41	1.33
2	A	695	2SM	C11-N16	4.71	1.40	1.34
3	A	696	FAB	C4X-N5	5.14	1.41	1.33
2	A	695	2SM	O17-S7	5.89	1.50	1.43
2	A	695	2SM	C14-C13	5.91	1.54	1.39
3	A	696	FAB	C2A-N1A	7.47	1.48	1.33
5	A	698	TDM	O28-C27	8.70	1.56	1.31
3	A	696	FAB	C2A-N3A	9.10	1.48	1.32
5	A	698	TDM	C2-N3	9.97	1.54	1.38

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	696	FAB	N3A-C2A-N1A	-21.89	112.13	128.89
2	A	695	2SM	O17-S7-O18	-7.89	109.07	119.54
2	A	695	2SM	N16-C11-N12	-7.19	119.01	126.67
2	A	695	2SM	C15-C14-C13	-5.23	113.15	117.67
2	A	695	2SM	C24-O23-C22	-4.11	107.39	115.84
2	A	695	2SM	O23-C22-O25	-3.63	115.89	123.44
5	A	698	TDM	CM4-C4-C5	-3.23	121.65	128.90
3	A	696	FAB	C1B-N9A-C4A	-2.82	122.69	126.94
5	A	698	TDM	N1'-C2'-N3'	-2.77	120.47	125.60
5	A	698	TDM	C29-C27-C2	-2.69	120.85	124.73
3	A	696	FAB	C4X-C4-N3	-2.66	119.95	123.59
2	A	695	2SM	C11-N10-C9	-2.66	126.82	130.19

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	696	FAB	C4A-C5A-N7A	-2.64	107.06	109.48
3	A	696	FAB	P-O3P-PA	-2.56	125.54	132.73
5	A	698	TDM	PA-O3A-PB	-2.49	124.32	132.67
5	A	698	TDM	C5'-C7'-N3	-2.26	109.56	113.31
2	A	695	2SM	O20-C9-N10	-2.08	120.43	123.58
3	A	696	FAB	C4X-N5-C5X	2.05	119.13	116.76
3	A	696	FAB	C5X-C9A-N10	2.09	119.21	117.62
5	A	698	TDM	O28-C27-C29	2.12	119.44	114.82
5	A	698	TDM	C2-N3-C4	2.18	112.18	109.14
4	A	697	NHE	O1-S-C2	2.26	108.84	106.91
2	A	695	2SM	N10-C11-N16	2.27	122.56	116.56
3	A	696	FAB	C2A-N1A-C6A	2.38	123.02	118.77
3	A	696	FAB	C4-C4X-N5	2.41	121.65	118.72
3	A	696	FAB	C2B-C1B-N9A	2.57	118.21	114.29
5	A	698	TDM	C6-C5-C4	2.67	129.96	127.56
5	A	698	TDM	C6'-N1'-C2'	2.77	120.61	115.77
5	A	698	TDM	CM2-C2'-N1'	2.90	120.51	117.03
5	A	698	TDM	C5-C4-N3	3.39	115.71	107.83
4	A	697	NHE	O2-S-C2	4.32	110.59	106.91
3	A	696	FAB	C4-N3-C2	5.10	119.65	115.25
2	A	695	2SM	O23-C22-C4	5.56	121.69	112.28
2	A	695	2SM	C15-N16-C11	7.56	122.06	115.49
2	A	695	2SM	C5-S7-N8	7.58	115.00	106.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	695	2SM	1	0
4	A	697	NHE	1	0
5	A	698	TDM	4	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	581/584 (99%)	-0.23	7 (1%) 81 73	39, 50, 66, 87	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	668	ILE	4.3
1	A	431	GLU	2.6
1	A	430	ALA	2.3
1	A	93	PRO	2.3
1	A	424	LYS	2.3
1	A	88	ILE	2.2
1	A	94	ASP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSD	A	340	8/9	0.92	0.18	-	64,65,67,67	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	TDM	A	698	29/29	0.94	0.20	1.35	52,69,70,70	0
2	2SM	A	695	24/24	0.96	0.17	0.15	62,62,63,64	0
4	NHE	A	697	13/13	0.96	0.17	0.02	56,57,58,58	0
3	FAB	A	696	53/58	0.98	0.11	-1.57	34,39,43,43	0
6	MG	A	699	1/1	0.95	0.08	-3.65	55,55,55,55	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.