



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:21 AM GMT

PDB ID : 3EBM
Title : Crystal structure of human translationally controlled tumour associated protein (hTCTP) mutant E12V
Authors : Yang, B.; Dong, X.; Zhong, C.; Ding, J.
Deposited on : 2008-08-28
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	 2% 67% 13% • 18%
1	B	180	 % 73% 8% • 18%
1	C	180	 % 70% 8% • 19%
1	D	180	 2% 71% 8% • 20%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Translationally-controlled tumor protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	0	0	0
			1206	775	197	222	12			
1	B	147	Total	C	N	O	S	0	0	0
			1212	777	202	221	12			
1	C	145	Total	C	N	O	S	0	0	0
			1192	766	195	219	12			
1	D	144	Total	C	N	O	S	0	0	0
			1178	757	190	219	12			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	12	VAL	GLU	ENGINEERED	UNP P13693
A	173	LEU	-	EXPRESSION TAG	UNP P13693
A	174	GLU	-	EXPRESSION TAG	UNP P13693
A	175	HIS	-	EXPRESSION TAG	UNP P13693
A	176	HIS	-	EXPRESSION TAG	UNP P13693
A	177	HIS	-	EXPRESSION TAG	UNP P13693
A	178	HIS	-	EXPRESSION TAG	UNP P13693
A	179	HIS	-	EXPRESSION TAG	UNP P13693
A	180	HIS	-	EXPRESSION TAG	UNP P13693
B	12	VAL	GLU	ENGINEERED	UNP P13693
B	173	LEU	-	EXPRESSION TAG	UNP P13693
B	174	GLU	-	EXPRESSION TAG	UNP P13693
B	175	HIS	-	EXPRESSION TAG	UNP P13693
B	176	HIS	-	EXPRESSION TAG	UNP P13693
B	177	HIS	-	EXPRESSION TAG	UNP P13693
B	178	HIS	-	EXPRESSION TAG	UNP P13693
B	179	HIS	-	EXPRESSION TAG	UNP P13693
B	180	HIS	-	EXPRESSION TAG	UNP P13693
C	12	VAL	GLU	ENGINEERED	UNP P13693
C	173	LEU	-	EXPRESSION TAG	UNP P13693
C	174	GLU	-	EXPRESSION TAG	UNP P13693


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	175	HIS	-	EXPRESSION TAG	UNP P13693
C	176	HIS	-	EXPRESSION TAG	UNP P13693
C	177	HIS	-	EXPRESSION TAG	UNP P13693
C	178	HIS	-	EXPRESSION TAG	UNP P13693
C	179	HIS	-	EXPRESSION TAG	UNP P13693
C	180	HIS	-	EXPRESSION TAG	UNP P13693
D	12	VAL	GLU	ENGINEERED	UNP P13693
D	173	LEU	-	EXPRESSION TAG	UNP P13693
D	174	GLU	-	EXPRESSION TAG	UNP P13693
D	175	HIS	-	EXPRESSION TAG	UNP P13693
D	176	HIS	-	EXPRESSION TAG	UNP P13693
D	177	HIS	-	EXPRESSION TAG	UNP P13693
D	178	HIS	-	EXPRESSION TAG	UNP P13693
D	179	HIS	-	EXPRESSION TAG	UNP P13693
D	180	HIS	-	EXPRESSION TAG	UNP P13693

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total O 5 5	0	0
2	B	4	Total O 4 4	0	0
2	C	9	Total O 9 9	0	0
2	D	19	Total O 19 19	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.19Å 81.54Å 139.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.00 – 2.60 35.95 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.4 (36.00-2.60) 97.6 (35.95-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.237 , 0.280 0.235 , 0.246	Depositor DCC
R_{free} test set	1391 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	53.7	Xtriage
Anisotropy	0.869	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.2	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 27482 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4825	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1230	0.48	0/1646
1	B	0.42	0/1237	0.48	0/1654
1	C	0.36	0/1216	0.48	0/1627
1	D	0.59	3/1200 (0.2%)	0.47	0/1605
All	All	0.45	3/4883 (0.1%)	0.48	0/6532

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	130	LYS	CE-NZ	11.34	1.77	1.49
1	D	130	LYS	C-O	5.17	1.33	1.23
1	D	174	GLU	C-O	5.02	1.32	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1206	0	1201	14	0
1	B	1212	0	1202	10	0
1	C	1192	0	1184	11	0
1	D	1178	0	1176	10	0
2	A	5	0	0	0	0
2	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	9	0	0	0	0
2	D	19	0	0	0	0
All	All	4825	0	4763	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:130:LYS:CE	1:D:130:LYS:NZ	1.77	1.45
1:D:92:ILE:HG12	1:D:122:ILE:HD11	1.51	0.90
1:B:118:ALA:O	1:B:122:ILE:HG12	1.83	0.78
1:C:96:MET:HE1	1:C:122:ILE:HD11	1.69	0.73
1:D:92:ILE:O	1:D:96:MET:HG2	1.91	0.71
1:A:3:ILE:HG13	1:A:170:GLU:HB3	1.71	0.71
1:A:118:ALA:O	1:A:122:ILE:HG12	1.90	0.70
1:A:35:MET:HE3	1:A:157:THR:HA	1.73	0.69
1:A:35:MET:SD	1:A:35:MET:N	2.53	0.68
1:D:92:ILE:HG12	1:D:122:ILE:CD1	2.22	0.66
1:C:92:ILE:HG23	1:C:96:MET:HE3	1.86	0.58
1:C:153:GLU:H	1:C:153:GLU:CD	2.08	0.56
1:B:5:ARG:HD2	1:B:10:HIS:O	2.05	0.56
1:D:112:LYS:HB3	1:D:113:PRO:HD3	1.91	0.53
1:A:92:ILE:HG12	1:A:122:ILE:HD12	1.91	0.52
1:A:171:LYS:HE3	1:A:173:LEU:HD11	1.91	0.52
1:A:92:ILE:HG12	1:A:122:ILE:CD1	2.41	0.51
1:A:3:ILE:CG1	1:A:170:GLU:HB3	2.38	0.51
1:C:107:ARG:HG3	1:C:109:GLU:OE1	2.13	0.49
1:D:22:GLU:HG2	1:D:26:GLY:HA2	1.96	0.47
1:D:77:HIS:O	1:D:138:GLU:HB3	2.15	0.47
1:C:13:MET:HG3	1:C:162:PHE:HE2	1.81	0.46
1:C:145:MET:HE1	1:C:167:LEU:HG	2.00	0.44
1:B:37:SER:O	1:B:38:ARG:HB2	2.17	0.44
1:D:130:LYS:CD	1:D:130:LYS:NZ	2.72	0.44
1:A:151:TYR:HA	1:A:157:THR:O	2.18	0.43
1:B:128:ASN:HB3	1:B:131:ASN:HD22	1.82	0.43
1:C:22:GLU:HG2	1:C:26:GLY:HA2	2.01	0.43
1:C:112:LYS:N	1:C:113:PRO:HD2	2.34	0.43
1:C:92:ILE:HD13	1:C:92:ILE:HA	1.84	0.42
1:B:37:SER:O	1:B:38:ARG:CB	2.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:PRO:O	1:A:111:VAL:HG22	2.19	0.42
1:D:110:ARG:C	1:D:113:PRO:HD2	2.40	0.42
1:C:98:SER:O	1:C:102:LYS:HG2	2.19	0.42
1:B:2:ILE:HG21	1:B:169:MET:HE3	2.02	0.42
1:D:15:SER:OG	1:D:17:ILE:HG12	2.20	0.41
1:A:99:ILE:HD12	1:A:166:GLY:HA2	2.01	0.41
1:B:92:ILE:HA	1:B:92:ILE:HD13	1.86	0.41
1:A:30:GLU:OE1	1:A:152:ARG:NH1	2.52	0.41
1:B:32:GLU:HG3	1:B:152:ARG:HH21	1.85	0.41
1:B:92:ILE:HG13	1:B:122:ILE:HD12	2.03	0.40
1:C:13:MET:HG3	1:C:162:PHE:CE2	2.56	0.40
1:A:18:TYR:OH	1:A:75:ASN:ND2	2.47	0.40
1:B:22:GLU:HG2	1:B:29:LEU:HD23	2.04	0.40
1:A:102:LYS:O	1:A:106:GLN:HG2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/180 (79%)	138 (96%)	5 (4%)	0	100	100
1	B	143/180 (79%)	140 (98%)	3 (2%)	0	100	100
1	C	141/180 (78%)	138 (98%)	3 (2%)	0	100	100
1	D	140/180 (78%)	135 (96%)	5 (4%)	0	100	100
All	All	567/720 (79%)	551 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/157 (84%)	123 (93%)	9 (7%)	20	39
1	B	132/157 (84%)	129 (98%)	3 (2%)	58	83
1	C	130/157 (83%)	122 (94%)	8 (6%)	23	45
1	D	129/157 (82%)	127 (98%)	2 (2%)	70	89
All	All	523/628 (83%)	501 (96%)	22 (4%)	36	65

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	ILE
1	A	34	LYS
1	A	35	MET
1	A	75	ASN
1	A	90	LYS
1	A	100	LYS
1	A	128	ASN
1	A	148	LEU
1	A	171	LYS
1	B	38	ARG
1	B	92	ILE
1	B	96	MET
1	C	7	LEU
1	C	13	MET
1	C	75	ASN
1	C	92	ILE
1	C	102	LYS
1	C	107	ARG
1	C	153	GLU
1	C	154	ASP
1	D	25	ASP
1	D	148	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	121	GLN
1	B	131	ASN
1	C	75	ASN
1	C	131	ASN
1	C	139	ASN
1	D	131	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/180 (81%)	-0.03	3 (2%) 68 63	32, 51, 78, 119	0
1	B	147/180 (81%)	0.11	2 (1%) 78 74	33, 55, 76, 100	0
1	C	145/180 (80%)	-0.03	2 (1%) 78 74	34, 52, 71, 98	0
1	D	144/180 (80%)	0.24	4 (2%) 56 49	35, 57, 84, 119	0
All	All	583/720 (80%)	0.07	11 (1%) 70 64	32, 54, 78, 119	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	153	GLU	4.1
1	D	21	ARG	3.9
1	A	67	ILE	3.8
1	D	156	VAL	3.3
1	B	153	GLU	2.8
1	B	97	LYS	2.4
1	A	68	THR	2.4
1	C	153	GLU	2.3
1	D	19	LYS	2.3
1	C	34	LYS	2.2
1	D	174	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.