



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 23, 2016 – 05:31 PM EDT

PDB ID : 5EBE  
Title : Structure of human sphingomyelinase phosphodiesterase like 3A (SMPDL3A) with 5' CMP  
Authors : Lim, S.M.; Yeung, K.; Tresaugues, L.; Teo, H.L.; Nordlund, P.  
Deposited on : 2015-10-19  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	418	<div> <div>80%</div> <div>17%</div> <div>..</div> </div>
2	B	415	<div> <div>4%</div> <div>67%</div> <div>29%</div> <div>.</div> </div>
2	C	415	<div> <div>4%</div> <div>74%</div> <div>24%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	C5P	A	607	-	-	-	X
7	MLI	B	610	-	-	-	X

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 10472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acid sphingomyelinase-like phosphodiesterase 3a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	418	Total	C	N	O	S	0	4	0
			3380	2182	541	644	13			

- Molecule 2 is a protein called Acid sphingomyelinase-like phosphodiesterase 3a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	415	Total	C	N	O	S	0	3	0
			3344	2157	536	638	13			
2	C	415	Total	C	N	O	S	0	4	0
			3350	2159	537	641	13			

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			13	8	1	4		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Zn	0	0
			2	2		
4	A	2	Total	Zn	0	0
			2	2		
4	C	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CYTIDINE-5'-MONOPHOSPHATE (three-letter code: C5P) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>3</sub>O<sub>8</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			21	9	3	8	1		
5	C	1	Total	C	N	O	P	0	0
			21	9	3	8	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is MALONATE ION (three-letter code: MLI) (formula:  $C_3H_2O_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			7	3	4		

- Molecule 8 is 5-O-phosphono-beta-D-ribofuranose (three-letter code: RP5) (formula:  $C_5H_{11}O_8P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	1	Total	C	O	P	0	0
			14	5	8	1		

- Molecule 9 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	C	1	Total	C	N	S	0	0
			3	1	1	1		
9	C	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 10 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	19	Total	O	0	0
			19	19		
10	B	60	Total	O	0	0
			60	60		
10	C	51	Total	O	0	0
			51	51		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acid sphingomyelinase-like phosphodiesterase 3a




- Molecule 2: Acid sphingomyelinase-like phosphodiesterase 3a



- Molecule 2: Acid sphingomyelinase-like phosphodiesterase 3a





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	147.65Å 147.65Å 142.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.56 – 3.00 47.56 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.56-3.00) 99.7 (47.56-3.00)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.47 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.251 , 0.295 0.251 , 0.295	Depositor DCC
$R_{free}$ test set	1790 reflections (5.20%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.1	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 13.5	EDS
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	11 of 36201 reflections (0.030%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	10472	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 75.71 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2003e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, SCN, RP5, NAG, C5P, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.68	1/3479 (0.0%)	0.83	5/4749 (0.1%)
2	B	0.70	0/3443	0.84	2/4702 (0.0%)
2	C	0.67	0/3448	0.84	1/4708 (0.0%)
All	All	0.68	1/10370 (0.0%)	0.84	8/14159 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	52	TYR	CE1-CZ	-5.98	1.30	1.38

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	446	GLN	CB-CA-C	6.25	122.89	110.40
1	A	53	HIS	CB-CA-C	6.03	122.46	110.40
1	A	447	LEU	N-CA-CB	-5.81	98.79	110.40
2	B	443	CYS	CA-CB-SG	5.68	124.22	114.00
2	B	444	LEU	N-CA-C	-5.42	96.35	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3380	0	3300	48	0
2	B	3344	0	3262	87	0
2	C	3350	0	3263	75	0
3	A	55	0	50	0	0
3	B	70	0	64	4	0
3	C	56	0	52	1	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
5	A	21	0	12	4	0
5	C	21	0	12	0	0
6	B	12	0	16	0	0
7	B	7	0	2	0	0
8	B	14	0	9	1	0
9	C	6	0	0	0	0
10	A	19	0	0	0	0
10	B	60	0	0	13	0
10	C	51	0	0	13	0
All	All	10472	0	10042	212	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 212 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:430:CYS:SG	2:B:443:CYS:CB	2.17	1.31
2:B:430:CYS:CB	2:B:443:CYS:SG	2.29	1.19
2:C:294[A]:THR:CG2	10:C:702:HOH:O	1.97	1.09
1:A:448:TYR:O	1:A:449:ILE:HD13	1.61	0.99
2:C:294[A]:THR:HG22	10:C:702:HOH:O	1.60	0.93

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	420/418 (100%)	380 (90%)	38 (9%)	2 (0%)	34	76
2	B	416/415 (100%)	367 (88%)	47 (11%)	2 (0%)	34	76
2	C	417/415 (100%)	381 (91%)	35 (8%)	1 (0%)	52	88
All	All	1253/1248 (100%)	1128 (90%)	120 (10%)	5 (0%)	39	80

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	35	PRO
2	B	445	LYS
2	C	414	TYR
1	A	164	THR
1	A	216	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	384/380 (101%)	359 (94%)	25 (6%)	21	58
2	B	380/377 (101%)	350 (92%)	30 (8%)	15	48
2	C	381/377 (101%)	357 (94%)	24 (6%)	22	60
All	All	1145/1134 (101%)	1066 (93%)	79 (7%)	19	56

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	269	GLU
2	B	398	LEU
2	C	323	LYS
2	B	284	ASP
2	B	312	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	97	ASN
2	B	200	ASN
2	C	211	ASN
2	B	127	ASN
2	B	169	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

Of 27 ligands modelled in this entry, 6 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	601	1	14,14,15	2.11	4 (28%)	15,19,21	2.13	5 (33%)
3	NAG	A	602	1	14,14,15	1.99	3 (21%)	15,19,21	2.02	2 (13%)
3	NAG	A	603	1	14,14,15	1.21	2 (14%)	15,19,21	2.14	3 (20%)
3	NAG	A	604	-	13,13,15	3.89	4 (30%)	13,17,21	2.85	6 (46%)
5	C5P	A	607	4	19,22,22	0.67	0	24,33,33	1.18	4 (16%)
3	NAG	B	601	2	14,14,15	1.52	3 (21%)	15,19,21	2.27	4 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	B	602	3,2	14,14,15	1.77	2 (14%)	15,19,21	1.91	3 (20%)
3	NAG	B	603	3	14,14,15	1.69	4 (28%)	15,19,21	2.59	5 (33%)
3	NAG	B	604	2	14,14,15	1.33	1 (7%)	15,19,21	1.46	2 (13%)
3	NAG	B	605	2	14,14,15	3.10	5 (35%)	15,19,21	1.82	5 (33%)
6	GOL	B	608	-	5,5,5	0.57	0	5,5,5	0.65	0
6	GOL	B	609	-	5,5,5	0.76	0	5,5,5	0.56	0
7	MLI	B	610	-	0,6,6	0.00	-	0,7,7	0.00	-
8	RP5	B	611	4	14,14,14	1.15	1 (7%)	19,21,21	1.62	4 (21%)
3	NAG	C	601	2	14,14,15	3.08	4 (28%)	15,19,21	2.39	2 (13%)
3	NAG	C	602	2	14,14,15	1.90	3 (21%)	15,19,21	1.49	1 (6%)
3	NAG	C	603	2	14,14,15	1.90	2 (14%)	15,19,21	2.23	2 (13%)
3	NAG	C	604	2	14,14,15	2.02	4 (28%)	15,19,21	1.62	3 (20%)
5	C5P	C	607	4	19,22,22	1.16	2 (10%)	24,33,33	1.87	5 (20%)
9	SCN	C	608	-	2,2,2	1.71	1 (50%)	1,1,1	0.53	0
9	SCN	C	609	-	2,2,2	1.84	1 (50%)	1,1,1	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
3	NAG	A	602	1	-	0/6/23/26	0/1/1/1
3	NAG	A	603	1	-	0/6/23/26	0/1/1/1
3	NAG	A	604	-	-	0/6/19/26	0/1/1/1
5	C5P	A	607	4	-	0/6/26/26	0/2/2/2
3	NAG	B	601	2	-	0/6/23/26	0/1/1/1
3	NAG	B	602	3,2	-	0/6/23/26	0/1/1/1
3	NAG	B	603	3	-	0/6/23/26	0/1/1/1
3	NAG	B	604	2	-	0/6/23/26	0/1/1/1
3	NAG	B	605	2	-	0/6/23/26	0/1/1/1
6	GOL	B	608	-	-	0/4/4/4	0/0/0/0
6	GOL	B	609	-	-	0/4/4/4	0/0/0/0
7	MLI	B	610	-	-	0/0/4/4	0/0/0/0
8	RP5	B	611	4	-	0/6/22/22	0/1/1/1
3	NAG	C	601	2	-	0/6/23/26	0/1/1/1
3	NAG	C	602	2	-	0/6/23/26	0/1/1/1
3	NAG	C	603	2	-	0/6/23/26	0/1/1/1
3	NAG	C	604	2	-	0/6/23/26	0/1/1/1

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C5P	C	607	4	-	0/6/26/26	0/2/2/2
9	SCN	C	608	-	-	0/0/0/0	0/0/0/0
9	SCN	C	609	-	-	0/0/0/0	0/0/0/0

The worst 5 of 46 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	C	609	SCN	C-S	-2.60	1.48	1.63
9	C	608	SCN	C-S	-2.39	1.49	1.63
8	B	611	RP5	O1'-C1'	-2.18	1.40	1.42
3	B	601	NAG	O4-C4	2.05	1.47	1.43
3	A	603	NAG	C8-C7	2.09	1.54	1.50

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	NAG	C4-C3-C2	-4.88	103.77	111.34
8	B	611	RP5	O1-C1'-O1'	-4.58	104.95	111.21
3	A	604	NAG	O6-C6-C5	-3.58	99.36	111.30
3	A	604	NAG	O5-C1-C2	-3.31	105.96	110.56
5	C	607	C5P	C6-N1-C2	-3.17	116.15	121.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	607	C5P	4	0
3	B	602	NAG	2	0
3	B	603	NAG	2	0
8	B	611	RP5	1	0
3	C	604	NAG	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	418/418 (100%)	-0.58	0	100	100	10, 26, 51, 85	0
2	B	415/415 (100%)	0.01	16 (3%)	43	18	23, 44, 72, 119	0
2	C	415/415 (100%)	-0.08	15 (3%)	46	20	22, 43, 63, 87	0
All	All	1248/1248 (100%)	-0.22	31 (2%)	61	30	10, 39, 65, 119	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	303	SER	4.8
2	C	283	SER	4.0
2	B	235	SER	3.6
2	B	224	THR	3.4
2	B	134	THR	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
7	MLI	B	610	7/7	0.80	0.45	7.39	52,63,65,67	0
5	C5P	A	607	21/21	0.94	0.22	7.00	48,77,83,84	0
3	NAG	C	601	14/15	0.89	0.28	1.65	43,47,51,52	0
8	RP5	B	611	14/14	0.87	0.25	1.34	61,68,78,81	0
5	C5P	C	607	21/21	0.93	0.25	1.07	53,75,86,90	0
3	NAG	A	601	14/15	0.93	0.20	0.41	26,27,29,30	0
4	ZN	A	606	1/1	1.00	0.13	0.24	14,14,14,14	0
3	NAG	B	605	14/15	0.84	0.25	0.15	42,47,52,52	0
3	NAG	A	604	13/15	0.91	0.16	-0.38	31,34,39,42	0
3	NAG	C	602	14/15	0.90	0.19	-0.71	50,52,54,55	0
3	NAG	C	604	14/15	0.86	0.16	-0.94	47,50,54,55	0
3	NAG	B	601	14/15	0.92	0.17	-0.98	38,43,46,46	0
3	NAG	A	602	14/15	0.93	0.13	-1.22	35,37,40,40	0
4	ZN	B	606	1/1	0.99	0.09	-1.98	35,35,35,35	0
4	ZN	C	606	1/1	0.99	0.08	-2.86	32,32,32,32	0
4	ZN	B	607	1/1	0.97	0.05	-2.86	40,40,40,40	0
4	ZN	A	605	1/1	0.99	0.10	-3.03	27,27,27,27	0
4	ZN	C	605	1/1	0.97	0.04	-4.02	50,50,50,50	0
6	GOL	B	608	6/6	0.72	0.38	-	41,42,44,44	0
3	NAG	A	603	14/15	0.89	0.16	-	45,52,57,61	0
3	NAG	B	602	14/15	0.89	0.27	-	46,57,70,74	0
3	NAG	C	603	14/15	0.83	0.18	-	53,63,68,72	0
6	GOL	B	609	6/6	0.48	0.28	-	52,63,67,68	0
3	NAG	B	603	14/15	0.87	0.28	-	62,70,78,78	0
3	NAG	B	604	14/15	0.78	0.32	-	72,76,79,85	0
9	SCN	C	608	3/3	0.53	0.30	-	36,36,44,48	0
9	SCN	C	609	3/3	0.86	0.21	-	40,40,43,52	0

## 6.5 Other polymers

There are no such residues in this entry.