



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2017 – 12:02 PM EST

PDB ID : 5EF6  
Title : Structure of HOXB13 complex with methylated DNA  
Authors : Morgunova, E.; Yin, Y.; Jolma, A.; Popov, A.; Taipale, J.  
Deposited on : 2015-10-23  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.









Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	62	<div> <div>2%</div> <div>82% 13% . .</div> </div>
1	B	62	<div> <div>3%</div> <div>81% 13% 5% .</div> </div>
1	G	62	<div> <div>2%</div> <div>84% 13% .</div> </div>
1	J	62	<div> <div>2%</div> <div>84% 10% 6%</div> </div>
2	C	18	<div> <div></div> <div>78% 22%</div> </div>
2	D	18	<div> <div></div> <div>89% 11%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	18	 89%11%
2	K	18	 83%17%
3	E	18	 72%28%
3	F	18	 89%11%
3	I	18	 6%78%22%
3	L	18	 72%28%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Homeobox protein Hox-B13.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	61	Total	C	N	O	0	0	0
			521	329	105	87			
1	B	61	Total	C	N	O	0	0	0
			521	329	105	87			
1	G	62	Total	C	N	O	0	0	0
			528	334	106	88			
1	J	62	Total	C	N	O	0	0	0
			528	334	106	88			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*TP\*TP\*GP\*TP\*GP\*TP\*TP\*TP\*AP\*(5CM)P\*GP\*AP\*GP\*GP\*TP\*CP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	18	Total	C	N	O	P	0	0	0
			370	178	60	114	18			
2	D	18	Total	C	N	O	P	0	0	0
			370	178	60	114	18			
2	H	18	Total	C	N	O	P	0	0	0
			370	178	60	114	18			
2	K	18	Total	C	N	O	P	0	0	0
			370	178	60	114	18			

- Molecule 3 is a DNA chain called DNA (5'-D(P\*GP\*GP\*AP\*CP\*CP\*TP\*(5CM)P\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*CP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	18	Total	C	N	O	P	0	0	0
			370	176	74	102	18			
3	E	18	Total	C	N	O	P	0	0	0
			370	176	74	102	18			
3	I	18	Total	C	N	O	P	0	0	0
			370	176	74	102	18			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	18	Total	C	N	O	P	0	0	0
			370	176	74	102	18			



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	8	Total	O	0	0
			8	8		
4	C	14	Total	O	0	0
			14	14		
4	F	5	Total	O	0	0
			5	5		
4	B	11	Total	O	0	0
			11	11		
4	D	6	Total	O	0	0
			6	6		
4	E	11	Total	O	0	0
			11	11		
4	G	7	Total	O	0	0
			7	7		
4	H	7	Total	O	0	0
			7	7		
4	I	6	Total	O	0	0
			6	6		
4	J	8	Total	O	0	0
			8	8		
4	K	11	Total	O	0	0
			11	11		
4	L	7	Total	O	0	0
			7	7		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Homeobox protein Hox-B13





- Molecule 1: Homeobox protein Hox-B13





- Molecule 1: Homeobox protein Hox-B13




- Molecule 1: Homeobox protein Hox-B13



- Molecule 2: DNA (5'-D(P\*TP\*TP\*GP\*TP\*GP\*TP\*TP\*TP\*TP\*AP\*(5CM)P\*GP\*AP\*GP\*GP\*TP\*CP\*C)-3')



- Molecule 2: DNA (5'-D(P\*TP\*TP\*GP\*TP\*GP\*TP\*TP\*TP\*TP\*AP\*(5CM)P\*GP\*AP\*GP\*GP\*TP\*CP\*C)-3')

Chain D:  89% 11%





- Molecule 2: DNA (5'-D(P\*TP\*TP\*GP\*TP\*GP\*TP\*TP\*TP\*AP\*(5CM)P\*GP\*AP\*GP\*GP\*TP\*CP\*C)-3')

Chain H:  89% 11%




- Molecule 2: DNA (5'-D(P\*TP\*TP\*GP\*TP\*GP\*TP\*TP\*TP\*AP\*(5CM)P\*GP\*AP\*GP\*GP\*TP\*CP\*C)-3')

Chain K:  83% 17%




- Molecule 3: DNA (5'-D(P\*GP\*GP\*AP\*CP\*CP\*TP\*(5CM)P\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*CP\*AP\*A)-3')

Chain F:  89% 11%





- Molecule 3: DNA (5'-D(P\*GP\*GP\*AP\*CP\*CP\*TP\*(5CM)P\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*CP\*AP\*A)-3')

Chain E:  72% 28%




- Molecule 3: DNA (5'-D(P\*GP\*GP\*AP\*CP\*CP\*TP\*(5CM)P\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*CP\*AP\*A)-3')

Chain I:  6% 78% 22%



- Molecule 3: DNA (5'-D(P\*GP\*GP\*AP\*CP\*CP\*TP\*(5CM)P\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*CP\*AP\*A)-3')

Chain L:  72% 28%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.97Å 55.56Å 102.01Å 90.00° 102.24° 90.00°	Depositor
Resolution (Å)	38.17 – 3.00 55.16 – 2.98	Depositor EDS
% Data completeness (in resolution range)	94.9 (38.17-3.00) 97.8 (55.16-2.98)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.31 (at 2.96Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.291 , 0.320 0.294 , 0.315	Depositor DCC
$R_{free}$ test set	878 reflections (5.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.9	Xtriage
Anisotropy	0.917	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 41.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	5159	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 58.78 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9459e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/527	0.51	0/697
1	B	0.31	0/527	0.53	0/697
1	G	0.36	0/534	0.70	1/707 (0.1%)
1	J	0.32	0/534	0.60	1/707 (0.1%)
2	C	0.59	0/389	1.04	0/597
2	D	0.59	0/389	1.04	0/597
2	H	0.58	0/389	1.05	0/597
2	K	0.58	0/389	1.04	0/597
3	E	0.58	0/393	0.88	0/601
3	F	0.58	0/393	0.89	0/601
3	I	0.57	0/393	0.89	0/601
3	L	0.58	0/393	0.90	0/601
All	All	0.50	0/5250	0.85	2/7600 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	276	ALA	C-N-CA	5.41	135.24	121.70
1	J	276	ALA	C-N-CA	5.12	134.50	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	521	0	568	7	0
1	B	521	0	568	8	0
1	G	528	0	577	7	0
1	J	528	0	577	9	0
2	C	370	0	209	3	0
2	D	370	0	209	1	0
2	H	370	0	209	1	1
2	K	370	0	209	1	1
3	E	370	0	203	4	0
3	F	370	0	203	1	0
3	I	370	0	203	2	0
3	L	370	0	203	4	0
4	A	8	0	0	0	0
4	B	11	0	0	2	0
4	C	14	0	0	0	0
4	D	6	0	0	0	0
4	E	11	0	0	0	0
4	F	5	0	0	0	0
4	G	7	0	0	0	0
4	H	7	0	0	0	0
4	I	6	0	0	0	0
4	J	8	0	0	0	0
4	K	11	0	0	0	0
4	L	7	0	0	1	0
All	All	5159	0	3938	43	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:17:DA:N7	4:L:101:HOH:O	2.17	0.77
1:G:258:ARG:NH2	2:H:11:5CM:OP2	2.22	0.72
1:A:218:LYS:N	1:A:219:LYS:HA	2.10	0.66
1:B:266:ASN:ND2	4:B:303:HOH:O	2.29	0.64
1:B:218:LYS:N	1:B:219:LYS:HA	2.13	0.64

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:18:DC:O3'	2:H:18:DC:O3'[2_655]	2.08	0.12
2:K:18:DC:O3'	2:K:18:DC:O3'[2_755]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	59/62 (95%)	55 (93%)	3 (5%)	1 (2%)	11	46
1	B	59/62 (95%)	56 (95%)	2 (3%)	1 (2%)	11	46
1	G	60/62 (97%)	56 (93%)	2 (3%)	2 (3%)	5	26
1	J	60/62 (97%)	56 (93%)	1 (2%)	3 (5%)	3	15
All	All	238/248 (96%)	223 (94%)	8 (3%)	7 (3%)	6	29

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	277	LYS
1	J	277	LYS
1	B	220	ARG
1	J	220	ARG
1	A	220	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	55/56 (98%)	51 (93%)	4 (7%)	17	52
1	B	55/56 (98%)	52 (94%)	3 (6%)	27	65
1	G	56/56 (100%)	53 (95%)	3 (5%)	27	66
1	J	56/56 (100%)	53 (95%)	3 (5%)	27	66
All	All	222/224 (99%)	209 (94%)	13 (6%)	24	63

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	225	LYS
1	B	232	GLU
1	J	219	LYS
1	B	219	LYS
1	G	232	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	5CM	C	11	3,2	13,21,22	1.43	2 (15%)	17,30,33	1.23	2 (11%)
2	5CM	D	11	3,2	13,21,22	1.37	2 (15%)	17,30,33	1.18	1 (5%)
3	5CM	E	7	3,2	13,21,22	1.35	1 (7%)	17,30,33	1.50	3 (17%)
3	5CM	F	7	3,2	13,21,22	1.36	1 (7%)	17,30,33	1.54	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	5CM	H	11	3,2	13,21,22	1.44	2 (15%)	17,30,33	1.28	2 (11%)
3	5CM	I	7	3,2	13,21,22	1.37	1 (7%)	17,30,33	1.53	2 (11%)
2	5CM	K	11	3,2	13,21,22	1.36	2 (15%)	17,30,33	1.24	2 (11%)
3	5CM	L	7	3,2	13,21,22	1.36	1 (7%)	17,30,33	1.51	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	C	11	3,2	-	0/3/21/22	0/2/2/2
2	5CM	D	11	3,2	-	0/3/21/22	0/2/2/2
3	5CM	E	7	3,2	-	0/3/21/22	0/2/2/2
3	5CM	F	7	3,2	-	0/3/21/22	0/2/2/2
2	5CM	H	11	3,2	-	0/3/21/22	0/2/2/2
3	5CM	I	7	3,2	-	0/3/21/22	0/2/2/2
2	5CM	K	11	3,2	-	0/3/21/22	0/2/2/2
3	5CM	L	7	3,2	-	0/3/21/22	0/2/2/2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	11	5CM	O3'-C3'	-2.44	1.37	1.43
2	C	11	5CM	O3'-C3'	-2.32	1.38	1.43
2	D	11	5CM	O3'-C3'	-2.07	1.38	1.43
2	K	11	5CM	O3'-C3'	-2.04	1.38	1.43
2	K	11	5CM	C5-C4	2.59	1.45	1.41

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	7	5CM	C2'-C1'-N1	-2.80	107.24	114.14
3	F	7	5CM	C2'-C1'-N1	-2.72	107.42	114.14
3	L	7	5CM	C2'-C1'-N1	-2.65	107.61	114.14
3	E	7	5CM	C2'-C1'-N1	-2.44	108.11	114.14
2	H	11	5CM	N4-C4-N3	-2.17	113.73	116.92

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	11	5CM	1	0
3	E	7	5CM	1	0
3	F	7	5CM	1	0
2	H	11	5CM	1	0
3	I	7	5CM	1	0
3	L	7	5CM	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	61/62 (98%)	-0.01	1 (1%) 74 47	15, 31, 59, 78	0
1	B	61/62 (98%)	-0.02	2 (3%) 50 22	20, 34, 64, 97	0
1	G	62/62 (100%)	0.01	1 (1%) 74 47	16, 31, 75, 102	0
1	J	62/62 (100%)	-0.09	1 (1%) 74 47	17, 33, 77, 90	0
2	C	17/18 (94%)	-0.26	0 100 100	21, 36, 69, 77	0
2	D	17/18 (94%)	-0.20	0 100 100	23, 40, 82, 110	0
2	H	17/18 (94%)	-0.16	0 100 100	25, 42, 74, 88	0
2	K	17/18 (94%)	-0.07	0 100 100	20, 39, 71, 84	0
3	E	17/18 (94%)	0.12	0 100 100	26, 40, 73, 74	0
3	F	17/18 (94%)	-0.10	0 100 100	25, 34, 64, 69	0
3	I	17/18 (94%)	-0.01	1 (5%) 26 10	27, 42, 69, 77	0
3	L	17/18 (94%)	0.12	0 100 100	25, 36, 66, 68	0
All	All	382/392 (97%)	-0.04	6 (1%) 74 47	15, 36, 77, 110	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	218	LYS	4.6
1	J	218	LYS	4.0
1	G	278	VAL	3.6
1	B	217	ARG	3.4
1	A	217	ARG	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron



density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	5CM	C	11	20/21	0.90	0.18	-	15,29,37,48	0
3	5CM	F	7	20/21	0.92	0.17	-	20,34,47,49	0
2	5CM	D	11	20/21	0.92	0.14	-	23,31,42,46	0
3	5CM	E	7	20/21	0.94	0.16	-	20,30,47,49	0
2	5CM	K	11	20/21	0.93	0.15	-	16,38,52,53	0
2	5CM	H	11	20/21	0.93	0.18	-	22,36,48,49	0
3	5CM	L	7	20/21	0.93	0.16	-	27,31,50,58	0
3	5CM	I	7	20/21	0.93	0.19	-	21,39,48,54	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.