



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2016 – 01:11 PM EDT

PDB ID : 5EGN
Title : Est816 as an N-Acyl homoserine lactone degrading enzyme
Authors : Xie, W.; Liu, X.; Cao, L.; Liu, Y.
Deposited on : 2015-10-27
Resolution : 2.64 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20027939
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3377 (2.68-2.60)
Clashscore	102246	3781 (2.68-2.60)
Ramachandran outliers	100387	3722 (2.68-2.60)
Sidechain outliers	100360	3722 (2.68-2.60)
RSRZ outliers	91569	3388 (2.68-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	271	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>..</div> </div> </div>
1	C	271	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>.</div> </div> </div>
1	D	271	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div>.</div> </div> </div>
1	E	271	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>.</div> </div> </div>
1	F	271	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	271	<div><div></div><div>11%</div><div>83%</div><div>13%</div><div></div></div>
1	H	271	<div><div></div><div>6%</div><div>81%</div><div>14%</div><div>5%</div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 16184 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			1983	1264	344	370	5			
1	B	260	Total	C	N	O	S	0	0	0
			1991	1268	345	373	5			
1	C	259	Total	C	N	O	S	0	0	0
			1986	1265	344	372	5			
1	D	259	Total	C	N	O	S	0	0	0
			1980	1259	344	372	5			
1	E	259	Total	C	N	O	S	0	0	0
			1980	1259	344	372	5			
1	F	260	Total	C	N	O	S	0	0	0
			1979	1258	345	371	5			
1	G	259	Total	C	N	O	S	0	0	0
			1950	1242	334	369	5			
1	H	258	Total	C	N	O	S	0	0	0
			1935	1233	335	362	5			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	216	VAL	ALA	engineered mutation	UNP I6YRG4
A	238	ASN	LYS	engineered mutation	UNP I6YRG4
A	262	ALA	-	expression tag	UNP I6YRG4
A	263	ALA	-	expression tag	UNP I6YRG4
A	264	LEU	-	expression tag	UNP I6YRG4
A	265	GLU	-	expression tag	UNP I6YRG4
A	266	HIS	-	expression tag	UNP I6YRG4
A	267	HIS	-	expression tag	UNP I6YRG4
A	268	HIS	-	expression tag	UNP I6YRG4
A	269	HIS	-	expression tag	UNP I6YRG4
A	270	HIS	-	expression tag	UNP I6YRG4
A	271	HIS	-	expression tag	UNP I6YRG4
B	216	VAL	ALA	engineered mutation	UNP I6YRG4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	238	ASN	LYS	engineered mutation	UNP I6YRG4
B	262	ALA	-	expression tag	UNP I6YRG4
B	263	ALA	-	expression tag	UNP I6YRG4
B	264	LEU	-	expression tag	UNP I6YRG4
B	265	GLU	-	expression tag	UNP I6YRG4
B	266	HIS	-	expression tag	UNP I6YRG4
B	267	HIS	-	expression tag	UNP I6YRG4
B	268	HIS	-	expression tag	UNP I6YRG4
B	269	HIS	-	expression tag	UNP I6YRG4
B	270	HIS	-	expression tag	UNP I6YRG4
B	271	HIS	-	expression tag	UNP I6YRG4
C	216	VAL	ALA	engineered mutation	UNP I6YRG4
C	238	ASN	LYS	engineered mutation	UNP I6YRG4
C	262	ALA	-	expression tag	UNP I6YRG4
C	263	ALA	-	expression tag	UNP I6YRG4
C	264	LEU	-	expression tag	UNP I6YRG4
C	265	GLU	-	expression tag	UNP I6YRG4
C	266	HIS	-	expression tag	UNP I6YRG4
C	267	HIS	-	expression tag	UNP I6YRG4
C	268	HIS	-	expression tag	UNP I6YRG4
C	269	HIS	-	expression tag	UNP I6YRG4
C	270	HIS	-	expression tag	UNP I6YRG4
C	271	HIS	-	expression tag	UNP I6YRG4
D	216	VAL	ALA	engineered mutation	UNP I6YRG4
D	238	ASN	LYS	engineered mutation	UNP I6YRG4
D	262	ALA	-	expression tag	UNP I6YRG4
D	263	ALA	-	expression tag	UNP I6YRG4
D	264	LEU	-	expression tag	UNP I6YRG4
D	265	GLU	-	expression tag	UNP I6YRG4
D	266	HIS	-	expression tag	UNP I6YRG4
D	267	HIS	-	expression tag	UNP I6YRG4
D	268	HIS	-	expression tag	UNP I6YRG4
D	269	HIS	-	expression tag	UNP I6YRG4
D	270	HIS	-	expression tag	UNP I6YRG4
D	271	HIS	-	expression tag	UNP I6YRG4
E	216	VAL	ALA	engineered mutation	UNP I6YRG4
E	238	ASN	LYS	engineered mutation	UNP I6YRG4
E	262	ALA	-	expression tag	UNP I6YRG4
E	263	ALA	-	expression tag	UNP I6YRG4
E	264	LEU	-	expression tag	UNP I6YRG4
E	265	GLU	-	expression tag	UNP I6YRG4
E	266	HIS	-	expression tag	UNP I6YRG4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	267	HIS	-	expression tag	UNP I6YRG4
E	268	HIS	-	expression tag	UNP I6YRG4
E	269	HIS	-	expression tag	UNP I6YRG4
E	270	HIS	-	expression tag	UNP I6YRG4
E	271	HIS	-	expression tag	UNP I6YRG4
F	216	VAL	ALA	engineered mutation	UNP I6YRG4
F	238	ASN	LYS	engineered mutation	UNP I6YRG4
F	262	ALA	-	expression tag	UNP I6YRG4
F	263	ALA	-	expression tag	UNP I6YRG4
F	264	LEU	-	expression tag	UNP I6YRG4
F	265	GLU	-	expression tag	UNP I6YRG4
F	266	HIS	-	expression tag	UNP I6YRG4
F	267	HIS	-	expression tag	UNP I6YRG4
F	268	HIS	-	expression tag	UNP I6YRG4
F	269	HIS	-	expression tag	UNP I6YRG4
F	270	HIS	-	expression tag	UNP I6YRG4
F	271	HIS	-	expression tag	UNP I6YRG4
G	216	VAL	ALA	engineered mutation	UNP I6YRG4
G	238	ASN	LYS	engineered mutation	UNP I6YRG4
G	262	ALA	-	expression tag	UNP I6YRG4
G	263	ALA	-	expression tag	UNP I6YRG4
G	264	LEU	-	expression tag	UNP I6YRG4
G	265	GLU	-	expression tag	UNP I6YRG4
G	266	HIS	-	expression tag	UNP I6YRG4
G	267	HIS	-	expression tag	UNP I6YRG4
G	268	HIS	-	expression tag	UNP I6YRG4
G	269	HIS	-	expression tag	UNP I6YRG4
G	270	HIS	-	expression tag	UNP I6YRG4
G	271	HIS	-	expression tag	UNP I6YRG4
H	216	VAL	ALA	engineered mutation	UNP I6YRG4
H	238	ASN	LYS	engineered mutation	UNP I6YRG4
H	262	ALA	-	expression tag	UNP I6YRG4
H	263	ALA	-	expression tag	UNP I6YRG4
H	264	LEU	-	expression tag	UNP I6YRG4
H	265	GLU	-	expression tag	UNP I6YRG4
H	266	HIS	-	expression tag	UNP I6YRG4
H	267	HIS	-	expression tag	UNP I6YRG4
H	268	HIS	-	expression tag	UNP I6YRG4
H	269	HIS	-	expression tag	UNP I6YRG4
H	270	HIS	-	expression tag	UNP I6YRG4
H	271	HIS	-	expression tag	UNP I6YRG4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	41	Total 41	O 41	0	0
2	B	67	Total 67	O 67	0	0
2	C	59	Total 59	O 59	0	0
2	D	49	Total 49	O 49	0	0
2	E	58	Total 58	O 58	0	0
2	F	61	Total 61	O 61	0	0
2	G	28	Total 28	O 28	0	0
2	H	37	Total 37	O 37	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Esterase




• Molecule 1: Esterase




• Molecule 1: Esterase





• Molecule 1: Esterase










- Molecule 1: Esterase






- Molecule 1: Esterase



- Molecule 1: Esterase



- Molecule 1: Esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	109.81Å 78.55Å 116.47Å 90.00° 99.37° 90.00°	Depositor
Resolution (Å)	44.82 – 2.64 44.82 – 2.64	Depositor EDS
% Data completeness (in resolution range)	98.6 (44.82-2.64) 91.2 (44.82-2.64)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.65Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.219 , 0.264 0.220 , 0.261	Depositor DCC
R_{free} test set	2725 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.471	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16184	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2033	0.40	0/2767
1	B	0.23	0/2041	0.40	0/2777
1	C	0.23	0/2036	0.41	0/2770
1	D	0.23	0/2029	0.40	0/2761
1	E	0.23	0/2029	0.41	0/2761
1	F	0.24	0/2027	0.42	0/2758
1	G	0.24	0/1997	0.42	0/2720
1	H	0.25	0/1983	0.43	0/2703
All	All	0.24	0/16175	0.41	0/22017

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1983	0	1921	16	0
1	B	1991	0	1931	22	0
1	C	1986	0	1926	18	0
1	D	1980	0	1919	17	1
1	E	1980	0	1919	26	0
1	F	1979	0	1914	14	0
1	G	1950	0	1860	21	0
1	H	1935	0	1849	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	41	0	0	1	0
2	B	67	0	0	2	0
2	C	59	0	0	0	0
2	D	49	0	0	1	0
2	E	58	0	0	0	0
2	F	61	0	0	0	0
2	G	28	0	0	3	0
2	H	37	0	0	0	0
All	All	16184	0	15239	146	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:55:GLY:O	2:G:301:HOH:O	1.91	0.86
1:E:80:ASP:OD1	1:E:109:ARG:NH2	2.14	0.81
1:G:61:LYS:NZ	1:G:173:SER:O	2.15	0.80
1:H:135:GLN:OE1	1:H:135:GLN:N	2.17	0.76
1:G:174:ASN:O	2:G:301:HOH:O	2.04	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:64:THR:OG1	1:D:238:ASN:ND2[2_547]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/271 (95%)	248 (96%)	10 (4%)	0	100	100
1	B	258/271 (95%)	250 (97%)	7 (3%)	1 (0%)	39	63
1	C	257/271 (95%)	245 (95%)	11 (4%)	1 (0%)	39	63
1	D	257/271 (95%)	249 (97%)	8 (3%)	0	100	100
1	E	257/271 (95%)	250 (97%)	7 (3%)	0	100	100
1	F	258/271 (95%)	251 (97%)	7 (3%)	0	100	100
1	G	257/271 (95%)	250 (97%)	7 (3%)	0	100	100
1	H	256/271 (94%)	243 (95%)	12 (5%)	1 (0%)	39	63
All	All	2058/2168 (95%)	1986 (96%)	69 (3%)	3 (0%)	56	79

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	197	ASP
1	B	260	VAL
1	C	135	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/218 (95%)	204 (99%)	3 (1%)	74	90
1	B	209/218 (96%)	206 (99%)	3 (1%)	74	90
1	C	209/218 (96%)	207 (99%)	2 (1%)	82	93
1	D	208/218 (95%)	205 (99%)	3 (1%)	74	90
1	E	208/218 (95%)	207 (100%)	1 (0%)	92	98
1	F	206/218 (94%)	203 (98%)	3 (2%)	72	89
1	G	200/218 (92%)	198 (99%)	2 (1%)	82	93
1	H	198/218 (91%)	195 (98%)	3 (2%)	72	89
All	All	1645/1744 (94%)	1625 (99%)	20 (1%)	78	92

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	86	ARG
1	D	129	GLU
1	G	197	ASP
1	C	238	ASN
1	D	69	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	222	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/271 (95%)	0.11	7 (2%) 58 52	20, 30, 48, 63	0
1	B	260/271 (95%)	-0.15	2 (0%) 87 85	18, 25, 39, 67	0
1	C	259/271 (95%)	-0.12	5 (1%) 70 65	19, 25, 44, 66	0
1	D	259/271 (95%)	-0.05	5 (1%) 70 65	18, 26, 46, 69	0
1	E	259/271 (95%)	-0.06	8 (3%) 52 46	18, 24, 45, 65	0
1	F	260/271 (95%)	-0.10	5 (1%) 70 65	20, 26, 43, 63	0
1	G	259/271 (95%)	0.73	29 (11%) 7 4	27, 44, 67, 73	0
1	H	258/271 (95%)	0.45	15 (5%) 26 20	25, 39, 55, 66	0
All	All	2074/2168 (95%)	0.10	76 (3%) 45 38	18, 28, 55, 73	0

The worst 5 of 76 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	130	ALA	5.2
1	G	130	ALA	5.2
1	G	66	TYR	4.9
1	G	185	ALA	4.8
1	D	130	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.