



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 11:57 PM GMT

PDB ID : 5EHZ  
Title : mAChE-syn TZ2PA5 complex from an equimolar mixture of the syn/anti isomers  
Authors : Bourne, Y.; Marchot, P.  
Deposited on : 2015-10-29  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	<div> <div>11%</div> <div>88%</div> <div>10%</div> <div>..</div> </div>
1	B	543	<div> <div>25%</div> <div>90%</div> <div>8%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	601	-	-	-	X
3	5NZ	B	603	-	-	-	X

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8903 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	538	Total	C	N	O	S	0	3	0
			4216	2705	732	765	14			
1	B	534	Total	C	N	O	S	0	3	0
			4182	2685	720	763	14			

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is 6-phenyl-5-[5-[3-[2-(1,2,3,4-tetrahydroacridin-9-ylamino)ethyl]-1,2,3-triazol-4-yl]pentyl]phenanthridin-5-ium-3,8-diamine (three-letter code: 5NZ) (formula: C<sub>41</sub>H<sub>43</sub>N<sub>8</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 49	C 41	N 8	0	0
3	B	1	Total 49	C 41	N 8	0	0

- Molecule 4 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula:  $\text{C}_{12}\text{H}_{26}\text{O}_7$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			19	12	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	203	Total 203	O 203	0	0
5	B	143	Total 143	O 143	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Acetylcholinesterase



#### • Molecule 1: Acetylcholinesterase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.11Å 110.97Å 227.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.00 – 2.50 46.12 – 2.46	Depositor EDS
% Data completeness (in resolution range)	99.6 (44.00-2.50) 96.3 (46.12-2.46)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.47 (at 2.45Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.176 , 0.197 0.312 , 0.312	Depositor DCC
$R_{free}$ test set	1400 reflections (2.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.4	Xtriage
Anisotropy	0.888	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 34.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	3 of 71018 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	8903	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5NZ, NAG, P6G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/4350	0.70	0/5944
1	B	0.50	0/4317	0.69	0/5900
All	All	0.51	0/8667	0.69	0/11844

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	409	CYS	Mainchain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4216	0	4106	30	0
1	B	4182	0	4065	27	0
2	A	28	0	26	0	0
2	B	14	0	13	0	0
3	A	49	0	43	4	0
3	B	49	0	43	2	0
4	B	19	0	26	1	0
5	A	203	0	0	2	0
5	B	143	0	0	0	0
All	All	8903	0	8322	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:VAL:H	1:B:223:HIS:HD2	1.27	0.83
1:A:197:VAL:H	1:A:223:HIS:HD2	1.28	0.79
1:B:497:SER:HB2	1:B:498:PRO:HA	1.65	0.78
1:B:497:SER:CB	1:B:498:PRO:HA	2.22	0.69
1:B:197:VAL:H	1:B:223:HIS:CD2	2.10	0.68
1:A:197:VAL:H	1:A:223:HIS:CD2	2.10	0.68
1:A:369[B]:GLN:HE22	1:A:405:HIS:CE1	2.15	0.65
1:B:113:PRO:HG2	1:B:485:ARG:HG2	1.82	0.62
1:A:113:PRO:HG2	1:A:485:ARG:HG2	1.84	0.58
1:B:497:SER:HB2	1:B:498:PRO:CA	2.32	0.58
1:B:68:VAL:HG23	1:B:90:ARG:HB2	1.87	0.57
1:A:68:VAL:HG11	1:A:88:PRO:HB3	1.87	0.57
1:B:224:ARG:HG2	1:B:325:GLN:HB2	1.87	0.56
1:B:497:SER:CB	1:B:498:PRO:CA	2.84	0.55
1:A:498:PRO:HG2	1:A:518:LEU:HB2	1.90	0.54
1:A:224:ARG:HG2	1:A:325:GLN:HB2	1.89	0.53
1:B:381:HIS:HA	4:B:602:P6G:H51	1.89	0.53
3:A:603:5NZ:H29	5:A:721:HOH:O	2.08	0.53
1:B:68:VAL:HG11	1:B:88:PRO:HB3	1.89	0.53
1:A:337:TYR:O	1:A:340:VAL:HG22	2.09	0.53
1:B:454:ILE:HD13	1:B:476:LEU:HB3	1.90	0.52
1:B:84:GLU:HG3	1:B:87:ASN:HD22	1.74	0.52
3:A:603:5NZ:C29	5:A:721:HOH:O	2.58	0.52
1:A:491:ASP:OD1	1:A:493:ARG:HG3	2.11	0.51
1:B:498:PRO:HG2	1:B:518:LEU:O	2.11	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:ASP:CG	3:A:603:5NZ:H8	2.32	0.49
1:B:328:VAL:O	1:B:427:ALA:HA	2.14	0.48
1:B:439:TRP:CZ2	3:B:603:5NZ:H35	2.49	0.47
1:A:211:MET:HG2	1:A:308:LEU:HD21	1.97	0.47
1:A:340:VAL:HG11	1:A:443:MET:CE	2.45	0.46
1:A:68:VAL:HG23	1:A:90:ARG:HB2	1.97	0.46
1:B:211:MET:HG2	1:B:308:LEU:HD21	1.96	0.46
1:A:5:ASP:HB3	1:A:8:LEU:HD12	1.97	0.46
1:A:340:VAL:HG11	1:A:443:MET:HE1	1.98	0.45
1:A:370:ALA:HA	1:A:540:LEU:HD11	1.99	0.45
1:B:353:LEU:HB3	1:B:391:PRO:HB2	1.98	0.45
1:B:374:ALA:HA	1:B:539:LEU:HD23	1.99	0.45
1:A:328:VAL:O	1:A:427:ALA:HA	2.16	0.44
1:A:353:LEU:HB3	1:A:391:PRO:HB2	1.98	0.44
1:A:202:GLU:HA	1:A:228:GLN:O	2.17	0.44
1:A:341:TYR:CE1	3:A:603:5NZ:H10	2.53	0.43
1:B:202:GLU:HA	1:B:228:GLN:O	2.18	0.43
1:A:374:ALA:HA	1:A:539:LEU:HD23	2.00	0.43
3:B:603:5NZ:H16	3:B:603:5NZ:H5	1.76	0.43
1:B:376:GLU:O	1:B:380:LEU:HG	2.19	0.43
1:B:81[B]:GLU:CD	1:B:81[B]:GLU:H	2.22	0.43
1:A:213:ILE:O	1:A:219[A]:ARG:HD3	2.17	0.43
1:B:536:LEU:HA	1:B:536:LEU:HD12	1.88	0.43
1:A:369[B]:GLN:HE22	1:A:405:HIS:HE1	1.66	0.42
1:A:376:GLU:O	1:A:380:LEU:HG	2.19	0.42
1:B:210:GLY:HA3	1:B:232:PRO:HD3	2.00	0.42
1:A:249:THR:HG23	1:A:259:PRO:HG3	2.02	0.42
1:A:210:GLY:HA3	1:A:232:PRO:HD3	2.01	0.41
1:B:352:SER:O	1:B:395:ARG:HG3	2.21	0.41
1:A:211:MET:HG3	1:A:232:PRO:HB3	2.03	0.40
1:B:211:MET:HG3	1:B:232:PRO:HB3	2.01	0.40
1:A:200:PHE:HB2	1:A:226:VAL:HB	2.02	0.40
1:A:386:LEU:HD23	1:A:386:LEU:HA	1.96	0.40
1:A:103:THR:HG21	1:A:190:PHE:HB3	2.03	0.40
1:B:227:LEU:HB2	1:B:328:VAL:HG12	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/543 (99%)	521 (97%)	15 (3%)	1 (0%)	52	75
1	B	533/543 (98%)	515 (97%)	17 (3%)	1 (0%)	52	75
All	All	1070/1086 (98%)	1036 (97%)	32 (3%)	2 (0%)	52	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	497	SER
1	A	523	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	444/443 (100%)	432 (97%)	12 (3%)	52	79
1	B	442/443 (100%)	429 (97%)	13 (3%)	50	77
All	All	886/886 (100%)	861 (97%)	25 (3%)	51	78

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	13	ARG
1	A	146	LEU
1	A	161	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	251	LEU
1	A	295	PHE
1	A	322	GLN
1	A	386	LEU
1	A	421	GLN
1	A	493	ARG
1	A	495	SER
1	A	540	LEU
1	B	60	LEU
1	B	84	GLU
1	B	146	LEU
1	B	161	LEU
1	B	200	PHE
1	B	251	LEU
1	B	295	PHE
1	B	313	GLU
1	B	386	LEU
1	B	421	GLN
1	B	437	LEU
1	B	495	SER
1	B	497	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	HIS
1	A	287	HIS
1	B	223	HIS
1	B	322	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	A	601	1	14,14,15	0.36	0	15,19,21	0.71	1 (6%)
2	NAG	A	602	1	14,14,15	0.37	0	15,19,21	0.77	1 (6%)
3	5NZ	A	603	-	54,56,56	2.32	12 (22%)	64,79,79	1.62	10 (15%)
2	NAG	B	601	1	14,14,15	0.37	0	15,19,21	1.12	1 (6%)
4	P6G	B	602	-	18,18,18	0.58	0	17,17,17	0.41	0
3	5NZ	B	603	-	54,56,56	2.47	15 (27%)	64,79,79	1.82	11 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
2	NAG	A	602	1	-	0/6/23/26	0/1/1/1
3	5NZ	A	603	-	-	0/18/25/25	0/8/8/8
2	NAG	B	601	1	-	0/6/23/26	0/1/1/1
4	P6G	B	602	-	-	0/16/16/16	0/0/0/0
3	5NZ	B	603	-	-	0/18/25/25	0/8/8/8

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	603	5NZ	C26-N4	-3.79	1.31	1.37
3	B	603	5NZ	C18-N3	-3.30	1.35	1.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	603	5NZ	C26-N4	-2.67	1.33	1.37
3	B	603	5NZ	C4-C5	2.02	1.51	1.45
3	B	603	5NZ	C6-C7	2.07	1.41	1.36
3	A	603	5NZ	C30-C31	2.85	1.47	1.43
3	A	603	5NZ	N5-N4	2.93	1.40	1.34
3	B	603	5NZ	N5-N4	3.06	1.40	1.34
3	B	603	5NZ	C30-C31	3.13	1.48	1.43
3	A	603	5NZ	N6-N5	3.13	1.39	1.34
3	A	603	5NZ	C11-N3	3.31	1.42	1.35
3	B	603	5NZ	C37-N8	3.40	1.36	1.32
3	B	603	5NZ	C11-C10	3.45	1.49	1.43
3	A	603	5NZ	C37-N8	3.48	1.36	1.32
3	B	603	5NZ	N6-N5	3.50	1.40	1.34
3	B	603	5NZ	C11-N3	3.52	1.43	1.35
3	A	603	5NZ	C11-C10	3.85	1.49	1.43
3	A	603	5NZ	C31-C33	4.67	1.49	1.42
3	B	603	5NZ	C31-C33	5.04	1.50	1.42
3	A	603	5NZ	C10-C4	5.36	1.50	1.42
3	B	603	5NZ	C10-C4	5.50	1.50	1.42
3	B	603	5NZ	C5-C18	5.56	1.50	1.41
3	A	603	5NZ	C5-C18	5.74	1.51	1.41
3	A	603	5NZ	C39-C37	6.98	1.50	1.40
3	B	603	5NZ	C39-C37	7.10	1.50	1.40
3	A	603	5NZ	C30-C39	7.23	1.49	1.39
3	B	603	5NZ	C30-C39	8.67	1.51	1.39

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	603	5NZ	C40-C39-C37	-4.30	117.73	120.95
3	B	603	5NZ	C39-C37-N8	-3.60	120.18	123.30
3	A	603	5NZ	C39-C37-N8	-3.06	120.64	123.30
3	A	603	5NZ	C32-C31-C30	-2.89	120.52	124.92
3	A	603	5NZ	C19-C18-C5	-2.88	116.34	120.69
3	B	603	5NZ	C24-C25-C26	-2.74	105.80	113.89
3	B	603	5NZ	C32-C31-C30	-2.64	120.90	124.92
3	B	603	5NZ	C31-C33-N8	-2.52	120.57	122.89
3	A	603	5NZ	C41-C40-C39	-2.40	108.34	113.14
3	B	603	5NZ	C10-C4-C5	-2.21	117.34	119.94
3	A	603	5NZ	C31-C33-N8	-2.03	121.03	122.89
3	B	603	5NZ	C13-C12-C11	-2.02	116.86	120.20
3	B	603	5NZ	C12-C11-N3	-2.01	118.05	120.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	NAG	C1-O5-C5	2.01	115.09	112.14
3	A	603	5NZ	C28-C29-N7	2.11	117.26	113.09
3	A	603	5NZ	C36-C35-C34	2.15	123.55	120.45
3	A	603	5NZ	C32-C31-C33	2.60	120.88	118.33
2	A	602	NAG	C1-O5-C5	2.61	115.98	112.14
3	B	603	5NZ	C25-C26-N4	2.73	125.78	121.95
2	B	601	NAG	C1-O5-C5	4.07	118.13	112.14
3	A	603	5NZ	C37-N8-C33	5.29	123.65	117.67
3	A	603	5NZ	C39-C30-N7	5.45	127.44	119.53
3	B	603	5NZ	C39-C30-N7	6.15	128.45	119.53
3	B	603	5NZ	C37-N8-C33	6.20	124.69	117.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	603	5NZ	4	0
4	B	602	P6G	1	0
3	B	603	5NZ	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	538/543 (99%)	0.89	61 (11%) <b>7</b> <b>7</b>	32, 47, 76, 132	0
1	B	534/543 (98%)	1.43	134 (25%) <b>1</b> <b>1</b>	33, 52, 83, 141	0
All	All	1072/1086 (98%)	1.16	195 (18%) <b>2</b> <b>2</b>	32, 49, 81, 141	0

All (195) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	341	TYR	8.7
1	B	543	THR	8.4
1	A	341	TYR	8.1
1	B	542	ALA	7.7
1	B	349	ASP	7.5
1	B	75	THR	7.5
1	B	394	LEU	7.4
1	B	391	PRO	7.2
1	B	72	TYR	6.9
1	B	124	TYR	6.7
1	B	365	ILE	6.3
1	B	344	PRO	6.1
1	B	78	PRO	5.8
1	B	337	TYR	5.7
1	B	345	GLY	5.6
1	B	395	ARG	5.4
1	B	353	LEU	5.4
1	A	540	LEU	5.2
1	A	496	LYS	5.1
1	B	342	GLY	4.8
1	A	267	THR	4.7
1	B	385	TRP	4.6
1	A	259	PRO	4.5
1	A	337	TYR	4.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	354	ILE	4.4
1	B	357	ALA	4.3
1	A	542	ALA	4.3
1	A	271	ALA	4.3
1	B	76	LEU	4.2
1	B	105	TYR	4.1
1	B	250	LEU	4.1
1	A	357	ALA	4.0
1	A	315	LEU	4.0
1	B	373	LEU	4.0
1	B	269	LEU	3.9
1	B	364	ARG	3.8
1	A	248	ALA	3.8
1	A	394	LEU	3.7
1	B	12	VAL	3.7
1	B	108	PRO	3.6
1	B	361	ALA	3.6
1	B	541	SER	3.5
1	A	471	ILE	3.5
1	B	358	GLN	3.5
1	B	340	VAL	3.4
1	A	249	THR	3.4
1	B	398	MET	3.4
1	A	541	SER	3.3
1	B	497	SER	3.3
1	A	373	LEU	3.3
1	B	446	PRO	3.3
1	B	347	SER	3.3
1	B	368	PRO	3.3
1	B	420	ALA	3.3
1	B	122	GLY	3.2
1	B	256	GLY	3.2
1	B	539	LEU	3.2
1	B	439	TRP	3.2
1	B	442	TRP	3.1
1	B	447	HIS	3.1
1	B	352	SER	3.1
1	A	472	PHE	3.1
1	B	25	PRO	3.1
1	B	321	PHE	3.1
1	A	253	ARG	3.0
1	B	300	VAL	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	216	LEU	3.0
1	B	191	GLY	3.0
1	B	350	ASN	3.0
1	B	123	PHE	3.0
1	B	77	TYR	3.0
1	A	314	ALA	3.0
1	B	355	SER	2.9
1	A	76	LEU	2.9
1	B	125	SER	2.9
1	A	268	GLU	2.8
1	B	335	GLY	2.8
1	B	20	ILE	2.8
1	B	360	LEU	2.8
1	B	339	LEU	2.8
1	B	338	PHE	2.8
1	B	73	VAL	2.7
1	A	365	ILE	2.7
1	A	165	ARG	2.7
1	B	392	THR	2.7
1	A	312	PRO	2.7
1	A	241	ALA	2.7
1	A	159	LEU	2.7
1	A	342	GLY	2.6
1	B	346	PHE	2.6
1	B	237	ALA	2.6
1	B	318	THR	2.6
1	A	252	ALA	2.6
1	B	79	GLY	2.6
1	B	463	LEU	2.6
1	B	70	TYR	2.6
1	A	72	TYR	2.6
1	B	383	THR	2.6
1	B	203	SER	2.6
1	A	217	PRO	2.6
1	A	422	GLY	2.6
1	B	119	TYR	2.6
1	B	430	PHE	2.5
1	B	438	THR	2.5
1	B	532	TRP	2.5
1	B	535	PHE	2.5
1	A	290	PRO	2.5
1	B	440	PRO	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	288	VAL	2.5
1	A	467	THR	2.5
1	B	380	LEU	2.5
1	A	385	TRP	2.5
1	B	328	VAL	2.5
1	A	82	GLY	2.5
1	B	389	GLU	2.5
1	B	204	ALA	2.4
1	B	436	THR	2.4
1	B	104	PRO	2.4
1	A	161	LEU	2.4
1	B	273	LEU	2.4
1	A	323	ASP	2.4
1	A	226	VAL	2.4
1	B	151	TYR	2.4
1	B	362	GLY	2.4
1	B	280	ASP	2.4
1	B	7	GLN	2.4
1	B	24	ALA	2.4
1	B	208	SER	2.4
1	A	141	VAL	2.4
1	A	535	PHE	2.4
1	B	258	PRO	2.4
1	B	382	TYR	2.4
1	B	336	SER	2.4
1	A	543	THR	2.3
1	B	292	GLU	2.3
1	A	309	SER	2.3
1	A	347	SER	2.3
1	B	284	HIS	2.3
1	B	401	VAL	2.3
1	B	443	MET	2.3
1	B	206	ALA	2.3
1	B	255	VAL	2.3
1	B	402	VAL	2.3
1	B	187	ILE	2.3
1	B	190	PHE	2.3
1	A	377	ALA	2.3
1	A	397	ALA	2.3
1	B	19	GLY	2.3
1	B	298	SER	2.3
1	A	318	THR	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	301	PRO	2.3
1	B	141	VAL	2.3
1	B	87	ASN	2.3
1	A	3	ARG	2.2
1	B	13	ARG	2.2
1	A	458	PRO	2.2
1	B	71	GLN	2.2
1	B	536	LEU	2.2
1	B	412	ALA	2.2
1	B	249	THR	2.2
1	A	171	VAL	2.2
1	A	288	VAL	2.2
1	B	434	ALA	2.2
1	A	137	PHE	2.2
1	B	157	GLY	2.2
1	B	376	GLU	2.2
1	B	156	PHE	2.2
1	B	230	GLY	2.2
1	A	354	ILE	2.1
1	B	451	ILE	2.1
1	A	380	LEU	2.1
1	A	352	SER	2.1
1	A	302	VAL	2.1
1	B	276	ARG	2.1
1	A	360	LEU	2.1
1	B	384	ASP	2.1
1	B	403	GLY	2.1
1	B	10	VAL	2.1
1	B	343	VAL	2.1
1	B	111	PRO	2.1
1	A	251	LEU	2.1
1	B	399	SER	2.1
1	B	120	GLY	2.1
1	A	322	GLN	2.1
1	B	160	ALA	2.1
1	B	277	PRO	2.1
1	B	226	VAL	2.1
1	B	363	VAL	2.1
1	B	515	LEU	2.1
1	A	205	GLY	2.1
1	A	206	ALA	2.0
1	A	506	ALA	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	281	LEU	2.0
1	B	229	SER	2.0
1	B	159	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	P6G	B	602	19/19	0.73	0.38	1.95	62,69,75,76	0
3	5NZ	A	603	49/49	0.78	0.30	0.83	30,45,78,79	0
3	5NZ	B	603	49/49	0.67	0.42	0.80	39,53,94,96	0
2	NAG	A	601	14/15	0.85	0.24	-0.03	78,84,88,88	0
2	NAG	B	601	14/15	0.60	0.42	-0.05	97,102,104,104	0
2	NAG	A	602	14/15	0.72	0.28	-	105,108,114,115	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.