



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:24 AM GMT

PDB ID : 3EIH  
Title : Crystal structure of S.cerevisiae Vps4 in the presence of ATPgammaS  
Authors : Gonciarz, M.D.; Whitby, F.G.; Eckert, D.M.; Kieffer, C.; Heroux, A.; Sundquist, W.I.; Hill, C.P.  
Deposited on : 2008-09-15  
Resolution : 3.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1624 (3.32-3.20)
Clashscore	102246	1806 (3.32-3.20)
Ramachandran outliers	100387	1773 (3.32-3.20)
Sidechain outliers	100360	1771 (3.32-3.20)
RSRZ outliers	91569	1632 (3.32-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	<div> <div>2%</div> <div>53%</div> <div>36%</div> <div>6%</div> </div>
1	B	340	<div> <div>2%</div> <div>50%</div> <div>39%</div> <div>7%</div> </div>
1	C	340	<div> <div>9%</div> <div>45%</div> <div>39%</div> <div>7%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	C	501	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7396 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein sorting-associated protein 4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2469	1552	419	489	9			
1	B	316	Total	C	N	O	S	0	0	0
			2449	1540	415	485	9			
1	C	310	Total	C	N	O	S	0	0	0
			2409	1518	407	475	9			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	98	GLY	-	EXPRESSION TAG	UNP P52917
A	99	ILE	-	EXPRESSION TAG	UNP P52917
A	100	ASP	-	EXPRESSION TAG	UNP P52917
A	101	PRO	-	EXPRESSION TAG	UNP P52917
A	102	PHE	-	EXPRESSION TAG	UNP P52917
A	103	THR	-	EXPRESSION TAG	UNP P52917
A	233	GLN	GLU	ENGINEERED	UNP P52917
B	98	GLY	-	EXPRESSION TAG	UNP P52917
B	99	ILE	-	EXPRESSION TAG	UNP P52917
B	100	ASP	-	EXPRESSION TAG	UNP P52917
B	101	PRO	-	EXPRESSION TAG	UNP P52917
B	102	PHE	-	EXPRESSION TAG	UNP P52917
B	103	THR	-	EXPRESSION TAG	UNP P52917
B	233	GLN	GLU	ENGINEERED	UNP P52917
C	98	GLY	-	EXPRESSION TAG	UNP P52917
C	99	ILE	-	EXPRESSION TAG	UNP P52917
C	100	ASP	-	EXPRESSION TAG	UNP P52917
C	101	PRO	-	EXPRESSION TAG	UNP P52917
C	102	PHE	-	EXPRESSION TAG	UNP P52917
C	103	THR	-	EXPRESSION TAG	UNP P52917
C	233	GLN	GLU	ENGINEERED	UNP P52917

- Molecule 2 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter

code: AGS) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vacuolar protein sorting-associated protein 4



- Molecule 1: Vacuolar protein sorting-associated protein 4



- Molecule 1: Vacuolar protein sorting-associated protein 4





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.96Å 119.80Å 156.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.00 – 3.25 49.93 – 3.25	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.00-3.25) 99.6 (49.93-3.25)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.77 (at 3.25Å)	Xtriage
Refinement program	REFMAC 5.5.0062	Depositor
R, $R_{free}$	0.226 , 0.305 0.242 , 0.287	Depositor DCC
$R_{free}$ test set	1207 reflections (5.42%)	DCC
Wilson B-factor (Å <sup>2</sup> )	79.8	Xtriage
Anisotropy	0.164	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 78.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 23501 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	7396	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AGS, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.64	0/2511	0.84	6/3395 (0.2%)
1	B	0.60	0/2491	0.81	5/3368 (0.1%)
1	C	0.55	0/2450	0.70	0/3312
All	All	0.60	0/7452	0.79	11/10075 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	156	PRO	N-CA-C	13.60	147.47	112.10
1	A	157	HIS	N-CA-C	-9.61	85.06	111.00
1	B	375	PRO	CA-N-CD	-9.44	98.29	111.50
1	A	319	LEU	CA-CB-CG	6.63	130.54	115.30
1	B	216	GLN	N-CA-C	6.25	127.87	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	157	HIS	Peptide
1	A	357	ALA	Peptide
1	B	216	GLN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2469	0	2476	122	0
1	B	2449	0	2457	145	0
1	C	2409	0	2417	216	0
2	A	31	0	12	1	0
2	B	31	0	12	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	C	4	0	6	10	0
All	All	7396	0	7380	476	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 476 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:154:LYS:HD3	1:C:155:PHE:CE1	1.61	1.34
1:B:386:MET:CE	1:B:391:ILE:HG13	1.63	1.27
1:B:386:MET:HE1	1:B:391:ILE:CG1	1.67	1.24
1:B:374:THR:HG23	1:B:375:PRO:CD	1.68	1.24
1:C:154:LYS:HE3	1:C:155:PHE:CZ	1.75	1.21

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/340 (93%)	267 (84%)	40 (13%)	10 (3%)	5	33
1	B	314/340 (92%)	252 (80%)	47 (15%)	15 (5%)	3	21
1	C	306/340 (90%)	227 (74%)	59 (19%)	20 (6%)	1	13
All	All	937/1020 (92%)	746 (80%)	146 (16%)	45 (5%)	3	21

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	266	ASP
1	B	191	ASN
1	B	210	SER
1	B	291	GLU
1	B	366	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/287 (95%)	262 (96%)	10 (4%)	41	77
1	B	270/287 (94%)	260 (96%)	10 (4%)	41	77
1	C	266/287 (93%)	246 (92%)	20 (8%)	17	53
All	All	808/861 (94%)	768 (95%)	40 (5%)	30	70

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	390	ASP
1	C	138	LEU
1	C	365	THR
1	C	130	VAL
1	C	158	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	261	ASN
1	C	425	GLN
1	C	268	GLN
1	B	261	ASN
1	C	265	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	AGS	A	501	-	24,33,33	2.02	3 (12%)	28,52,52	1.94	5 (17%)
2	AGS	B	501	-	24,33,33	2.19	3 (12%)	28,52,52	2.45	5 (17%)
4	EDO	C	501	-	3,3,3	0.50	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AGS	A	501	-	-	0/15/38/38	0/3/3/3
2	AGS	B	501	-	-	0/15/38/38	0/3/3/3
4	EDO	C	501	-	-	0/1/1/1	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	AGS	O4'-C1'	2.15	1.43	1.41
2	B	501	AGS	O4'-C1'	2.19	1.44	1.41
2	A	501	AGS	C5-C4	3.24	1.47	1.40
2	B	501	AGS	C5-C4	3.40	1.48	1.40
2	A	501	AGS	PG-S1G	8.39	2.06	1.90

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	AGS	N3-C2-N1	-8.03	122.75	128.89
2	B	501	AGS	C2'-C1'-N9	-6.77	103.95	114.29
2	A	501	AGS	N3-C2-N1	-6.48	123.93	128.89
2	A	501	AGS	PB-O3B-PG	-4.71	116.89	132.67
2	B	501	AGS	PB-O3B-PG	-4.50	117.57	132.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	AGS	1	0
2	B	501	AGS	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	501	EDO	10	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/340 (93%)	0.26	7 (2%) 65 55	42, 68, 108, 124	0
1	B	316/340 (92%)	0.31	8 (2%) 61 51	51, 82, 122, 138	0
1	C	310/340 (91%)	0.62	29 (9%) 11 8	74, 105, 130, 136	0
All	All	945/1020 (92%)	0.40	44 (4%) 35 27	42, 84, 126, 138	0

The worst 5 of 44 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	243	GLU	5.7
1	B	240	THR	5.3
1	C	181	TYR	5.0
1	A	242	GLY	4.9
1	B	242	GLY	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	AGS	B	501	31/31	0.91	0.16	-1.20	103,107,113,114	0
2	AGS	A	501	31/31	0.94	0.16	-1.80	64,67,81,82	0
4	EDO	C	501	4/4	0.95	0.08	-2.40	54,56,58,61	0
3	MG	C	502	1/1	0.84	0.12	-	100,100,100,100	0
3	MG	B	502	1/1	0.96	0.14	-	59,59,59,59	0
3	MG	A	502	1/1	0.94	0.18	-	51,51,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.